



## Genetics/Génétique

Cytogenetic data on the threatened leafcutter ant *Atta robusta* Borgmeier, 1939 (Formicidae: Myrmicinae: Attini)<sup>☆</sup>

*Données cytogénétiques sur la fourmi coupeuse de feuilles menacée d'extinction Atta robusta Borgmeier, 1939 (Formicidae : Myrmicinae : Attini)*

Luísa Antônia Campos Barros<sup>a,\*b</sup>, Hilton Jeferson Alves Cardoso de Aguiar<sup>b,c</sup>, Gisele Amaro Teixeira<sup>a</sup>, Cléa dos Santos Ferreira Mariano<sup>d,e</sup>, Marcos da Cunha Teixeira<sup>g</sup>, Jacques Hubert Charles Delabie<sup>d,f</sup>, Silvia das Graças Pompolo<sup>a</sup>

<sup>a</sup> Laboratório de Citogenética de Insetos, Departamento de Biologia Geral, Universidade Federal de Viçosa, Viçosa, MG 36570-000, Brazil

<sup>b</sup> Programa de Pós-graduação em Genética e Melhoramento, Universidade Federal de Viçosa, Viçosa, MG 36570-000, Brazil

<sup>c</sup> Universidade Federal do Amapá, Campus Binacional, Oiapoque, AP 68980-000, Brazil

<sup>d</sup> Laboratório de Mirmecologia, CEPEC/CEPLAC, Itabuna, BA CP 7, 45600-000, Brazil

<sup>e</sup> Departamento de Ciências Biológicas, Universidade Estadual de Santa Cruz, Ilhéus, BA 45650-000, Brazil

<sup>f</sup> Departamento de Ciências Agrárias e Ambientais, Universidade Estadual de Santa Cruz, Ilhéus, BA 45650-000, Brazil

<sup>g</sup> Departamento de Ciências Agrárias e Ambientais, Universidade Federal do Espírito Santo, ES, Brazil

## ARTICLE INFO

## Article history:

Received 12 April 2015

Accepted after revision 2 July 2015

Available online 24 August 2015

## Keywords:

Banding techniques

Conservation

FISH

Karyotype

Restinga

## ABSTRACT

The karyotype of the threatened ant species *Atta robusta* is described so as to establish the evolutionary relationships of this taxon with other leafcutter ants. Standard Giemsa staining, C-banding, NOR banding, fluorochromes CMA<sub>3</sub>/DAPI, Hsc-FA technique and Fluorescence *in situ* Hybridization (FISH) using 18S rDNA probe were conducted on a population from Aracruz, state of Espírito Santo, Brazil, allowing for comparisons with data available on *Atta* and other fungus-growing ant species. The diploid chromosome number observed for *A. robusta* was  $2n = 22$ , and the karyotypic formula was  $18m + 2sm + 2st$ . Heterochromatic blocks were observed in the centromeric region of most chromosomes, where one pair of metacentric chromosomes is characterized by a GC-rich heterochromatic band in the interstitial region of its long arm. The detection of 18S rDNA using FISH confirmed the presence of single NOR for *A. robusta*. This is the first report of rDNA 18S detection using FISH for leafcutter ants. The cytogenetic results of this study confirm the information available for *Atta* and allow us to confirm the conserved chromosome number, morphology and banding pattern within the genus for the taxa studied to date, which included species from three out of the four groups of *Atta* indicated by molecular data. The accumulation of cytogenetic data on fungus-growing ants enhances the understanding of the genomic evolutionary patterns of *Atta*, since it belongs to a group of recent origin between the most well studied ants. Cytogenetic data does not indicate restrictions in relocation or reintroduction in areas where populations were extinct due to the conserved

<sup>☆</sup> Barros et al.: Cytogenetic data of *Atta robusta*.

\* Corresponding author.

E-mail address: [luufv@yahoo.com.br](mailto:luufv@yahoo.com.br) (L.A.C. Barros).

karyotype. This study allows for cytogenetic comparison of *A. robusta* with other ants of *Atta*, emphasizing the importance of chromosomal information for species conservation.  
© 2015 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

## RÉSUMÉ

Mots clés:  
Techniques de marcation  
Conservation  
FISH  
Caryotype  
Littoral.

Le caryotype de la fourmi *Atta robusta*, menacée d'extinction, a été décrit dans le but d'établir les relations évolutives de ce taxon avec les autres fourmis coupeuses de feuilles. La coloration au Giemsa, bande C, bande NOR, fluorochromes CMA<sub>3</sub>/DAPI, la technique Hsc-FA et l'hybridation fluorescente *in situ* (FISH) utilisant une sonde de l'ADN ribosomique 18S ont été mises en œuvre sur une population d'Aracruz, État du Espírito Santo, au Brésil, permettant des comparaisons avec les données disponibles sur *Atta* et d'autres espèces de fourmis cultivatrices de champignons. Le nombre de chromosomes diploïdes observés pour *A. robusta* était  $2n=22$  et la formule chromosomique  $18m + 2sm + 2st$ . Des blocs hétérochromatiques ont été rencontrés dans la région centromérique de la plupart des chromosomes, où une paire de chromosomes métacentriques se caractérise par une bande hétérochromatique riche en GC dans la région interstitielle de son bras long. La détection d'ADNr 18S utilisant la technique du FISH a confirmé la présence d'une unique paire de NOR pour *A. robusta*. Les résultats cytogénétiques de cette étude confirment l'information disponible pour *Atta* ainsi que la conservation du nombre de chromosomes, la morphologie et le modèle de bande dans le genre pour les taxons déjà étudiés jusqu'à présent, et qui concerne des espèces de trois des quatre groupes d'*Atta* suggérés par les données moléculaires. En raison de ce caryotype conservé, les données cytogénétiques ne présentent aucune restriction quant à la relocation ou la réintroduction de l'espèce dans les zones où les populations ont été éteintes. L'accumulation de données cytogénétiques sur les espèces de fourmis cultivatrices de champignons contribue ainsi à la compréhension des tendances évolutives génomiques dans le genre *Atta*, car il appartient à un groupe d'origine récente parmi toutes les fourmis les plus étudiées. Cette étude a permis la comparaison cytogénétique de *A. robusta* avec d'autres fourmis du genre *Atta*, soulignant l'importance de l'information chromosomique pour la conservation des espèces.

© 2015 Académie des sciences. Publié par Elsevier Masson SAS. Tous droits réservés.

## 1. Introduction

Species belonging to the fungus-growing ants comprise the erstwhile tribe Attini, and were included recently in the clade Attini that corresponds to a wide group constituting a diverse selection of predominantly Neotropical taxa [1].

The leafcutter ants represented by the genera *Atta* and *Acromyrmex* are considered the dominant herbivores of the Neotropics [2,3]. These ants in particular are the most well-known and studied species of the genus *Atta* because they significantly impact human activity, because they cut plant material from cultivated plants in large quantities for the cultivation of the symbiotic fungus on which they feed [2,4]. However, in natural ecosystems, the leafcutter ants are especially important for ecosystem structure and dynamics [5].

*Atta robusta* Borgmeier, 1939, an endemic leafcutter ant of the “restinga” ecosystem (sandy open vegetation along the Brazilian coastline), occurs from northern regions of the state of Espírito Santo to the south of the state of Rio de Janeiro, Brazil [6]. It is even one of the few ants of this genus endemic to the restricted area [7]. The species was included in the list of threatened fauna, mainly because of the intense human occupation of the ecosystem in its region of occurrence, as well as because of the indiscriminate chemical use for control of the leafcutter ants [5]. Furthermore, it is possible that plant cover removal

from the restinga directly influences the temperature and humidity control necessary for cultivation of the fungus on which they feed [5]. *A. robusta* plays a significant role in vegetation regeneration because it interacts with at least 36 plant species from the restinga of Guriri Island, state of Espírito Santo, throughout the year (reviewed in [5]). Because it does not exert any kind of economic impact and cannot be considered a pest, most of the studies conducted on this species are recent [5,6,8,9].

Of the various fields of genetics, cytogenetics is considered an important tool because chromosomal rearrangements may induce developmental changes in individuals without causing genomic alterations [10], thus directly participating in the evolutionary mechanisms generating variability and modifications in individual fitness [11,12]. Cytogenetic studies conducted on the family Formicidae highlight the importance of this field of genetics in evolutionary, phylogenetic and taxonomic studies of this diverse group [reviewed in 13]. Cytogenetic data on threatened species reveals the relevance of the referred tool for a better understanding of the structure and isolation of distinct populations, as well as in furnishing arguments for the species conservation programs [14,15]. For example, cytogenetic studies were performed on the threatened ant *D. lucida* at 15 sites of Bahia and Espírito Santo, revealing the occurrence of a great variation in the chromosome number:  $2n = 106, 114,$

116, 118 and 120 in the state of Bahia and  $2n = 118$  in Espírito Santo. In the southern Bahia forest, the fragments are smaller and much more isolated due to human occupation and deforestation, which increase the chance of the evolutionary divergence of karyotypes among the different populations. Cytogenetic results determined from a range of locations indicated the importance of this tool for conservation of this threatened species [15].

To date, chromosomal studies have been conducted in five taxa of leafcutter ants of the genus *Atta*. All presented  $2n = 22$  chromosomes, including the species collected in Brazil: *A. bispshaerica*, *A. laevigata*, *A. sexdens rubropilosa* [16,17], *A. sexdens piriventris* [18] and in Panama: *A. colombica* [19]. Data on chromosome banding are available for *A. bispshaerica*, *A. laevigata*, *A. sexdens rubropilosa* [17], and *A. colombica* [19]. Chromosome banding techniques are of fundamental importance for the localization of specific chromosomal regions, allowing for the performance of additional evolutionary inferences not permitted with conventional Giemsa staining alone. Cytogenetic data shows constant karyotypes for the different *Atta* spp. not only with respect to chromosome number, but also their morphology, pattern and heterochromatin composition [17].

The species of the genus *Atta* cytogenetically studied to date include ants presenting broad geographic distribution. No endemic ant species of this genus with restricted distribution has been studied at the cytogenetic level, which allows for comparative studies on the evolutionary dynamics of the group. Incorporating the cytogenetic results with those of other current studies on this ant (Teixeira MC, personal communication) will enable a better understanding of this group of leafcutter ants, and also assist in conservation strategies.

Knowledge on the degree of genetic variability within natural populations of threatened species can usually furnish arguments on structuring conservationist practices and politics, thus avoiding the disaster of species extinction. Habitat fragmentation and other anthropic interventions reduce the gene flow as well as genetic diversity [20], affecting the chances of survival of species, like *A. robusta*.

The present study sought to describe the karyotype of the threatened leafcutter ant *A. robusta*, endemic to restinga areas, in order to provide cytogenetic knowledge for its conservation and also to provide a better understanding of the evolution of the leafcutter ants.

## 2. Materials and methods

Two colonies of *A. robusta* were collected in the city of Aracruz, located in the Brazilian state of Espírito Santo ( $19^{\circ}51'S$   $40^{\circ}04'W$ ), during January 2012. Specimen collection was authorized by the Brazilian Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio), according to a special collecting permit (SISBio number 31861). This collecting permit was issued to Luísa Antônia Campos Barros. Ant vouchers (workers) were deposited in the reference collection of the Laboratório de Mirmecologia, Centro de Pesquisas do Cacau (CPDC/Brazil), under the record #5727.


Mitotic metaphases were obtained according to Imai et al. [21] by dissecting the cerebral ganglia of the larvae after the meconium elimination. A total of 50 individuals were analyzed. The metaphases were observed and photographed using a BX 60 microscope with a  $100\times$  objective, coupled with a Q-Color3 Olympus® image capture system. A total of 15 metaphases with similar degrees of condensation, without overlapping and evident centromeres, were organized by pairing the chromosomes in order of size, measuring and classifying them according to Levan et al. [22], based on the chromosome arm ratio ( $r$ ) using the following features: long arm length ( $L$ ), short arm length ( $S$ ) and arm ratio between the long and short arms ( $r = L/S$ ). The chromosomes were classified as:  $m$  = metacentric,  $sm$  = submetacentric,  $st$  = subtelocentric and  $a$  = acrocentric. Chromosomes were measured and organized using Image Pro Plus® and Corel Photopaint X3 software packages, respectively. The C-banding technique was performed according to Sumner [23]. Regions rich in GC and AT base pairs were detected using the fluorochromes Chromomycin A<sub>3</sub> (CMA<sub>3</sub>) and 4',6-diamidin-2-phenylindole (DAPI), respectively [24]. The Hsc-FA technique (secondary constriction heterochromatin-associated bands by fluorescence using Acridine Orange) was applied according to [25]. The NOR banding technique was performed according to Howell and Black [26]. Ribosomal gene clusters were detected by FISH according to Pinkel et al. [27], with the use of a 18S rDNA probe isolated from the bee *Melipona quinquefasciata* amplified by PCR using the primers 18SF1 (5'-GTCATATGCTGTCAAAGA-3') and 18SR1.1 (3'-TCTAATTTCAAAGTAAACGC-5') designed by Pereira [28]. The probes 18S rDNA were labeled by indirect method using digoxigenin-11-dUTP (Roche Applied Science) and the signal was detected with anti-digoxigenin-rhodamine (Roche Applied Science). The metaphases were analyzed with an epifluorescence microscope using WU (330–385 nm) and WG filters (510–550 nm), for DAPI and rhodamine, respectively.

## 3. Results

The chromosome number observed was  $2n = 22$  with the karyotypic formula:  $18m + 2sm + 2st$  (Fig. 1a). A secondary constriction was observed on the fourth metacentric pair (Fig. 1a). The centromeric regions of the chromosomes indicated the presence of weak and little visible blocks of heterochromatin (Fig. 2). The long arm of the fourth pair of metacentric chromosomes showed a region rich in GC base pairs using CMA<sub>3</sub> (Fig. 3a). Regions with differential staining using DAPI were absent, with a uniform pattern along the chromosomes (Fig. 3b). Markings resembling CMA<sub>3</sub> pattern were also observed on the fourth pair of chromosomes when the Hsc-FA technique was applied using Acridine Orange (AO) (Fig. 3c), NOR banding technique (Fig. 3a – inbox) and FISH using 18S rDNA (Fig. 3d).

## 4. Discussion

The chromosome number of *A. robusta* showed the same general pattern described for other *Atta* spp.:  $2n = 22$ ,



**Fig. 1.** Conventional staining with Giemsa of *Atta robusta*. a: diploid karyotype  $2n = 22$ ; b: the respective metaphase. Arrowhead indicates the presence of a secondary constriction in the fourth metacentric pair. m = metacentric, sm = submetacentric, st = subtelocentric. Bar = 5  $\mu\text{m}$ .

with a predominance of metacentric chromosomes. The referred species also presented a conserved morphology of chromosomes when compared with the other species of the genus [17], which was based on the chromosome arm ratio according to Levan et al. [22]. Cytogenetic results of this study confirm the information available for *Atta* and allow for confirming the conserved chromosome number, morphology and banding pattern within the genus for taxa studied to date, which included species from three of the four groups of *Atta* indicated by molecular data [29]. It included species in three of the four groups of the genus *Atta* pertaining to *A. laevigata* and *A. bisphaerica* in the Epiatta group; *A. colombica* in the *Atta* sensu stricto group; and *A. sexdens rubropilosa*, *A. sexdens piriventris*, and *A. robusta* in the Neoatta group.

The morphological constancy of the chromosomes revealed for *Atta* spp. differs from that observed for species of the other leafcutter ant genus *Acromyrmex*, which presented  $2n = 38$  chromosomes [Barros et al. in prep.]. Morphological differences in the chromosomes may be a consequence of the greater chromosomal number of *Acromyrmex* spp., which, according to the most robust chromosomal evolution theories of Formicidae proposed by Imai et al. [21], would result from centric fissions that allow for variations in the heterochromatin blocks between different species. The karyotype of *A. striatus*, however, bears closer similarity to *Atta* spp. ( $2n = 22$ ) studied earlier and, associated with the molecular data, it provided new insights into the phylogenetic position of

this species among leafcutter ants [30]. According to the authors, it is a sister group of the remainder leafcutter ants that split before the divergence between *Acromyrmex*.


The heterochromatin distribution pattern of *A. robusta* is similar to those of other *Atta* spp., weak centromeric markings, differing from the pattern observed for *A. striatus* ( $2n = 22$ ), which is the only *Acromyrmex* having the same chromosome number as *Atta* spp. In this ant, cytogenetic studies not only presented weak centromeric markings, but also more evident additional pericentromeric, interstitial, and telomeric heterochromatic GC-rich blocks [27]. Markings observed using the Hsc-FA technique was similar to those observed using the fluorochrome CMA<sub>3</sub> in the present study. Hsc-FA using (AO) marks the NOR-associated heterochromatin (for details, see [31]). The same pattern was also observed for the ants *Dinoponera lucida* [31] and *Wasmannia auropunctata* [32]. The only pair of *A. robusta* chromosomes that showed heterochromatic blocks rich in GC base pairs using the fluorochromes CMA<sub>3</sub> and AO has a pattern similar to that observed in other *Atta* spp. The negative interstitial region of the fourth chromosome pair was observed with DAPI, indicating complementarity with CMA<sub>3</sub>. The same non-specific pattern observed with DAPI along the chromosomes was observed for other fungus-growing ants, e.g., *Mycoceropurus goeldii* [33], *A. striatus* [30], *Trachymyrmex fuscus* [34] and *Atta* spp. [17].

The markings obtained with the FISH technique confirmed the results obtained when employing the fluorochromes CMA<sub>3</sub> and AO, as well as the NOR banding, complementing the data and allowing us to conclude that there is a single NOR for *A. robusta*. Studies conducted with ants that presented markings in one pair of chromosomes coincided with single NOR e.g., *D. lucida* [15,31]. It is possible that a single NOR will be found in the genus *Atta*. Only *M. goeldii* among fungus-growing ants had its 18S rDNA sites mapped [35], and this study presents the first data for leafcutter ants.

Multiple markings rich in GC base pairs were described in *A. striatus* [30]. However, they were absent in *A. bisphaerica*, *A. laevigata* and *A. sexdens rubropilosa* [17] and also in the chromosomes of *A. robusta* stained with CMA<sub>3</sub>. A single NOR was confirmed for *A. robusta*; however, the characteristics of the heterochromatin blocks rich in GC base pairs remain unknown for *A. striatus* and only further molecular cytogenetic techniques (FISH) could confirm the presence or absence of rDNA in the chromosomes. The fungus-growing ant *M. goeldii* showed single NOR [35], although it presented multiple GC-rich regions [33].



**Fig. 2.** Metaphase of *Atta robusta* submitted to the C-banding protocol. Centromeric markings denote weak heterochromatin. Bar = 5  $\mu\text{m}$ .



**Fig. 3.** (Color online.) Cytogenetic markings in the fourth chromosome pair of *Atta robusta*. Metaphase stained with fluorochromes (a) CMA<sub>3</sub> that match GC-rich regions indicated by the arrows and (b) DAPI with uniform pattern of AT base pairs along the chromosomes. NOR banding (b, inbox), indicates the nucleolus organizer regions bearer pair. Hsc-FA using Acridine Orange marks the NOR-associated heterochromatin indicated by the arrows (c) and the FISH technique to detect ribosomal genes 18S (d). Bar = 5  $\mu$ m.

At the long arm of the fourth chromosome pair markings were observed in its interstitial region using the NOR banding technique, indicating the presence of active ribosomal genes in the last interphase in this pair. The repeatability obtained in the metaphases subjected to this technique is low. However, the clear secondary constriction associated with the presence of heterochromatic blocks rich in GC base pairs, using the CMA<sub>3</sub> and AO, in addition to 18S rDNA-FISH, confirm that this chromosome pair is the NOR bearer. Imai et al. [36] raised the possibility that, somehow, the chromatin of the ribosomal genes was little accessible, which hinders the argentophilic proteins from associating with the chromatin and that silver nitrate later binds to them.

The endemic status of *A. robusta* suggests peculiarities in its biology if compared to other leafcutter ants. Moreover, strong anthropic pressure on the biome of this species may greatly reduce the variability of its populations by subjecting them to a bottleneck effect capable of reducing its frequency of occurrence. This effect could be expected in function of its cytogenetic configuration; however, the results of this study, which represent the first cytogenetic information on an endemic species of *Atta* with a restricted

distribution, showed no differences in the karyotype compared to other *Atta* spp. Although samples were not collected in other populations of *A. robusta*, the constancy of the chromosomal characters found in *Atta* suggests that variations in the karyotype of *A. robusta* are unlikely.

Populations of *A. robusta* have been management target for conservation purposes since 2008 due to the intense process of plant cover reduction in restinga areas. The use of cytogenetic data in animal relocation and breeding programs is of great significance to prevent the reduction of fitness or infertility, because chromosomally divergent groups need to be avoided [14,37]. Therefore, it is important to ensure, especially when dealing with threatened species, that the animals used as founders do not carry chromosomal rearrangements [28]. Considering the constant karyotype (number, morphology and banding patterns) of the threatened *A. robusta* and other *Atta* spp., cytogenetic data do not indicate restrictions in relocation or reintroduction in areas where populations were extinct due to the conserved karyotype. However, other data need to be considered, including molecular biology that indicates haplogroups for the species [13,17]. It differs from the giant ant *Dinoponera lucida* that presents high chromosome number variation in

some populations of the state of Bahia (Brazil). In this case, cytogenetic data need to be considered for relocation or reintroduction in areas where the populations were extinct in association with other fields of knowledge. Considering the status of the threatened *A. robusta*, all information is important for conservation and chromosome data is not a restriction. Furthermore, efforts for preserving restinga environments are needed to preserve *A. robusta* populations along with many other organisms.

## Disclosure of interest

The authors declare that they have no conflicts of interest concerning this article.

## Acknowledgments

We are grateful to Filipe Pola Vargas and José Teixeira for field support at Aracruz, Riudo de Paiva Ferreira for technical support. Access to the biological material was allowed by the Brazilian Instituto Chico Mendes de Conservação da Biodiversidade (permit SISBio number 31861). LACB and JHCD acknowledge the grants from Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq). This research was supported by Fundação de Amparo à Pesquisa de Minas Gerais (FAPE-MIG) process: APQ-00141-2012 and Fundação de Amparo à Pesquisa da Bahia (FAPESB).

## References

- [1] P.S. Ward, S.G. Brady, B.L. Fisher, T.R. Schultz, The evolution of myrmicine ants: phylogeny and biogeography of a hyperdiverse ant clade (Hymenoptera: Formicidae), *Syst. Entomol.* 40 (2015) 61–81.
- [2] N.A. Weber, Gardening ants, the Attines, *Mem. Amer. Philos. Soc.* 92 (1972) 1–146.
- [3] B. Hölldobler, E.O. Wilson, *The Ants*, Harvard University Press, USA, 1990.
- [4] N.J. Mehdiabadi, T.R. Schultz, Natural history and phylogeny of the fungus-farming ants (Hymenoptera: Formicidae: Myrmicinae: Attini), *Myrmecol. News* 13 (2009) 37–55.
- [5] M.C. Teixeira, I.A. Santos, J.H. Schoereder, *Atta robusta*: endemismo, extinção ou ausência de estudos? in: E.F. Vilela, I.A. Santos, J.H. Schoereder, J. Lino-Neto, L.A.O. Campos, J.E. Serrão (Eds.), *Insetos Sociais: da biologia à aplicação*, Editora da Universidade Federal de Viçosa, Viçosa-MG, 2008, pp. 359–367.
- [6] M.C. Teixeira, J.H. Schoereder, A.J. Mayhé-Nunes, Geographic distribution of *Atta robusta* Borgmeier (Hymenoptera: Formicidae), *Neotrop. Entomol.* 32 (2003) 719–721.
- [7] J.H.C. Delabie, H.S.R. Alves, G.M.R. Strenzel, A.F.R. Carmo, I.C. Nascimento, in: T.M.C. Della-Lucia (Ed.), *Distribuição das formigas cortadeiras dos gêneros Acromyrmex e Atta no Novo Mundo. Formigas cortadeiras: da bioecologia ao manejo*, 1<sup>a</sup> edição, Editora da Universidade Federal de Viçosa, Viçosa-MG, 2011, pp. 80–101.
- [8] D.E.E. Pinto, Diversidade genética, relações filogenéticas e morfologia das sensilas antenais da formiga ameaçada de extinção *Atta robusta* Borgmeier, 1939 (Hymenoptera, Formicidae), (D. Sc. Thesis), in: Programa de Pós-graduação em Genética e Melhoramento, Universidade Federal de Viçosa, Viçosa-MG, Brazil, 2012.
- [9] E.P. Reis, T.M.F. Salomão, L.A.O. Campos, M.G. Tavares, Genetic diversity and structure of *Atta robusta* (Hymenoptera, Formicidae, Attini), an endangered species endemic to the restinga ecoregion, *Genet. Mol. Biol.* 37 (2014) 581–586.
- [10] G.L. Bush, Modes of speciation, *Annu. Rev. Ecol. Syst.* 6 (1975) 339–364.
- [11] M.J.D. White, Cytogenetics of speciation, *Aust. J. Entomol.* 9 (1970) 1–6.
- [12] A.A. Hoffmann, L.H. Rieseberg, Revisiting the impact of inversions in evolution: from population genetic markers to drivers of adaptive shifts and speciation? *Annu. Rev. Ecol. Evol. Syst.* 39 (2008) 21–42.
- [13] P. Lorite, T. Palomeque, Karyotype evolution in ants (Hymenoptera: Formicidae), with a review of the known ant chromosome numbers, *Myrmecol. News* 13 (2010) 89–102.
- [14] T.J. Robinson, F.F.B. Elder, Cytogenetics: its role in wildlife management and the genetic conservation of mammals, *Biol. Conserv.* 63 (1993) 47–51.
- [15] C.S.F. Mariano, S.G. Pompolo, L.A.C. Barros, E. Mariano-Neto, S. Campiolo, J.H.C. Delabie, A biogeographical study of the threatened ant *Dinoponera lucida* Emery (Hymenoptera: Formicidae: Ponerinae) using a cytogenetic approach, *Insect Conserv. Divers.* 1 (2008) 161–168.
- [16] M.A.M. Fadini, S.G. Pompolo, Cytogenetics of some ant species of the tribe Attini (Hymenoptera, Formicidae) from the region of Viçosa, MG, *Rev. Bras. Genet.* 19 (1996) 53–55.
- [17] L.A.C. Barros, G.A. Teixeira, H.J.A.C. Aguiar, C.S.F. Mariano, J.H.C. Delabie, S.G. Pompolo, Banding patterns of three leafcutter ant species of the genus *Atta* (Formicidae: Myrmicinae) and chromosomal inferences, *Fla. Entomol.* 97 (2014) 1694–1701.
- [18] M.C. Santos-Colares, J. Viégas, M.G. Martino Roth, A.E. Loeck, Preparation of mitotic chromosomes of leaf-cutting ants from the genera *Atta* and *Acromyrmex*, *Braz. J. Genet.* 20 (1997) 25–27.
- [19] T. Murakami, A. Fujiwara, M.C. Yoshida, Cytogenetics of ten ant species of the tribe Attini (Hymenoptera, Formicidae) in Barro Colorado Island, Panama, *Chromosome Sci.* 2 (1998) 135–139.
- [20] N.S. Sudhi, P.R. Ehrlich, *Conservation biology for all*, Oxford University Press, Oxford, United Kingdom, 2010.
- [21] H. Imai, R.W. Taylor, M.W. Croziers, Modes of spontaneous chromosomal mutation and karyotype evolution in ants with reference to the minimum interaction hypothesis, *Jpn. J. Genet.* 63 (1988) 159–185.
- [22] A. Levan, K. Fredga, A. Sandberg, Nomenclature for centromeric position on chromosomes, *Hereditas* 52 (1964) 201–220.
- [23] A.T. Sumner, A simple technique for demonstrating centromeric heterochromatin, *Exp. Cell Res.* 83 (1972) 438–442.
- [24] D. Schweizer, Simultaneous fluorescent staining of R bands and specific heterochromatic regions (DA/DAPI-bands) in human chromosomes, *Cytogenet. Cell Genet.* 27 (1980) 190–193.
- [25] R.S. Verma, A. Babu, *Human chromosomes. Principles and Techniques*, McGraw-Hill, Inc., New York, 1995.
- [26] W.M. Howell, D.A. Black, Controlled silver-staining of nucleolus organizer regions with a protective colloidal developer: a 1-step method, *Experientia* 36 (1980) 1014–1015.
- [27] D. Pinkel, T. Straume, J.W. Gray, Cytogenetic analysis using quantitative, high-sensitivity, fluorescence hybridization, *Proc. Natl. Acad. Sci. USA* 83 (1986) 2934–2938.
- [28] J.O.P. Pereira, Diversidade genética da abelha sem ferrão *Melipona quinquefasciata* baseada no seqüenciamento das regiões ITS1 parcial e 18S do DNA ribossômico nuclear, (D. Sc. Thesis), Programa de Pós-graduação em Zootecnia, Universidade Federal do Ceará, Fortaleza-CE, Brazil, 2006.
- [29] M. Bacci Jr., S.E. Solomon, U.G. Mueller, V.G. Martins, A.O. Carvalho, L.G. Vieira, A.C.O. Silva-Pinhati, Phylogeny of leafcutter ants in the genus *Atta fabricius* (Formicidae: Attini) based on mitochondrial and nuclear DNA sequences, *Mol. Phylogenet. Evol.* 5 (2009) 427–437.
- [30] M.P. Cristiano, D.C. Cardoso, T.M. Fernandes-Salomão, Cytogenetic and molecular analyses reveal a divergence between *Acromyrmex striatus* (Roger, 1863) and other congeneric species: taxonomic implications, *PLoS ONE* 8 (2013) e59784.
- [31] L.A.C. Barros, C.S.F. Mariano, S.G. Pompolo, J.H.C. Delabie, Hsc-FA and NOR bandings on chromosomes of the giant ant *Dinoponera lucida* Emery, 1901 (Hymenoptera: Formicidae), *Comp. Cytogenet.* 3 (2009) 97–102.
- [32] A.L.B. Souza, C.S.F. Mariano, J.H.C. Delabie, S.G. Pompolo, J.E. Serrão, Cytogenetic studies on workers of the Neotropical ant *Wasmannia auropunctata* (Roger, 1863) (Hymenoptera: Formicidae: Myrmicinae), *Ann. Soc. Entomol. Fr.* 47 (2011) 507–513.
- [33] L.A.C. Barros, H.J.A.C. Aguiar, C.S.F. Mariano, J.H.C.D. Delabie, S.G. Pompolo, Cytogenetic characterization of the lower-Attine *Mycocepurus goeldii* (Formicidae: Myrmicinae: Attini), *Sociobiology* 56 (2010) 57–66.
- [34] L.A.C. Barros, H.J.A.C. Aguiar, C.S.F. Mariano, J.H.C. Delabie, S.G. Pompolo, Cytogenetic characterization of the ant *Trachymyrmex fuscus* Emery, 1934 (Formicidae: Myrmicinae: Attini) with the description of a chromosomal polymorphism, *Ann. Soc. Entomol. Fr.* 50 (2014) 1–7.
- [35] L.A.C. Barros, H.J.A.C. Aguiar, V. Andrade-Souza, C.S.F. Mariano, J.H.C. Delabie, S.G. Pompolo, Occurrence of pre-nucleolar bodies and 45S rRNA location on the chromosomes of the ant *Mycocepurus goeldii* (Forel) (Formicidae, Myrmicinae, Attini), *Hereditas* 149 (2012) 50–54.
- [36] H.T. Imai, H. Hirai, Y. Satta, T. Shiroischi, M. Yamada, R.W. Taylor, Phase specific Ag-staining of nucleolar organizer regions (NORs) and kinetochores in the Australian and *Myrmecia crozlandi*, *Jap. J. Genet.* 67 (1992) 437–447.
- [37] K. Benirschke, A.T. Kumamoto, Mammalian cytogenetics and conservation of species, *J. Hered.* 82 (1991) 187–191.