***Harpactor angulosus* (Reduviidae: Harpactorinae),
A Predator of Neotropical Saturniids, *Hylesia* spp. in
Brazil**

Author(s): Alexandre I. A. Pereira, José C. Zanuncio, Hélcio R. Gil-Santana, Francisco S. Ramalho, Germano L. D. Leite, and José E. Serrão

Source: Entomological News, 120(2):206-212.

Published By: The American Entomological Society

<https://doi.org/10.3157/021.120.0213>

URL: <http://www.bioone.org/doi/full/10.3157/021.120.0213>

BioOne (www.bioone.org) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/page/terms_of_use.

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

***HARPACTOR ANGULOSUS* (REDUVIIDAE: HARPACTORINAE), A PREDATOR OF NEOTROPICAL SATURNIIDS, *HYLESIA* SPP. IN BRAZIL¹**

Alexandre I. A. Pereira,² José C. Zanuncio,² Hélcio R. Gil-Santana,³ Francisco S. Ramalho,⁴ Germano L. D. Leite,⁵ and José E. Serrão⁶

ABSTRACT: Caterpillars of the genus *Hylesia* (Lepidoptera: Saturniidae: Hemileucinae) can cause agricultural damages and dermatological lesions by direct contact to its urticating bristles. The biological control can regulate populations of these insects, but their natural enemies are poorly known. The occurrence of *Harpactor angulosus* (Lepeletier and Serville, 1825) (Reduviidae: Harpactorinae) predating caterpillars of *Hylesia* spp. is described. Adults of this predator were captured in an area of secondary forest in Viçosa, Minas Gerais State, Brazil and their pairs individualized in the field to obtain eggs. Adults of this predator presented sex dimorphism, with smaller males than females. They were found predating caterpillars of *Hylesia* spp., isolated from others to avoid the aposematic defense of this prey, when grouped. The presence of this predator, in the field, demonstrates its potential for the biological control of defoliating caterpillars. Studies on biological aspects and alimentary habits are important to understand the importance of *H. angulosus* in the biological control of defoliating caterpillars.

KEY WORDS: Biological control, predator, Harpactorinae, Saturniidae

Arthropods are important for the public health in several areas of the world because some of its species are vectors of infectious microorganisms (Pereira et al., 1998; Lounibos, 2002; Costa et al., 2007) and the use of poisonous substances to defend themselves against natural enemies. These compounds can cause irritating allergic reactions when in direct contact with humans (Carrijo-Carvalho and Chudzinski-Tavassi, 2007; Pereira et al., 2007).

Neotropical moths of the genus *Hylesia* are distributed from Mexico to Argentina, and they have urticating bristles. The burns caused by its urticating bristles are one of the main causes of dermatological burns in humans in America (Iserhard et al., 2007). These burns occur because the moths possess night habit, and they are attracted by public illumination. Such areas present high population

¹ Submitted on February 3, 2008. Accepted on May 5, 2008.

² Departamento de Biologia Animal/BIOAGRO; Universidade Federal de Viçosa; 36.571-000; Viçosa, Minas Gerais State, Brasil. E-mails: aiapereira@yahoo.com.br, zanuncio@ufv.br, respectively.

³ Departamento de Entomologia, Instituto Oswaldo Cruz, Avenida Brasil, 4365, Manguinhos, 21045-900; Rio de Janeiro State, Brasil. E-mail: helciogil@uol.com.br

⁴ Unidade de Controle Biológico (UCB)/Embrapa Algodão, Caixa Postal 174, 58107-720, Campina Grande, Paraíba State, Brazil. E-mail: framalho@pesquisador.cnpq.br

⁵ Setor de Fitotecnia, Núcleo de Ciências Agrárias (NCA), UFMG, Av. Osmane Barbosa, s/n, B. JK, Caixa Postal 135, CEP 39404-006 Montes Claros, Minas Gerais State, Brasil. E-mail: gldleite@ufmg.br

⁶ Departamento de Biologia Geral/BIOAGRO; Universidade Federal de Viçosa; 36.571-000; Viçosa, Minas Gerais State, Brasil. E-mail: jeserrao@ufv.br

and increased possibilities of direct contact with their larvae due to its aposematic aspect is what makes children and adults pay more attention to them.

Caterpillars of *Hylesia* spp. (Lepidoptera: Saturniidae), also, present potential as agricultural and forest pests in Brazil (Specht et al., 2006) and high taxonomic diversity (Lemaire, 2002), which hinders its recognition and control. Few chemical products are registered for controlling these insects in Brazil, which are considered secondary pests in eucalyptus plantations (Zanuncio et al., 1994) and *Ilex paraguaiensis* (Borges et al., 2003).

The knowledge of potential biological control agents of *Hylesia* spp. can be important for controlling these insects and reducing ecological disadvantages of chemical products application (Sengonca, 1998). *Hylesia* spp. have been controlled in Venezuela with aerial application of *Bacillus thuringiensis* var. *kurstaki* (Osborn et al., 2002), but this technique is unviable in some situations.

Caterpillars of *Hylesia* spp. are parasitized by tachinids (Diptera) as well as scelionids and ichneumonids (both Hymenoptera) (Pleigler, 1994). Caterpillars of *Hylesia metabus* Cramer (Lepidoptera: Saturniidae) were infected by microsporide in Ecuador (Osborn, 2002). However, predators of *Hylesia* spp. are poorly known, and the number of natural enemies of these Saturniidae is underestimated. The objective of this work was to register the occurrence of the predator *Harpactor angulosus* (Lepeletier and Serville, 1825) (Reduviidae: Harpactorinae) predating caterpillars of *Hylesia* spp. (Lepidoptera: Saturniidae) in the municipality of Viçosa, Minas Gerais State, Brazil.

METHODS

Three transects in line were established in an area along existent trails where insects were collected. Each one of these transects had 200 m length, 20 m width and a total sampled area of 12000 m². Each transect was walked, slowly, from 09:00 to 12:00 A.M., three times per week in September, October and November 2007, being considered, mainly, the transect and their margins. The collections were done by sweeping the vegetation because the use of light traps does not allow defining the alimentary diet of reduviids (Gil-Santana and Zeraik, 2003).

Specimens of harpactorine reduviids were collected in the experimental area of the beekeeping section of the Federal University of Viçosa (UFV) in Viçosa, Minas Gerais State, Brazil. This area had a secondary forest located at the southeast region of the Minas Gerais State (Figure 1) with geographical coordinates of 20° 28' and 21° 20' South Latitude and 42° 20' and 43° 13' West Longitude and 651 m altitude above sea level and with steep or soft and slopes hillsides. The climate is of Cwb type of tropical altitude with rainy summers and cold and dry winters with monthly mean annual temperature from 20 to 22°C and mean annual rainfall of 1,221.4 mm (Meira Neto and Martins, 2002). This area presents a fragment of secondary forest with native and exotic herbaceous and arboreal species including *Psidium guajava* (Myrtaceae) and *Eucalyptus* spp. (Myrtaceae) around 100 meters from the beehives.

The individuals collected were identified as *Harpactor angulosus* (Lepeletier and Serville, 1825) (Wygodzinsky 1946) by comparing them with the material deposited at the Entomological Collections of the National Museum of Rio de Janeiro (MNRJ) and Oswaldo Cruz Institute (IOC).

Figure 1. Location of the Municipality of Viçosa, Minas Gerais State, Brasil, and the area where *Harpactor angulosus* (Reduviidae: Harpactorinae) was collected.

RESULTS AND DISCUSSION

Predators of the subfamily Harpactorinae are very common with 2059 species in almost 300 genera (Brailovsky et al., 2007) and with large geographical distribution in America, Asia and Oceania (Forero, 2006; Zhao et al., 2007, respectively). However, the real number of species of this subfamily and information about the biology and ecology of many of their species are still ignored (Zhao et al., 2007).

Three nymphs and four adults of *H. angulosus* were found preying on isolated caterpillars of *Hylesia* spp. Also, specimens of the reduviid were collected under leaves of plants with vacuum cleaners entomological suckers (250 ml) and brought alive to the Laboratory of Biological Control of the Institute of Applied Biotechnology to the Agriculture (BIOAGRO) of UFV where nymphs and adults of this predator received, every 48 hours, sixth instar caterpillars of *Hylesia paulex* Dognin (Lepidoptera: Saturniidae). Two pairs of *H. angulosus* were put in white organza bags (20 x 30 cm) in the field involving extremities of guava tree branches and closed with strings (Zanuncio et al., 2004; Zanuncio et al., 2006) to obtain eggs to rear this Reduviidae predator.

Fig. 2. A. *Harpactor angulosus* (Reduviidae: Harpactorinae) clutch under a *Psidium guajava* L. (Myrtaceae) leaf. B. Egg, detail. C. Late instar nymph. D. *H. angulosus* female. E. Two *H. angulosus* reared in laboratory conditions. F. Natural predation on *Hylesia* sp. (Lepidoptera: Saturniidae) larvae by *H. angulosus*.

Harpactor angulosus laid its first egg batches five days after mating on the abaxial area of guava leaves in the organza bags (Figure 2A). However, egg masses of this predator were not found on guava leaves in the field which indicates that the oviposition substratum of this species remains unknown.

The eggs of each egg mass were glued sidelong to the substratum by an adherent substance which also linked them to the oviposition substratum. The production and/or use of adherent substances for individuals of the subfamily Harpactorinae has ecological importance to fix the eggs to the oviposition substratum and to reduce predation such as reported for species of the tribe Apiomerini (Gil-Santana et al., 2003) or to join soil or vegetable remains particles under the body as a disguise against natural enemies as found for nymphs of several species such as *Rhynocoris erythropus* Linnaeus (Heteroptera: Reduviidae) (Weirauch, 2006). Eggs of *H. angulosus* (Figure 2B) presented dark brown color, except for the operculum that is clear beige (Wygodzinsky, 1946).

Nymphs of *H. angulosus* (Figure 2C) were, easily, identified in the field by their clear white color while the adults were grayish (Figure 2D). The adults of this species present spiniform processes of variable sizes in the lobes of pronotum and lateral salencies at the abdomen, which are more or less apparent (Wygodzinsky, 1946). The females are larger than the males (Figure 2E) which is common for Harpactorinae (Gil-Santana and Milano, 2007). However, a useful and secure way to determine the sex is the observation of the globe aspect of the external male genital capsule (pygophore), which has greater importance to sex *H. angulosus* due to intraespecific variability in size in the same sex as observed for *H. angulosus*. Males were commonly found on the back of the females in laboratory, and they only abandoned this behavior to feed. Females fed with a male on its back (Figure 2E). Many Reduviidae species present parental care with the offspring (Ambrose, 1999; Tallamy et al., 2004), but this was not observed for *H. angulosus*. Males and females of this predator did not approach the egg mass, and they stayed indifferent when the egg masses were removed with a brush.

Caterpillars of *H. paulex* were preyed on almost immediately after they were offered to *H. angulosus* in the laboratory. This predator was observed feeding only on isolated caterpillars of *Hylesia* spp., suggesting larger success with this behavior to avoid the aposematic defense of this prey in colonies (Specht et al., 2006).

The attacks of *H. angulosus* on caterpillars of *Hylesia* spp. were done in the medium area of the body of this prey (Figure 2F). These preys become immobilized soon after being attacked by the probes of this predator which suggests a reduced handling time. The typical behavior of attack, with elevation of the first pair of paw legs as observed for *H. angulosus* is similar to that for *Apiomerus pilipes* Fabricius (Heteroptera: Harpactorinae) (Silva and Gil-Santana, 2004), with subsequent capture and inserting the probes of the rostrum into the prey.

Harpactor angulosus preyed on *Tenebrio molitor* L. (Coleoptera: Tenebrionidae) pupae and *Anticarsia gemmatalis* Huebner (Lepidoptera: Noctuidae) caterpillars in the laboratory. This reinforces the generalist feeding habit of this predator. This agrees with previous observations that Reduviidae species are usually polyphagous (Sahayaraj and Paulraj, 2001; Lioni and Cividanes, 2004) and also feeding on beneficial insects as bees and ladybugs (Gil-Santana et al., 2003; Silva and Gil-Santana, 2004; Marques et al., 2006). *Harpactus angulosus* is probably polyphagous too, since *Hylesia* spp. did not occur during the whole year but only in the period from the beginning of the spring to the end of the summer in Brazil.

The number of prey and biological aspects of *H. angulosus* should be studied to evaluate the potential of this natural enemy in biological control programs of defoliating caterpillars in agricultural and forest systems.

ACKNOWLEDGMENTS

The authors thank the Brazilian agencies “Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq)” and the “Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG)” for funding.

LITERATURE CITED

- Ambrose, D. P. 1999. Assassin Bugs. Science Publishers, Inc., New Hampshire, U.S.A. 337 pp.
- Borges, L. R., S. M. N. Lázari, and F. A. Lázari. 2003. Comparação dos sistemas de cultivo nativo e adensado de erva mate, *Ilex paraguariensis* St. Hil., quanto à ocorrência e flutuação populacional de insetos. Revista Brasileira de Entomologia 47: 563-568.
- Brailovsky, H., R. Marino, and E. Barrera. 2007. Cinco especies de *Pselliopus* (Hemiptera: Heteroptera: Reduviidae: Harpactorinae: Harpactorini) para México. Revista Mexicana de Biodiversidad 78: 85-98.
- Carrijo-Carvalho, L. C. and A. M. Chudzinski-Tavassi. 2007. The venom of the *Lonomia* caterpillar: An overview. Toxicon 49: 741-757.
- Costa, S. M., M. Cechinel, V. Bandeira, J. C. Zanuncio, R. Lainson, and E. F. Rangel. 2007. *Lutzomyia* (*Nyssomyia*) *whitmani* s.l. (Antunes & Coutinho, 1939) (Diptera: Psychodidae: Phlebotominae): geographical distribution and the epidemiology of American cutaneous leishmaniasis in Brazil – Mini-review. Memórias do Instituto Oswaldo Cruz 102: 149-153.
- Forero, D. 2006. New records of Reduviidae (Hemiptera: Heteroptera) from Colombia and other Neotropical countries. Zootaxa 1107: 1-47.
- Gil-Santana, H. R. and S. O. Zeraik. 2003. Reduviidae de Cabo Frio, Rio de Janeiro, Brasil (Hemiptera, Heteroptera). Revista Brasileira de Zoociências 5: 101-120.
- Gil-Santana, H. R., L. A. A. Costa, D. Forero, and S. O. Zeraik. 2003. Sinopse dos Apiomerini, com chave ilustrada para os gêneros (Hemiptera-Heteroptera, Reduviidae, Harpactorinae). Publicações Avulsas do Museu Nacional 97: 1-24.
- Gil-Santana, H. R. and P. Milano. 2007. Descrição de *Apiomerus costai* sp. nov. do Mato Grosso, com notas taxonômicas sobre *Apiomerus* Hahn (Hemiptera, Heteroptera: Reduviidae, Harpactorinae, Apiomerini). Neotropical Entomology 36: 314-316.
- Iserhard, C. A., L. A. Kaminski, M. O. Marchiori, E. C. Teixeira, and H. P. Romanowski. 2007. Occurrence of lepidopterism caused by the moth *Hylesia nigricans* (Berg) (Lepidoptera: Saturniidae) in Rio Grande do Sul State, Brazil. Neotropical Entomology 36(4): 612-615.
- Lemaire, C. 2002. The Saturniidae of America – Hemileucinae Part A. Keltern, Goecke & Evers, Keltern, Germany. Deutschland. 688 pp.
- Lioni, L. P. and F. J. Cividanes. 2004. Ecological life table of the citrus leafminer *Phyllocnistis citrella* Stainton (Lepidoptera: Gracilidae). Neotropical Entomology 33: 407-415.

- Lounibos, L. P.** 2002. Invasions by insect vectors of human disease. *Annual Review of Entomology* 47: 66-233.
- Marques, O. M., H. R. Gil-Santana, M. L. Coutinho, and D. D. Silva Júnior.** 2006. Percevejos predadores (Hemiptera, Reduviidae, Harpactorinae) em fumo (*Nicotiana tabacum* L.) no município de Cruz das Almas, Bahia. *Revista Brasileira de Zoociências* 8: 55-60.59.
- Meira-Neto, J. A. A and F. R. Martins.** 2002. Composição florística de uma floresta estacional semidecidual Montana no município de Viçosa-MG. *Árvore* 26: 437-446.
- Osborn, F.** 2002. Primer reporte de un microsporidio parasítico en larvas de *Hylesia metabus* Cramer, 1775 (Lepidoptera: Saturniidae). *Entomotopica* 17: 185-188.
- Osborn, F., L. Berlioz, J. Vitelli-Flores, W. Monsalve, B. Dorta, and V. R. Lemoine.** 2002. Pathogenic effects of bacteria isolated from larvae of *Hylesia metabus* Cramer (Lepidoptera: Saturniidae). *Journal of Invertebrate Pathology* 80: 7-12.
- Pereira, H., C. M. Penido, M. S. Martins, and L. Diotaiuti.** 1998. Comparative kinetics of blood-meal intake by *Triatoma infestans* and *Rhodnius prolixus*, the two principal vectors of Chagas disease. *Medical and Veterinary Entomology* 12: 84-88.
- Pereira, F. F., J. P. M. Felipe, G. C. Canevari, O. H. H. Mielke, J. C. Zanuncio, and J. E. Serrão.** 2007. Biological aspects of *Dirphia moderata* Bouvier, 1929 (Lepidoptera: Saturniidae) in *Eucalyptus cloeziana* and *Psidium guajava*. *Brazilian Archives of Biology and Technology* 49: 232-243.
- Pleiger, R. S.** 1994. Catalog of parasitoids of Saturniidae of the world. *Journal of Research on the Lepidoptera* 33: 1-121.
- Sahayaraj, K. and M. G. Paulraj.** 2001. Rearing and life table of reduviid predator *Rhynocoris marginatus* Fab. (Het., Reduviidae) on *Spodoptera litura* Fab. (Lep., Noctuidae) larvae. *Journal of Applied Entomology* 125: 321-325.
- Sengonca, C.** 1998. Conservation and enhancement of natural enemies in biological control. *Phytoparasitica* 26: 187-190.
- Silva, A. C. and H. R. Gil-Santana.** 2004. Predation of *Apiomerus pilipes* (Fabricius) (Hemiptera, Reduviidae, Harpactorinae, Apiomerini) over Meliponinae bees (Hymenoptera, Apidae), in the State of Amazonas, Brazil. *Revista Brasileira de Zoologia* 21: 769-774.
- Specht, A., A. C. Formentini, and E. Corseuil.** 2006. Biologia de *Hylesia nigricans* (Berg) (Lepidoptera, Saturniidae, Hemileucinae). *Revista Brasileira de Zoologia* 23: 248-255.
- Tallamy, D. W., E. Walsh, and D. C. Peck.** 2004. Revisiting paternal care in the assassin bug, *Atopozelus pallens* (Heteroptera: Reduviidae). *Journal of Insect Behavior* 17: 431-436.
- Weirauch, C.** 2006. Anatomy of disguise: camouflaging structures in nymphs of some Reduviidae (Heteroptera). *American Museum Novitates*. 3542: 1-18.
- Wygodzinsky, P.** 1946. Sobre um novo gênero de Harpactorinae do Brasil, com notas sobre os gêneros *Harpactor* Laporte e *Erbessus* Stål (Reduviidae, Hemiptera). *Revista de Entomologia* 17: 401-424.
- Zanuncio, J. C., E. C. Nascimento, J. F. Garcia, and T. V. Zanuncio.** 1994. Major lepidopterous defoliators of eucalypt in southeast Brazil. *Forest Ecology and Management* 65: 53-63.
- Zanuncio, J. C., M. C. Lacerda, J. S. Zanuncio Júnior, T. V. Zanuncio, A. M. C. Silva, and M. C. Espindula.** 2004. Fertility table and rate of population growth of the predator *Supputius cincticeps* (Heteroptera: Pentatomidae) on one plant of *Eucalyptus cloeziana* in the field. *Annals of Applied Biology* 144: 357-361.
- Zanuncio, J. C., W. P. Lemos, M. C. Lacerda, T. V. Zanuncio, J. E. Serrão, and E. Bauce.** 2006. Age-dependent fecundity and fertility life tables of the Predator *Brontocoris tabidus* (Heteroptera: Pentatomidae) under field conditions. *Journal of Economic Entomology* 99: 401-407.
- Zhao, P., J. Constant, and W. Cai.** 2007. Revision of the Papua New Guinea genus *Elemacoris* Miller (Hemiptera: Reduviidae: Harpactorinae). *Zootaxa* 1556: 51-60.