

## Moderate Continuous Aerobic Exercise Training Improves Cardiomyocyte Contractility in $\beta_1$ Adrenergic Receptor Knockout Mice

Aurora Corrêa Rodrigues,<sup>1</sup> Antônio José Natali,<sup>1</sup> Daise Nunes Queiroz da Cunha,<sup>1</sup> Alexandre Jayme Lopes Dantas Costa,<sup>1</sup> Anselmo Gomes de Moura,<sup>1</sup> Miguel Araújo Carneiro-Júnior,<sup>1</sup> Leonardo Bonato Félix,<sup>1</sup> Patrícia Chakur Brum,<sup>2</sup> Thales Nicolau Prímola-Gomes<sup>1</sup>

Universidade Federal de Viçosa (UFV),<sup>1</sup> Viçosa, MG; Universidade de São Paulo (USP),<sup>2</sup> São Paulo, SP – Brazil

### Abstract

**Background:** The lack of cardiac  $\beta_1$ -adrenergic receptors ( $\beta_1$ -AR) negatively affects the regulation of both cardiac inotropy and lusitropy, leading, in the long term, to heart failure (HF). Moderate-intensity aerobic exercise (MCAE) is recommended as an adjunctive therapy for patients with HF.

**Objective:** We tested the effects of MCAE on the contractile properties of left ventricular (LV) myocytes from  $\beta_1$  adrenergic receptor knockout ( $\beta_1$ ARKO) mice.

**Methods:** Four- to five-month-old male wild type (WT) and  $\beta_1$ ARKO mice were divided into groups: WT control (WTc) and trained (WTt); and  $\beta_1$ ARKO control ( $\beta_1$ ARKOc) and trained ( $\beta_1$ ARKOt). Animals from trained groups were submitted to a MCAE regimen (60 min/day; 60% of maximal speed, 5 days/week) on a treadmill, for 8 weeks.  $P \leq 0.05$  was considered significant in all comparisons.

**Results:** The  $\beta_1$ ARKO and exercised mice exhibited a higher ( $p < 0.05$ ) running capacity than WT and sedentary ones, respectively. The  $\beta_1$ ARKO mice showed higher body (BW), heart (HW) and left ventricle (LVW) weights, as well as the HW/BW and LVW/BW than WT mice. However, the MCAE did not affect these parameters. Left ventricular myocytes from  $\beta_1$ ARKO mice showed increased ( $p < 0.05$ ) amplitude and velocities of contraction and relaxation than those from WT. In addition, MCAE increased ( $p < 0.05$ ) amplitude and velocities of contraction and relaxation in  $\beta_1$ ARKO mice.

**Conclusion:** MCAE improves myocyte contractility in the left ventricle of  $\beta_1$ ARKO mice. This is evidence to support the therapeutic value of this type of exercise training in the treatment of heart diseases involving  $\beta_1$ -AR desensitization or reduction. (Arq Bras Cardiol. 2018; [online].ahead print, PP.0-0)

**Keywords:** Heart Failure; Exercise; Myocardial Contraction; Myocytes, Cardiac; Adrenergic beta 1 Receptor Antagonists; Mice.

### Introduction

Chronic sympathetic hyperactivity resulting from altered autonomic nervous system balance is common in many cardiovascular disease states, ending up in heart failure (HF), and is related to a higher incidence of morbidity and mortality.<sup>1,2</sup> Such hyperactivity is paralleled by a decrease in  $\beta$ -adrenergic receptors ( $\beta$ -AR) density and desensitization of the remaining  $\beta$ -AR, thus leading to a reduced cardiac contractile response to  $\beta$ -AR activation.<sup>3</sup> In this framework,  $\beta_1$ -AR, predominant in the heart, is selectively reduced, resulting in a modified ratio of  $\beta_1$  to  $\beta_2$  subtypes,<sup>4</sup> and  $\beta_2$ -AR are markedly coupled to inhibitory G protein. Consequently, inasmuch as the  $\beta_1$ -AR phosphorylates several  $\text{Ca}^{2+}$  regulatory

proteins involved in cardiomyocyte excitation-contraction coupling,<sup>5-7</sup> cardiac chronotropism, inotropism and lusitropism are impaired under adrenergic stimulation.<sup>8</sup>

Exercise training in cardiac rehabilitation is very important in several cardiovascular diseases, including chronic HF.<sup>9</sup> Continuous moderate-intensity aerobic exercise (MCAE) is, at present, the best-established form of exercise for this population because of its efficacy and safety.<sup>10</sup> For example, aerobic exercise training recovers the resting autonomic balance in HF patients by reducing the resting sympathetic nerve activity,<sup>11</sup> and restoring the parasympathetic tone to the heart.<sup>12,13</sup> In the myocardium, aerobic exercise training increases stroke volume and, hence, cardiac output in patients<sup>14,15</sup> and in animal models of HF,<sup>8</sup> although some studies failed to confirm such benefits.<sup>11,12</sup> At the cellular level, studies on animal models for sympathetic hyperactivity have demonstrated aerobic exercise training improves the net balance of cardiac  $\text{Ca}^{2+}$  handling proteins either alone<sup>8,16</sup> or in combination with  $\beta$ -blockers.<sup>17</sup> Nevertheless, whether MCAE training affects mechanical properties of single myocytes in a heart lacking  $\beta_1$ -AR remains to be elucidated.

Therefore, the aim of this study was to test the effects of an MCAE program on mechanical properties of single

**Mailing Address:** Thales Nicolau Prímola Gomes •

Universidade Federal de Viçosa - Departamento de Educação Física - Av. Ph. Rolfs, s/n. Postal Code 36570-000, Campus Universitário, Viçosa, MG – Brazil

E-mail: thales.gomes@ufv.br, thalesprimola@gmail.com

Manuscript received May 14, 2017, revised manuscript September 15, 2017, accepted September 22, 2017

**DOI:** 10.5935/abc.20180025

left ventricular (LV) myocytes in  $\beta$ <sub>1</sub>AR knockout ( $\beta$ <sub>1</sub>ARKO) mice. We hypothesized that MCAE training positively affects mechanical properties of LV myocytes from  $\beta$ <sub>1</sub>ARKO mice.

## Methods

### Experimental animals

A cohort of 4- to 5-month-old male wild type (WT) and congenic  $\beta$ <sub>1</sub>ARKO mice in the C57Bl6/J genetic background were studied. Mice were maintained in cages under a 12:12-h light-dark cycle in a temperature-controlled room (22°C), with free access to water and standard rodent diet. WT and  $\beta$ <sub>1</sub>ARKO mice were randomly assigned into one of the following groups by using the simple random sampling: WT control (WTc), WT trained (WTt),  $\beta$ <sub>1</sub>ARKO control ( $\beta$ <sub>1</sub>ARKOc) and  $\beta$ <sub>1</sub>ARKO trained ( $\beta$ <sub>1</sub>ARKOt). The sample size was defined by convenience. All groups initiated the experimental period with eight animals, however, during the cardiomyocyte isolation procedure, some animals/hearts were lost. Thus, the final number of animals in each group is specified in figures and table. Body weight (BW) was measured every week. The experimental protocols were approved by the Ethics Committee for Animal Use at the Viçosa Federal University (protocol #59/2012) in accordance with the Guide for the Care and Use of Laboratory Animals/2011.

### Exercise training protocol and graded treadmill exercise test

MCAE was performed on a motor treadmill (Insight Equipamentos Científicos, Brazil) 5 days/week (Monday to Friday), 60 min/day, for 8 weeks. Over the first week, the duration and running speed of exercise were progressively increased from 10 minutes and 10% of the maximal speed until 60 minutes and 60% of the maximal speed, achieved during a graded treadmill exercise test. At the end of the fourth week of aerobic exercise training, graded treadmill exercise tests were repeated to readjust the running speed. This intensity was maintained during the rest of the training period. During the training period, animals from the untrained groups were handled every day and subjected to a short period of mild exercise (5 min, 0% grade, 5 m/min, 3 days/week). The exercise capacity estimated by total distance run was evaluated using a graded treadmill exercise protocol for mice (Panlab/Harvard Apparatus, Spain), as described previously.<sup>18</sup> Briefly, after being adapted to the treadmill for 1 week (10 min/day, 0% grade, 0.3 km/h), mice were placed in the exercise streak and allowed to acclimatize for at least 30 minutes. The graded treadmill exercise test began at 6 m/min with no grade and increased by 3 m/min every 3 minutes until fatigue, which was defined as when the test was interrupted because the animals could no longer keep pace with the treadmill speed. The graded treadmill exercise test was performed in WT and  $\beta$ 1ARKO untrained and exercise-trained groups before and after the exercise training period.

### Cardiomyocyte isolation

Forty-eight hours after the last exercise training session, mice were weighed and killed by decapitation, and their hearts were removed quickly. Left ventricular myocytes were enzymatically isolated as described previously.<sup>19</sup> Briefly, hearts

were mounted onto a Langendorff system and perfused with calcium-free HEPES-Tyrode solution for 6 minutes with the following composition (in mM): 130 NaCl, 1.43 MgCl<sub>2</sub>, 5.4 KCl, 0.4 NaH<sub>2</sub>PO<sub>4</sub>, 0.75 CaCl<sub>2</sub>, 25 HEPES, 22 glucose, 0.01  $\mu$ g/ml insulin, 0.1 EGTA, pH 7.4, at 37°C. Afterwards, the hearts were perfused for 7-10 minutes with a solution containing 1 mg/ml collagenase type II (Worthington, USA) and CaCl<sub>2</sub> (0.8  $\mu$ M). The digested heart was then removed from the perfusion apparatus and the heart and left ventricle were carefully weighed. Left ventricle was cut into small pieces and placed into conical flasks with collagenase-containing solution. The cells were dispersed by agitating the flasks for periods of 3 minutes at 37°C. Single cells were separated from the non-dispersed tissue by filtration. The resulting cell suspension was centrifuged and resuspended in HEPES-Tyrode solution containing CaCl<sub>2</sub> (2.5 and 5  $\mu$ M, subsequently). The isolated cells were stored in HEPES-Tyrode solution containing 10  $\mu$ M CaCl<sub>2</sub> at room temperature until use. Only calcium-tolerant, quiescent, rod-shaped cardiomyocytes showing clear cross-striations were studied. The isolated cardiomyocytes were used within 2-3 hours of isolation.

### Cell contractility measurement

Cell contractility was evaluated as described previously.<sup>20</sup> Briefly, the isolated cells were placed in a chamber with a glass coverslip base mounted onto the stage of an inverted microscope (Nikon Eclipse, TS100). The chamber was perfused with HEPES-Tyrode solution plus 10  $\mu$ M CaCl<sub>2</sub> at 37°C. Steady-state 1-Hz contractions were elicited via platinum bath electrodes (Myopacer, Field Stimulator, IonOptix) with 5-ms voltage pulses and an intensity of 40 V. The cells were visualized on a personal computer monitor with a NTSC camera (MyoCam, IonOptix) in partial scanning mode. The image was used to measure cell shortening (our index of contractility) in response to electrical stimulation using a video motion edge detector (IonWizard, IonOptix). The cell image was sampled at 240 Hz. Cell shortening was calculated from the output of the edge detector using an A/D converter (IonOptix, Milton, MA). Cell shortening (expressed as percentage of resting cell length) and the velocities of shortening and relaxation were calculated.

### Statistics

Data were subjected to Shapiro-Wilk or Kolmogorov-Smirnov normality tests as appropriate. Paired *t* test was used to compare initial and final BW in each group. The comparisons among groups of the values of BW, heart weight (HW), left ventricular weight (LVW) and ratios, as well as cell contraction were made using a two-way ANOVA followed by Tukey test using software SigmaPlot®, 12.5 version (Systat Software, San Jose, CA). Data are presented as means  $\pm$  SD. A statistical significance level of 5% was adopted. Numbers of mice, hearts, and myocytes used are given in the relevant table and figure legends.


## Results

Table 1 shows BW and LVW. The initial BW of  $\beta$ <sub>1</sub>ARKO animals was higher as compared to their respective control

**Table 1 – Body and left ventricular weights**

	WTc (n = 7)	WTt (n = 6)	$\beta_1$ ARKOc (n = 7)	$\beta_1$ ARKOt (n = 6)
Initial BW, g	27.43 ± 2.46	26.50 ± 2.45	33.86 ± 2.46	32.67 ± 2.23
Final BW, g	29.86 ± 2.64*	28.67 ± 2.64*	37.14 ± 2.64*	34.33 ± 2.55*
HW, mg	231.00 ± 37.57	226.00 ± 37.48	302.00 ± 37.57	317.00 ± 37.48
LVW, mg	146.00 ± 20.82	141.00 ± 20.82	184.00 ± 20.82	194.00 ± 20.82
HW/BW, mg/g	7.73 ± 0.85	7.86 ± 0.86	8.12 ± 0.85	9.22 ± 0.86
LVW/BW, mg/g	4.89 ± 0.48	4.94 ± 0.49	4.96 ± 0.48	5.66 ± 0.49

Values are means ± SD; WTc: wild-type control; WTt: wild-type trained;  $\beta_1$ ARKOc: knockout  $\beta_1$ -ARs control;  $\beta_1$ ARKOt: knockout  $\beta_1$ -ARs trained; BW: body weight; HW: heart weight; LVW: left ventricular weight; N: number of animals; \* $p < 0.05$  vs. initial BW within the same group. Statistical differences were determined by paired t test.



**Figure 1 – Total distance run.** Values are means ± SD of 8 mice in each group. \* $p < 0.05$  vs. WTc group; § $p < 0.05$  vs. WTt group; # $p < 0.05$  vs.  $\beta_1$ ARKOc group.

WT animals. As expected, the final BW of each group was higher, compared to their respective initial BW. The final BW was higher ( $p < 0.05$ ) in  $\beta_1$ ARKO ( $\beta_1$ ARKOc +  $\beta_1$ ARKOt), compared to WT mice (WTc + WTt). However, the final BW was not affected ( $p > 0.05$ ) by the MCAE. Likewise, HW was higher in  $\beta_1$ ARKO than in WT mice, but no effect of MCAE was observed ( $p > 0.05$ ). Regarding LVW,  $\beta_1$ ARKO presented higher values than WT mice; nevertheless, no effect of MCAE was found ( $p > 0.05$ ). As for the ratios,  $\beta_1$ ARKO mice presented higher HW to BW ratio than WT mice. However, it was not affected by MCAE ( $p > 0.05$ ). The LVW to BW ratio was higher in  $\beta_1$ ARKO mice, compared to WT mice, but there was no effect of MCAE.

Figure 1 shows the physical capacity.  $\beta_1$ ARKO animals ( $\beta_1$ ARKOc +  $\beta_1$ ARKOt) exhibited a longer running distance, compared to WT animals (WTc + WTt). In addition, trained animals presented a longer running distance, compared to their respective controls.


The contractile properties of single LV myocytes are presented in Figure 2.  $\beta_1$ ARKO myocytes ( $\beta_1$ ARKOc +  $\beta_1$ ARKOt) had higher shortening amplitude than WT cells (WTc + WTt). The amplitude of shortening in  $\beta_1$ ARKOt myocytes was higher, compared to  $\beta_1$ ARKOc and WTt cells; and in WTc cells, compared to WTt cells (Figure 2A). Regarding the contractile time course,  $\beta_1$ ARKOc myocytes exhibited higher velocity of shortening than WTc cells. In addition,  $\beta_1$ ARKOt myocytes exhibited higher velocity of shortening than  $\beta_1$ ARKOc and WTt

cells (Figure 2B). As for the velocity of relaxation,  $\beta_1$ ARKOc myocytes exhibited higher values than WTc cells. Moreover,  $\beta_1$ ARKOt myocytes exhibited higher velocity of relaxation than  $\beta_1$ ARKOc and WTt cells (Figure 2C).

## Discussion

In this study, we tested the effects of MCAE on mechanical properties of LV myocytes from  $\beta_1$ ARKO mice. The main finding was that MCAE increased the amplitude of shortening and velocities of shortening and relaxation in  $\beta_1$ ARKO mice myocytes.

The initial and final BWs were higher in  $\beta_1$ ARKO than in WT mice. Similar results have been observed elsewhere.<sup>21</sup>  $\beta$ -AR activation in adipose tissue leads to cyclic adenosine monophosphate (cAMP) production, which activates protein kinase A (PKA) and stimulates lipolysis. Even though  $\beta_3$ -AR is the predominant receptor in rodent adipose tissue, mice overexpressing  $\beta_1$ -AR exhibit increased adipocyte lipolytic activity.<sup>22</sup> Therefore,  $\beta_1$ ARKO mice may have reduced lipolysis, which would influence the amount of body fat and, consequently, BW.<sup>23</sup> Nevertheless, our MCAE did not affect the final BW. Regarding HW,  $\beta_1$ ARKO mice exhibited heavier hearts and left ventricles than WT mice, as well as higher HW to BW and LVW to BW ratios. Our MCAE, nevertheless, did not modify these cardiac parameters. Exercise-induced cardiac hypertrophy in WT mice has been demonstrated elsewhere;<sup>24-26</sup> nevertheless in  $\beta_1$ ARKO mice, as far as we know, no data have been reported.



**Figure 2 – Cell contractility.** A) Shortening. B) Velocity of shortening. C) Velocity of relaxation. WTc, wild-type control (n = 7; N = 14-39 cells from each mouse); WTt, wild-type trained (n = 6; N = 8-27 cells from each mouse);  $\beta_1$ ARKOc, knockout  $\beta_1$ -AR control (n = 7; N = 24-31 cells from each mouse);  $\beta_1$ ARKOt, knockout  $\beta_1$ -AR trained (n = 6; N = 17-29 cells from each mouse). Values are means  $\pm$  SD. \*p < 0.05 vs. WTc group; §p < 0.05 vs. WTt group; #p < 0.05 vs.  $\beta_1$ ARKOc group.

We observed that trained mice (WTt and  $\beta_1$ ARKOt) showed longer total running distance than their respective controls (WTc and  $\beta_1$ ARKOc). This MCAE-induced increase may be associated with cardiovascular adaptations, which are known features of aerobic exercise training.<sup>27</sup> Previous studies using the same aerobic exercise training protocol also observed increased exercise capacity in trained animals.<sup>8,17</sup> Specifically, the  $\beta_1$ ARKO groups showed longer total running distance than the WT groups. It is known that sympathetic activation during aerobic exercise promotes glycogenolysis by  $\beta$ -AR pathway.<sup>28,29</sup> Probably, the  $\beta_1$ ARKO mice have compensatory mechanisms in the skeletal muscle, such as modified  $\beta_2$  and  $\alpha_1$  adrenergic signaling pathways, which could improve glycogenolysis,

gluconeogenesis, insulin-independent glucose uptake and lipolysis in the skeletal muscles.<sup>30</sup> These compensatory mechanisms may have led to increased exercise performance in  $\beta_1$ ARKO mice. However, inasmuch as this issue is not the focus of this study, further investigations are needed to test the hypothesis that  $\beta_1$ ARKO mice increase exercise performance by altering  $\beta_2$  and  $\alpha_1$  adrenergic signaling pathways.

Although myocytes from  $\beta_1$ ARKO mice had a higher amplitude of shortening than cells from WT mice, an independent factor effect, LV myocytes from  $\beta_1$ ARKOc and WTc groups had similar contractile properties. Although  $\beta_1$ AR is the predominant adrenergic receptor subtype expressed in the heart in terms of density and modulation of cardiac contraction,<sup>31,32</sup> its deletion had little impact on resting cardiac function, but had significant effects on cardiac function after  $\beta$ -agonist stimulation.<sup>33</sup> Other studies did not observe changes in cardiomyocyte contractility upon loss of  $\beta_1$ -AR<sup>34</sup> or  $\beta_{1/2}$ -AR under basal conditions.<sup>35</sup> Therefore, the similarity between  $\beta_1$ ARKOc and WTc groups suggests that  $\beta_1$ -AR has little impact on the contractile properties of cardiomyocytes under basal conditions.

More important, the MCAE program increased the amplitude of shortening of LV myocytes from  $\beta_1$ ARKO mice. The MCAE may have triggered two compensatory mechanisms in the heart of  $\beta_1$ ARKO mice. First, an increase in  $\alpha_1$ -ARs signaling is common under situations of  $\beta_1$ -ARs desensitization when the reduction of  $\beta_1$ -adrenergic signaling is compensated by an increase in  $\alpha_1$ -adrenergic signaling pathway, which could help preserve cardiac function.<sup>36</sup> Although not evaluated here, an increased inotropic responsiveness of rat cardiomyocytes via  $\alpha_1$ -AR stimulation was found as an adaptation to aerobic exercise training.<sup>37,38</sup> Moreover, the potential therapeutic role of  $\alpha_1$ -ARs to maintain normal cardiac function, especially in terms of commitment of the  $\beta_1$ -adrenergic signaling pathway, has been proposed in previous studies.<sup>37-40</sup> Second, MCAE may have reduced the responsiveness of  $\beta_2$ -AR in myocytes of  $\beta_1$ ARKO mice. When  $\beta_2$ -AR coupling to  $G_i$  protein is reduced, the inhibitory effect of the receptor to adenylate cyclase activation is also reduced,<sup>5</sup> which causes an increased cAMP production and phosphorylation of proteins involved in cardiomyocyte excitation-contraction coupling.<sup>6</sup>

The time courses of  $\beta_1$ ARKO LV myocyte contraction and relaxation were also improved by MCAE, indicating enhanced systolic and diastolic functions. The  $Ca^{2+}$  regulatory proteins modulate cardiomyocyte mechanical properties. While faster myocyte contraction is associated with increased density and/or activity of L-type  $Ca^{2+}$  channels and  $RyR_2$ , quicker relaxation is dependent on the increased activity and/or density of SERCA2a, PLB and NCX.<sup>6</sup> Although not measured in the present study, MCAE may have improved the net balance of cardiac  $Ca^{2+}$  handling proteins in  $\beta_1$ ARKO mice. Such adaptations have been demonstrated previously in a different model for sympathetic hyperactivity.<sup>8,16</sup> In addition, endurance-exercise training may have reduced the  $\beta/\alpha$ -MHC ratio,<sup>20</sup> which would also help explain the increased velocities of LV myocyte contraction and relaxation.

In recent years, high-intensity interval training (HIIT) has emerged as the method that leads to significant benefits to cardiac function. For instance, mice submitted to HIIT

presented higher cardiomyocyte contractile function by increasing the expression and activity of calcium cycle regulatory proteins, as compared to those submitted to MCAE.<sup>41-43</sup> Thus, it is possible that cardiomyocytes from  $\beta_1$ ARKO mice might be more responsive to HIIT. However, in the present study, we chose the MCAE because the effects of such exercise protocol on the single cardiomyocyte contractility in  $\beta_1$ ARKO mice are not known. We believe that future studies using HIIT would provide interesting findings in this animal model.

This study has limitations. First, we used global KO mice and systemic alterations confounding the exercise effects may have occurred, thus these results have to be interpreted with caution. Second, although WTt animals had improved their exercise capacity, unexpectedly their LV myocytes presented lower cell shortening than WTc mice. This finding really intrigued us, and, unfortunately, we cannot explain it.

## Conclusion

In conclusion, MCAE training improves myocyte contractility in the left ventricle of  $\beta_1$ ARKO mice. This finding has potential clinical implications and supports the therapeutic value of this type of exercise training in the treatment of heart diseases involving  $\beta_1$ -AR desensitization or reduction.

## References

1. Barretto AC, Santos AC, Munhoz R, Rondon MU, Franco FG, Trombetta IC, et al. Increased muscle sympathetic nerve activity predicts mortality in heart failure patients. *Int J Cardiol.* 2009;135(3):302-7. doi: 10.1016/j.ijcard.2008.03.056.
2. Braunwald E. Heart failure. *JACC Heart Fail.* 2013 Feb;1(1):1-20. doi: 10.1016/j.jchf.2012.10.002.
3. Bristow MR, Ginsburg R, Minobe W, Cubicciotti RS, Sageman WS, Lurie K, et al. Decreased catecholamine sensitivity and beta-adrenergic-receptor density in failing human hearts. *N Engl J Med.* 1982;307(4):205-11. doi: 10.1056/NEJM198207223070401.
4. Wallukat G. The beta-adrenergic receptors. *Herz.* 2002;27(7):683-90. doi: 10.1007/s00059-002-2434-z.
5. Xiang Y, Kobilka BK. Myocyte adrenoceptor signaling pathways. *Science.* 2003;300(5625):1530-2. doi: 10.1126/science.1079206.
6. Bers DM. Cardiac excitation-contraction coupling. *Nature.* 2002;415(6868):198-205. doi: 10.1038/415198a.
7. Kubalova Z, Terentyev D, Viatchenko-Karpinski S, Nishijima Y, Gyorke I, Terentyeva R, et al. Abnormal intrastore calcium signaling in chronic heart failure. *Proc Natl Acad Sci U S A.* 2005;102(39):14104-9. doi: 10.1073/pnas.0504298102.
8. Rolim NP, Medeiros A, Rosa KT, Mattos KC, Irigoyen MC, Krieger EM, et al. Exercise training improves the net balance of cardiac Ca<sup>2+</sup> handling protein expression in heart failure. *Physiol Genomics.* 2007;29(3):246-52. doi: 10.1152/physiolgenomics.00188.2006.
9. Gielen S, Laughlin MH, O'Conner C, Duncker DJ. Exercise training in patients with heart disease: review of beneficial effects and clinical

## Author contributions

Conception and design of the research: Rodrigues AC, Natali AJ, Brum PC, Prímola-Gomes TN; Acquisition of data: Rodrigues AC, Cunha DNQ, Costa AJLD, Moura AG; Analysis and interpretation of the data: Rodrigues AC, Natali AJ, Carneiro-Júnior MA, Prímola-Gomes TN; Statistical analysis: Rodrigues AC, Félix LB; Writing of the manuscript: Rodrigues AC, Natali AJ, Prímola-Gomes TN; Critical revision of the manuscript for intellectual content: Natali AJ, Prímola-Gomes TN.

## Potential Conflict of Interest

No potential conflict of interest relevant to this article was reported.

## Sources of Funding

This study was funded by CNPq, Fapemig, Capes and Fapesp.

## Study Association

This article is part of the thesis of master submitted by Aurora Corrêa Rodrigues, from Universidade Federal de Viçosa.

## Ethics approval and consent to participate

This study was approved by the Ethics Committee on Animal Experiments of the Universidade Federal de Viçosa under the protocol number #59/2012.

- recommendations. *Prog Cardiovasc Dis.* 2015;57(4):347-55. doi: 10.1016/j.pcad.2014.10.001.
10. O'Connor CM, Whellan DJ, Lee KL, Keteyian SJ, Cooper LS, Ellis SJ, et al. Efficacy and safety of exercise training in patients with chronic heart failure: HF-ACTION randomized controlled trial. *JAMA.* 2009;301(14):1439-50. doi: 10.1001/jama.2009.454.
11. Roveda F, Middlekauff HR, Rondon MU, Reis SF, Souza M, Nastari L, et al. The effects of exercise training on sympathetic neural activation in advanced heart failure: a randomized controlled trial. *J Am Coll Cardiol.* 2003;42(5):854-60. PMID: 12957432.
12. Ichige MH, Santos CR, Jordao CP, Ceroni A, Negrao CE, Michelini LC. Exercise training preserves vagal preganglionic neurones and restores parasympathetic tonus in heart failure. *J Physiol.* 2016;594(21):6241-54. doi: 10.1113/jp272730.
13. Negrao CE, Middlekauff HR, Gomes-Santos IL, Antunes-Correa LM. Effects of exercise training on neurovascular control and skeletal myopathy in systolic heart failure. *Am J Physiol Heart Circ Physiol.* 2015;308(8):H792-802. doi: 10.1152/ajpheart.00830.2014
14. Erbs S, Linke A, Gielen S, Fiehn E, Walther C, Yu J, et al. Exercise training in patients with severe chronic heart failure: impact on left ventricular performance and cardiac size. A retrospective analysis of the Leipzig Heart Failure Training Trial. *Eur J Cardiovasc Prev Rehabil.* 2003 Oct;10(5):336-44. doi: 10.1097/01.hjr.0000099031.38268.27.
15. Freimark D, Adler Y, Feinberg MS, Regev T, Rotstein Z, Eldar M, et al. Impact of left ventricular filling properties on the benefit of exercise training in patients with advanced chronic heart failure secondary to ischemic or nonischemic cardiomyopathy. *Am J Cardiol.* 2005;95(1):136-40. doi: 10.1016/j.amjcard.2004.08.081.

16. Medeiros A, Rolim NP, Oliveira RS, Rosa KT, Mattos KC, Casarini DE, et al. Exercise training delays cardiac dysfunction and prevents calcium handling abnormalities in sympathetic hyperactivity-induced heart failure mice. *J Appl Physiol* (1985). 2008 Jan;104(1):103-9. doi: 10.1152/japplphysiol.00493.2007.
17. Vanzelli AS, Medeiros A, Rolim N, Bartholomeu JB, Cunha TF, Bechara LR, et al. Integrative effect of carvedilol and aerobic exercise training therapies on improving cardiac contractility and remodeling in heart failure mice. *PLoS One*. 2013;8(5):e62452. doi: 10.1371/journal.pone.0062452.
18. Ferreira JC, Rolim NP, Bartholomeu JB, Gobatto CA, Kokubun E, Brum PC. Maximal lactate steady state in running mice: effect of exercise training. *Clin Exp Pharmacol Physiol*. 2007;34(8):760-5. doi: 10.1111/j.1440-1681.2007.04635.x.
19. Natali AJ, Turner DL, Harrison SM, White E. Regional effects of voluntary exercise on cell size and contraction-frequency responses in rat cardiac myocytes. *J Exp Biol*. 2001;204(Pt 6):1191-9. PMID: 11222134.
20. Carneiro-Junior MA, Quintao-Junior JF, Drummond LR, Lavorato VN, Drummond FR, da Cunha DN, et al. The benefits of endurance training in cardiomyocyte function in hypertensive rats are reversed within four weeks of detraining. *J Mol Cell Cardiol*. 2013 Apr;57:119-28. doi: 10.1016/j.yjmcc.2013.01.013.
21. Ueta CB, Fernandes GW, Capelo LP, Fonseca TL, Maculan FD, Gouveia CH, et al. beta(1) Adrenergic receptor is key to cold- and diet-induced thermogenesis in mice. *J Endocrinol*. 2012;214(3):359-65. doi: 10.1530/JOE-12-0155.
22. Soloveva V, Graves RA, Rasenick MM, Spiegelman BM, Ross SR. Transgenic mice overexpressing the beta 1-adrenergic receptor in adipose tissue are resistant to obesity. *Mol Endocrinol*. 1997;11(1):27-38. doi: 10.1210/mend.11.1.9870.
23. Lafontan M, Berlan M. Fat cell adrenergic receptors and the control of white and brown fat cell function. *J Lipid Res*. 1993;34(7):1057-91. PMID: 8371057.
24. Allen DL, Harrison BC, Maass A, Bell ML, Byrnes WC, Leinwand LA. Cardiac and skeletal muscle adaptations to voluntary wheel running in the mouse. *J Appl Physiol* (1985). 2001;90(5):1900-8. PMID: 11299284.
25. Kaplan ML, Cheslow Y, Vikstrom K, Malhotra A, Geenen DL, Nakouzi A, et al. Cardiac adaptations to chronic exercise in mice. *Am J Physiol*. 1994;267(3 Pt 2):H1167-73. PMID: 8092282.
26. Kemi OJ, Loennechen JP, Wisloff U, Ellingsen O. Intensity-controlled treadmill running in mice: cardiac and skeletal muscle hypertrophy. *J Appl Physiol* (1985). 2002;93(4):1301-9. doi: 10.1152/japplphysiol.00231.2002.
27. Moore RL, Korzick DH. Cellular adaptations of the myocardium to chronic exercise. *Prog Cardiovasc Dis*. 1995;37(6):371-96. PMID: 7777668.
28. Chruscinski AJ, Rohrer DK, Schauble E, Desai KH, Bernstein D, Kobilka BK. Targeted disruption of the beta2 adrenergic receptor gene. *J Biol Chem*. 1999;274(24):16694-700. PMID: 10358008.
29. Rohrer DK, Chruscinski A, Schauble EH, Bernstein D, Kobilka BK. Cardiovascular and metabolic alterations in mice lacking both beta1- and beta2-adrenergic receptors. *J Biol Chem*. 1999;274(24):16701-8. PMID: 10358009.
30. Boyda HN, Procyshyn RM, Pang CC, Barr AM. Peripheral adrenoceptors: the impetus behind glucose dysregulation and insulin resistance. *J Neuroendocrinol*. 2013;25(3):217-28. doi: 10.1111/jne.12002.
31. Bristow MR, Ginsburg R, Umans V, Fowler M, Minobe W, Rasmussen R, et al. Beta 1- and beta 2-adrenergic-receptor subpopulations in nonfailing and failing human ventricular myocardium: coupling of both receptor subtypes to muscle contraction and selective beta 1-receptor down-regulation in heart failure. *Circ Res*. 1986;59(3):297-309. PMID: 2876788.
32. Zhu WZ, Zheng M, Koch WJ, Lefkowitz RJ, Kobilka BK, Xiao RP. Dual modulation of cell survival and cell death by beta(2)-adrenergic signaling in adult mouse cardiac myocytes. *Proc Natl Acad Sci U S A*. 2001;98(4):1607-12. doi: 10.1073/pnas.98.4.1607.
33. Rohrer DK, Desai KH, Jasper JR, Stevens ME, Regula DP Jr, Barsh GS, et al. Targeted disruption of the mouse beta1-adrenergic receptor gene: developmental and cardiovascular effects. *Proc Natl Acad Sci U S A*. 1996;93(14):7375-80. PMID: 8693001.
34. Zhu WZ, Chakir K, Zhang S, Yang D, Lavoie C, Bouvier M, et al. Heterodimerization of beta1- and beta2-adrenergic receptor subtypes optimizes beta-adrenergic modulation of cardiac contractility. *Circ Res*. 2005;97(3):244-51. doi: 10.1161/01.RES.0000176764.38934.86
35. Zhou YY, Yang D, Zhu WZ, Zhang SJ, Wang DJ, Rohrer DK, et al. Spontaneous activation of beta(2)- but not beta(1)-adrenoceptors expressed in cardiac myocytes from beta(1)beta(2) double knockout mice. *Mol Pharmacol*. 2000;58(5):887-94. PMID: 11040034.
36. O'Connell TD, Jensen BC, Baker AJ, Simpson PC. Cardiac alpha1-adrenergic receptors: novel aspects of expression, signaling mechanisms, physiologic function, and clinical importance. *Pharmacol Rev*. 2014;66(1):308-33. doi: 10.1124/pr.112.007203.
37. Korzick DH, Hunter JC, McDowell MK, Delp MD, Tickerhoof MM, Carson LD. Chronic exercise improves myocardial inotropic reserve capacity through alpha1-adrenergic and protein kinase C-dependent effects in Senescent rats. *J Gerontol A Biol Sci Med Sci*. 2004;59(11):1089-98. PMID: 15602054.
38. Korzick DH, Moore RL. Chronic exercise enhances cardiac alpha 1-adrenergic inotropic responsiveness in rats with mild hypertension. *Am J Physiol*. 1996;271(6 Pt 2):H2599-608. PMID: 8997321.
39. Beaulieu M, Brakier-Gingras L, Bouvier M. Upregulation of alpha1A- and alpha1B-adrenergic receptor mRNAs in the heart of cardiomyopathic hamsters. *J Mol Cell Cardiol*. 1997;29(1):111-9. doi: 10.1006/jmcc.1996.0256.
40. Milligan G, Svoboda P, Brown CM. Why are there so many adrenoceptor subtypes? *Biochem Pharmacol*. 1994;48(6):1059-71. PMID: 7945399.
41. Kemi OJ, Haram PM, Loennechen JP, Osnes JB, Skomedal T, Wisloff U, et al. Moderate vs. high exercise intensity: differential effects on aerobic fitness, cardiomyocyte contractility, and endothelial function. *Cardiovasc Res*. 2005;67(1):161-72. doi: 10.1016/j.cardiores.2005.03.010.
42. Kemi OJ, Ellingsen O, Ceci M, Grimaldi S, Smith GL, Condorelli G, et al. Aerobic interval training enhances cardiomyocyte contractility and Ca2+ cycling by phosphorylation of CaMKII and Thr-17 of phospholamban. *J Mol Cell Cardiol*. 2007 Sep;43(3):354-61. doi: 10.1016/j.yjmcc.2007.06.013.
43. Kemi OJ, Ceci M, Condorelli G, Smith GL, Wisloff U. Myocardial sarcoplasmic reticulum Ca2+ ATPase function is increased by aerobic interval training. *Eur J Cardiovasc Prev Rehabil*. 2008 Apr;15(2):145-8. doi: 10.1097/HJR.0b013e3282ef4e0.



This is an open-access article distributed under the terms of the Creative Commons Attribution License