

UNIVERSIDADE FEDERAL DE VIÇOSA

Effects of liquid delactosed permeate in nursery piglet diets: improvements in feed conversion with increased post-weaning diarrhea occurrence

Hellen Lazarino Oliveira Vilela
Magister Scientiae

**VIÇOSA - MINAS GERAIS
2025**

HELLEN LAZARINO OLIVEIRA VILELA

Effects of liquid delactosed permeate in nursery piglet diets: improvements in feed conversion with increased post-weaning diarrhea occurrence

Dissertation submitted to the Animal Science Graduate Program of the Universidade Federal de Viçosa in partial fulfillment of the requirements for the degree of *Magister Scientiae*.

Adviser: Jansller Luiz Genova

Co-adviser: Gabriel Cipriano Rocha

**VIÇOSA - MINAS GERAIS
2025**

**Ficha catalográfica elaborada pela Biblioteca Central da Universidade
Federal de Viçosa - Campus Viçosa**

T

V699e
2025 Vilela, Hellen Lazarino Oliveira, 1999-
Effects of liquid delactosed permeate in nursery piglet diets
: improvements in feed conversion with increased post-weaning
diarrhea occurrence / Hellen Lazarino Oliveira Vilela. – Viçosa,
MG, 2025.

1 dissertação eletrônica (42 f.)

Texto em inglês.

Orientador: Jansller Luiz Genova.

Dissertação (mestrado) - Universidade Federal de Viçosa,
Departamento de Zootecnia, 2025.

Referências bibliográficas: f. 38-42.

DOI: <https://doi.org/10.47328/ufvbbt.2025.242>

Modo de acesso: World Wide Web.

1. Suínos - Alimentação e rações - Efeito da lactose.
2. Suínos - Registros de desempenho. I. Genova, Jansller Luiz,
1991-. II. Universidade Federal de Viçosa. Departamento de
Zootecnia. Programa de Pós-Graduação em Zootecnia.
III. Título.

CDD 22. ed. 636.40855

HELLEN LAZARINO OLIVEIRA VILELA

Effects of liquid delactosed permeate in nursery piglet diets: improvements in feed conversion with increased post-weaning diarrhea occurrence

Dissertation submitted to the Animal Science Graduate Program of the Universidade Federal de Viçosa in partial fulfillment of the requirements for the degree of *Magister Scientiae*.

APPROVED: February 19, 2025.

Assent:

Hellen Lazarino Oliveira Vilela
Author

Jansller Luiz Genova
Adviser

Essa dissertação foi assinada digitalmente pela autora em 06/05/2025 às 19:38:54 e pelo orientador em 07/05/2025 às 08:56:16. As assinaturas têm validade legal, conforme o disposto na Medida Provisória 2.200-2/2001 e na Resolução nº 37/2012 do CONARQ. Para conferir a autenticidade, acesse <https://siadoc.ufv.br/validar-documento>. No campo 'Código de registro', informe o código **I5SQ.85CK.WY4V** e clique no botão 'Validar documento'.

I dedicate this work to the memory of my beloved grandmother, Dona Maria da Conceição, whose wisdom, strength and unconditional love were inexhaustible sources of inspiration throughout my life. Her example of perseverance, dedication and hard work taught me never to give up on my dreams, even in the face of challenges. This is more than a title, it is the realization of a dream that she, with all her strength and example, helped to build

ACKNOWLEDGMENTS

This work has been sponsored by the following Brazilian research agencies: Coordination for the Improvement of Higher Education Personnel (CAPES; Financing code 001), Minas Gerais State Foundation for Research Aid (FAPEMIG) and National Council of Scientific and Technological Development (CNPq).

First of all, I thank God, who gave me the strength, wisdom and courage to walk this path. My faith was the foundation that supported each step, especially in the most challenging moments, always reminding me that no obstacle is greater than God's will in our lives.

To my family, my foundation and safe haven, who always believed in me, even when the uncertainties seemed greater than the certainties.

I couldn't fail to mention the women in my life: Ernestina, Eliane and Maria (in memoriam). Each one, in their own way, taught me about love, resilience and the importance of never giving up. You are a fundamental part of this achievement.

To the memory of my aunt Arlinda, whose dedication to her studies has always been an inspiration to me. Even though she was physically absent, her story and example are still alive and present in my every achievement.

To my aunt Elane, for taking me in during a challenging period in my life. Your support and affection were fundamental in helping me to move forward with courage and determination.

To my soul friend, Damares, who has always been by my side in times of joy and also in times of challenge, with words of support and affection that have made all the difference in this journey.

To my dear sorority friends, Julya, Isabella, Sthephanny and Luana. Living with you made this phase lighter and more fun. The moments of sharing, laughter and mutual support will remain in my memory forever

To my friends Ruan, Danilo and Laynne, who have always been one of the best fruits that Animal Science has given me.

To my teammates, Pedro, Rayanne, Paloma and Marina, your collaboration and companionship have been fundamental to my professional and personal growth.

To Professor Paulo Levi, for his patience and valuable contributions throughout my academic career. Your dedication and encouragement were fundamental to my development.

To the team of postgraduate students at UNIOESTE, for carrying out this

work and for their collaboration, which allowed me to share countless experiences.

To my advisor, Jansller, and co-advisor, Gabriel, for their trust, guidance and continuous encouragement. Your guidance was essential to the development of this work, and I am grateful for all the learning and support I received from you.

To my in-laws, Adriano and Neusa, for their support, welcome and for making me feel part of the family. Your support was fundamental in enabling me to dedicate myself to my studies with peace of mind.

To my boyfriend, Gabriel, for being by my side every step of the way. Your patience, understanding and unconditional support were essential for me to overcome the challenges. Thank you for believing in me even when I doubted, and for being my partner in every way.

To the Coordination for the Improvement of Higher Education Personnel (CAPES), for awarding the scholarship.

To everyone who has been part of this journey in any way, my sincere thanks. This work is the result not only of my efforts, but also of the support and inspiration of each and every one of you.

ABSTRACT

VILELA, Hellen Lazarino Oliveira, M.Sc., Universidade Federal de Viçosa, February, 2025. **Effects of liquid delactosed permeate in nursery piglet diets: improvements in feed conversion with increased post-weaning diarrhea occurrence.** Adviser: Jansller Luiz Genova. Co-adviser: Gabriel Cipriano Rocha.

The objective of this study was to assess the effects of liquid delactosed permeate (DLP) alone or mixed as baby feed with a conventional lactose-containing diet for nursery piglets on growth performance, diarrhea occurrence (DO), blood profile, gastrointestinal tract traits, and bacterial population counts. Ninety male hybrid piglets (7.59 ± 0.63 kg) weaned at 25 days old (day 0) were assigned for 35 days in a complete block design based on body weight (BW) to one of three treatments: control diet (CD) containing lactose via whey permeate powder, supplied in mash and dry form, without DLP; supplemented diet (SD): CD + DLP provided in specific feeders and separated from the mash diet; moist diet (MD): CD moistened with DLP in a 1:1 ratio, 10 replicates, and three piglets per pen. Nursery phases were defined as pre-starter I and II (days 0 to 7, and 8 to 21) and starter (day 22 to 35). Piglets fed MD diet had better feed conversion (FCR) compared with those fed CD in the pre-starter I and II and total period. Piglets in pre-starter I fed MD diet had a higher DO compared with those fed CD, whereas piglets fed SD had intermediate results. Piglets in pre-starter II fed SD had a higher total protein concentration than those fed MD, and CD provided intermediate results. Piglets on MD had higher leukocyte concentrations compared with those fed CD, but piglets fed SD had intermediate results. Piglets fed CD or SD diets had higher colonic content pH than those fed MD. Piglets fed SD or MD diets had longer large intestine lengths than animals fed CD. Delactosed permeate improves FCR in piglets in the pre-starter phases and promotes minor changes in other variables; however, a greater DO was observed in the first few days post-weaning

Keywords: bacterial population; blood profile; dairy co-products; diarrhea prevalence; growth performance; nursery pig

RESUMO

VILELA, Hellen Lazarino Oliveira, M.Sc., Universidade Federal de Viçosa, fevereiro de 2025. **Efeitos do permeado líquido deslactosado em dietas de leitões de creche: Melhorias na conversão alimentar com aumento da ocorrência de diarreia pós-desmame.** Orientador: Jansller Luiz Genova. Coorientador: Gabriel Cipriano Rocha.

O objetivo deste estudo foi o de avaliar os efeitos do permeado deslactosado líquido (PDL) sozinho ou misturado como “papinha” com uma dieta convencional contendo lactose para leitões de creche sobre o desempenho zootécnico, a ocorrência de diarreia (OD), o perfil sanguíneo, as características do trato gastrointestinal e as contagens de população bacteriana. Noventa leitões híbridos machos ($7,59 \pm 0,63$ kg) desmamados aos 25 dias de idade (dia 0) foram designados por 35 dias em um delineamento de blocos completos baseado no peso corporal (PC) para um dos três tratamentos: dieta controle (DC) contendo lactose via permeado de soro de leite em pó, fornecido na forma farelada e seca, sem PDL; dieta suplementada (DS): DC + PDL fornecido em alimentadores específicos e separado da dieta farelada; dieta úmida (DU): DC umedecida com PDL na proporção de 1:1, 10 repetições e três leitões por baía. As fases de creche foram definidas como pré-inicial I e II (dias 0 a 7 e 8 a 21) e inicial (dia 22 a 35). Leitões alimentados com a dieta DU tiveram melhor conversão alimentar (CA) em comparação com àqueles alimentados com DC na pré-inicial I e II e no período total. Leitões na pré-inicial I alimentados com dieta DU tiveram maior OD em comparação com àqueles alimentados com DC, enquanto leitões alimentados com DS tiveram resultados intermediários. Leitões na pré-inicial II alimentados com DS tiveram uma concentração de proteína total maior do que àqueles alimentados com DU, e a DC forneceu resultados intermediários. Leitões com DU tiveram maiores concentrações de leucócitos em comparação com àqueles alimentados com DC, mas leitões alimentados com DS tiveram resultados intermediários. Leitões alimentados com dietas DC ou DS tiveram mai

Palavras-chave: população bacteriana; perfil sanguíneo; co-produtos lácteos; prevalência de diarreia; desempenho em termos de crescimento; suínos em creche

SUMÁRIO

1. INTRODUCTION	9
2. MATERIAL AND METHODS.....	11
2.1. Local and ethical information	11
2.2. Animals, experimental design, housing, and dietary treatments.....	11
2.3. Chemical composition and supply of delactosed permeate	14
2.4. Growth performance and diarrhea occurrence	16
2.5. Sampling and analysis of blood metabolites	16
2.6. Slaughter procedures and gastrointestinal tract traits.....	17
2.7. pH of tract contents, relative weight, and length of organs	17
2.8. Intestinal morphometry	18
2.9. Bacterial population counts.....	18
2.10. Statistical procedures.....	19
3. RESULTS	21
3.1. Growth performance and diarrhea occurrence.....	21
3.2. Blood metabolites.....	23
3.3. pH of digestive tract contents, length, and relative weight of organs.....	26
3.4. Intestinal morphometry	28
3.5. Bacterial population counts.....	30
4. DISCUSSION.....	32
5. CONCLUSIONS.....	37
REFERENCES.....	38

1. INTRODUCTION

Co-products from the dairy industry can be strategic in diets for newly weaned piglets by maintaining a healthier intestinal environment (Yoo et al., 2018). Due to the considerable increase in the use of liquid feed and fermented liquids in pig diets (Zhang et al., 2020), these co-products are alternatives to expensive and high-quality ingredients (Sutera et al., 2023). Delactosed permeate (DLP) is a co-product of the lactose manufacturing process. It is considered an economic burden for processing companies, and its disposal poses environmental concerns due to its high organic matter content and treatment costs; besides, it is a product of negligible value and, therefore, there is an opportunity to use this dairy co-product in animal feed (Oliveira et al., 2019).

Delactosed permeate has a high concentration of minerals (ADPI, 2015) and oligosaccharides (3'-sialyllactose, 6'-sialyl-lactose, and lactosamine) (Dallas et al., 2014), which have functional properties for the gastrointestinal tract (GIT) health (Barile et al., 2009). For example, the effects include inhibiting binding of pathogenic bacteria (e.g., Enterobacteriaceae) and improving growth of beneficial bacteria (e.g., Lactobacillaceae) in the small intestine (Yu et al., 2013), stimulating the immunity of the intestinal mucosa and enterocyte proliferation (Jang et al., 2021a), and producing short-chain fatty acids as an energy source for intestinal cells (Venema, 2012). In addition, diets moistened with dairy co-products (e.g., baby feed) can improve piglet performance by increasing feed intake and reducing the dehydration risk (Russell et al., 1996) and sudden dietary changes (Lepine et al., 1991).

Jang et al. (2021a) obtained better results in feed efficiency, crypt cell proliferation rate, and bacterial modulation when 45.9 g of lactose day⁻¹ was given to piglets weaned at 21 days old, while Jang et al. (2021b) observed that reducing the daily intake of lactose from 63 to 43 g day⁻¹ in diets for piglets weaned at 25 days old was positive. Pierce et al. (2006) observed that 150 g of lactose kg⁻¹ diet increased villus height and reduced intestinal pH, but 330 g of lactose kg⁻¹ diet increased Lactobacillus in piglets weaned at 21 days old.

The form of supply, the inclusion level of these co-products, and the response to lactose content according to weaning age for optimal piglet performance are a theme of debate (Fondevila et al., 2021). The ingestion and excessive lactose fermentation can cause intolerance symptoms, resulting in diarrhea (Forsgård, 2019). This study hypothesizes that weaned piglets fed DLP would improve growth performance through benefits in intestinal health. Therefore, the present study assessed the effects of DLP fed alone or mixed as baby

feed with a conventional lactose-containing diet for nursery piglets on performance, diarrhea occurrence (DO), blood profile, GIT traits, and bacterial population counts.

2. MATERIAL AND METHODS

2.1. Local and ethical information

The study was conducted in an experimental unit located in Marechal Cândido Rondon, PR, Brazil (24°31'52" S and 54°01'03" W). Research on animals was conducted according to the institutional committee on animal use (protocol no. P026/22).

2.2. Animals, experimental design, housing, and dietary treatments

The experiment lasted 35 days, with no adaptation period. A total of 90 entire male hybrid piglets of commercial lineage (Landrace × Large White), weaned at 25 days old (day 0) and weighing 7.59 ± 0.63 kg, were allotted to one of three treatments in a completely block design based on the initial body weight (BW) categorized into light (up to 7.57 kg BW) or heavy (over 7.57 kg BW) piglets. Each treatment had 10 replicates pens (n = five replicate pens per block) of three piglets of the same BW category per pen as experimental unit. The blocks (n = 2) consisted of 15 pens (experimental unit) and 45 piglets in each block (15 piglets/treatment), totaling 30 pens.

On arrival at the facility, the piglets were weighed and individually identified with numbered earrings and housed in a masonry installation with ceramic tiles and a plastic tarpaulin lining. The elevated pens (1.5 × 1.03 m, 1.54 m²) were equipped with plastic slatted floor, trough-type feeders, and pacifier-type drinkers, arranged in two rows divided by a central aisle. The pens and the facility were cleaned daily in the afternoon. The ambient temperature and relative humidity were recorded daily in the morning (08:00 and 10:30 h) and afternoon (14:00 and 16:30 h) using a datalogger (UT330B digital USB; UNI-T, Vketech, Beijing, China) installed in the central part of the facility.

Fans, exhaust fans, and side-hinged glass windows controlled ventilation inside the facility. Individual incandescent and infrared lamps located at the top of each pen were used to heat the environment. An automation system for the heating lamps and camera monitoring controlled via Wi-Fi was used to help control temperature and humidity and observe the animals during the experimental period. A combination of daylight and artificial light was used, with a 12-h light-dark cycle.

The dietary treatments (Table 1) consisted of: control diet (CD) containing lactose via whey permeate powder, supplied in mash and dry form, without liquid DLP; supplemented diet (SD): CD + DLP provided in specific feeders (e.g., polyethylene plastic piglet feeder attached to the floor) and separated from the mash diet; and moist diet (MD): CD moistened with DLP in a 1:1 ratio. All the diets were corn and soybean meal-based with industrial amino acids, and mixed for 15 min in a vertical mixer. The CD was formulated to meet the nutritional and energy recommendations for piglets in the nursery phase, being pre-starter I (0 to 7 days) and II (8 to 21 days), and starter (22 to 35 days) (Rostagno et al., 2017).

Table 1 - Calculated and chemical composition of the experimental diets offered to piglets in the nursery phase (as-fed basis)

Item	Experimental phase		
	Pre-starter I	Pre-starter II	Starter
Ingredient (%)			
Ground corn (7.8%)	40.34	44.26	61.09
Soybean meal (45.4%)	21.69	20.65	24.21
Whole deactivated soybean (37.3%)	10.00	10.00	6.00
Whey powder permeate (3.0%)	14.02	9.41	-
Cooking sugar	5.00	5.00	-
Fish meal (53.0%)	3.00	3.00	3.00
Soybean oil	1.63	3.00	1.62
Dicalcium phosphate	1.48	1.57	1.54
Calcitic limestone	0.89	0.692	0.669
Common salt	0.11	0.37	0.42
Mineral-vitamin premix ¹	0.30	0.30	0.30
Lysine sulphate (54.6%)	0.71	0.89	0.57
DL-methionine (99.5%)	0.36	0.27	0.23
L-threonine (96.8%)	0.42	0.34	0.27
L-tryptophan (99.0%)	0.09	0.05	0.05
L-valine (95.5%)	-	0.182	-
Halquinol (60.0%)	0.01	0.01	0.01
Calculated chemical composition			
Crude protein (%)	21.42	19.87	20.55
Metabolizable energy (kcal kg ⁻¹)	3,400	3,380	3,350
Total calcium (%)	1.07	0.97	0.91
Available phosphorus (%)	0.52	0.48	0.45
Lactose (%)	11.00	8.00	-
Total sodium (%)	0.22	0.21	0.20
Digestible lysine (%)	1.45	1.35	1.28
Digestible methionine + cysteine (%)	0.81	0.75	0.73
Digestible threonine (%)	0.97	0.90	0.83
Digestible tryptophan (%)	0.27	0.25	0.24
Digestible valine (%)	1.00	0.92	0.88

¹Mineral premix (content per kg of product): zinc oxide, 26.66 g; iron sulphate, 10 g; copper sulphate, 2,708 mg; iodine, 333 mg; biotin, 26.66 mg; manganese sulphate, 10 g; selenium (sodium selenite), 100 mg. Vitamin premix (content per kg of product): vitamin A, 2,000,000 IU; vitamin D₃, 533,330 IU; vitamin E, 6,583 IU; vitamin K₃, 1,066 mg; vitamin B₁, 334 mg; vitamin B₂, 1,333 mg; vitamin B₆, 466 mg; vitamin B₁₂, 5,333 mcg; niacin, 9,372 mg; pantothenic acid, 5,022 mg; folic acid, 167 mg

2.3. Chemical composition and supply of delactosed permeate

The DLP (Table 2) was manufactured by Sooro Renner Nutrição S.A. (Marechal Cândido Rondon, PR, Brazil). The co-product was packed in plastic bottles to facilitate handling and kept refrigerated (4 °C) for a maximum of seven days. It was given to the animals in the SD or MD treatment in the morning (08:00 h) and afternoon (14:00 h) periods. The amount offered to the piglets in the SD treatment was altered in each nursery phase: pre-starter I (morning: 200 mL; afternoon: 250 mL), pre-starter II (morning: 275 mL; afternoon: 275 mL), and starter (morning: 400 mL; afternoon: 400 mL). The ratio (1:1) of CD to DLP given to the piglets in the MD treatment was altered based on the total daily feed intake of the animals in each phase. Before feeding, the CD was weighed and mixed with the DLP to ensure the animals' total intake in each of the two periods.

Table 2 - Chemical composition of liquid delactosed permeate

Item	Content
Dry matter (%)	37.00
Total solids (%)	28.57 (68.25% de lactose)
Ash (%)	4.46
Sodium (mg/100 g)	371.18
Potassium (mg/100 g)	499.06
Chlorides (%)	0.80
Crude protein (%)	2.44
Ethereal extract (%)	0.00
Carbohydrates (%)	21.67
Lactose (%)	19.50
Galactose (%)	1.08
Glicose (%)	1.08
pH	5.56
Acidity (°D)	105.88

2.4. Growth performance and diarrhea occurrence

The animals were fed ad libitum and had free access to water throughout the experiment. The piglets were weighed at the beginning and end of each phase using a digital scale (Model UL-50, Brand DIGI-TRON, Curitiba, PR, Brazil) to monitor final BW (FBW) and calculate average daily BW gain (ADG, kg day^{-1}). The diets and leftovers in the feeder and on the floor were collected manually, weighed daily, and deducted from the total supply to calculate the average daily feed intake (ADFI, kg day^{-1}) and feed conversion ratio (FCR, kg kg^{-1}). In the case of leftovers from the MD treatment at the change of each phase, these were dried in a forced-air circulation oven (Tecnal, SF-325 NM; Piracicaba, SP, Brazil) and then weighed. The intake of DLP provided alone was not considered in the ADFI of piglets fed the SD.

Diarrhea occurrence was assessed throughout the experimental period. The fecal consistency of the animals was observed daily (08:00 h) at pen level by a single trained person through visual analysis, assigning scores from 0 to 3 according to the physical appearance of the feces, in which 0 = normal feces, 1 = soft feces, 2 = watery/pasty feces, and 3 = liquid feces (Pérez-Calvo et al., 2019). Scores 2 and 3 were considered diarrheal feces.

2.5. Sampling and analysis of blood metabolites

Blood samples ($\cong 10$ mL, 08:30 h) were taken at the end of the pre-starter II (day 21) and starter (day 35) phases. All the piglets were fasted for 6 h, and 10 piglets per treatment (one animal per experimental unit) were chosen at the end of each phase for blood collection based on the BW closest to the average BW of their respective pen. The collection procedure was carried out via puncture of the anterior cranial vena cava, using 20 mL plastic syringes and 0.70×30 mm gauge needles. The same animals were used for both blood samples. The blood was then transferred to four sterile glass tubes containing a type of anticoagulant (heparin) for analysis of urea (heparin), complete blood count (EDTA), glucose (potassium fluoride), and albumin, total cholesterol, triacylglycerols, and total protein (serum).

The tubes containing the collected samples were placed in a thermal box containing ice (4°C) and sent to the laboratory for analysis. The blood samples were centrifuged (analog centrifuge, model 80-2B, Centrilab, Maringá, PR, Brazil) at 3,000 rpm for 10 min at room

temperature. Approximately 3 mL of the supernatant was then transferred to Eppendorf-type polyethylene microtubes, duly labeled, and stored in a freezer at $-20\text{ }^{\circ}\text{C}$.

The analytical procedures for the analysis of albumin (enzymatic-colorimetric method), glucose (enzymatic-colorimetric method, Trinder), total protein (colorimetric-biuret method), and total cholesterol, triacylglycerols, and urea (enzymatic-colorimetric method) were carried out in technical duplicate by spectrophotometry (Bel SPECTRO S05, Ramos, RJ, Brazil), using specific kits from the Gold Analisa Diagnostic[®] brand (Belo Horizonte, MG, Brazil).

To assess the complete blood count, the blood samples were sent to the Lotus Veterinary Laboratory (Cascavel, PR, Brazil) for quantification of red blood cells, hemoglobin, hematocrit, mean corpuscular volume, red blood cell distribution width, mean corpuscular hemoglobin concentration, plasma proteins, platelets, total leukocytes, segmented neutrophils, eosinophils, lymphocytes, and monocytes.

2.6. Slaughter procedures and gastrointestinal tract traits

At the end of the experimental period (day 35), seven animals from each treatment were slaughtered based on the BW closest to the average BW of their respective pen. The animals were not the same blood donors. They were fasted for 8 h and electrically stunned (240 volts for 3 s), followed by bleeding. Slaughter was carried out in a commercial slaughterhouse (Marechal Cândido Rondon, PR, Brazil).

2.7. pH of tract contents, relative weight, and length of organs

The viscera were exposed through a central incision. The GIT portions were aseptically isolated with a double ligature, and the pH was measured in the homogenized contents of the medial region of the stomach, jejunum (150 cm from the ileocecal junction), ileum (15 cm from the ileocecal junction), cecum, and medial colon using a digital pH meter (HI 99163, Hanna Instruments Inc., Rhodes Island, USA). After measuring the pH, the organs of the GIT (stomach, small intestine, cecum, empty large intestine, and liver with gallbladder) were weighed (digital scale, model 9094, Toledo, Brazil) to calculate the relative weight of the organs based on the BW of the animals at slaughter. A common tape measure was used to measure the length of the small and intestines.

2.8. Intestinal morphometry

The fragments (\cong 3 cm long) of the jejunum and ileum of all the slaughtered animals were collected, washed with distilled water and physiological solution (0.9% sodium chloride), and then stored in 50-mL sterile plastic containers containing a prepared solution of 10.0% neutral buffered formalin (100 mL of 37.5% formaldehyde, 900 mL of distilled water, 4 g of monobasic sodium phosphate, and 6.5 g of dibasic sodium phosphate) for 48 h (Genova et al., 2021). The samples were then placed in a 70.0% alcohol solution and sent to a commercial laboratory (Mercolab, Cascavel, PR, Brazil). The samples were washed in saline solution, fixed in Bouin's solution, dehydrated in a gradual series of ethanol, and cleaned with xylene. The tissues were embedded in paraffin blocks to prepare the slides. Five-micrometer sections were cut from the paraffin blocks using a rotary microtome (Leica RM2245, Leica Biosystems, São Paulo, SP, Brazil), for subsequent staining with hematoxylin and eosin.

The images of the slides were captured by a digital camera (Pro Series, Silver Spring, MD, USA) coupled to an Olympus BX40 microscope (Olympus, Tokyo, Japan). The morphometric examination of the images was carried out using imaging software (Image-Pro plus 4.1, Media Cybernetics[®], Silver Spring, Maryland). Ten measurements of villus height (V) and their respective crypt depths (C) in each intestinal fragment (jejunum and ileum) were evaluated to calculate the average value for each animal and then obtain the V:C ratio.

2.9. Bacterial population counts

After evisceration, the jejunal and ileal mucosa were exposed using a scalpel. Samples (\cong 5 g) of the contents of the jejunum and ileum of the piglets were collected using the technique of bacterial isolation, cultivation, counting, and characterization by the traditional plating method. The samples were analyzed according to the procedures of ISO 16649-2:2001 (ISO, 2006). Then, they were immediately packed in sterile plastic containers, identified, and transported under refrigeration (4 °C) for laboratory analysis (Mercolab, Cascavel, PR, Brazil).

In the commercial laboratory, 1 g of each sample was serially diluted in 1.0% peptone water. Each dilution was vortexed (Phoenix, model AP 56; Araraquara, SP, Brazil) for 30 s. An aliquot of 100 μ L of each diluted sample was spread evenly on the surface of the plates containing specific agar for each bacterium. The microbial populations evaluated were

Escherichia coli (eosin methylene blue levine agar, Kasvi brand; São José dos Pinhais, PR, Brazil) and *Lactobacillus* (Man, Rogosa and Sharpe agar, Acumedia brand; São Paulo, SP, Brazil). The plates were incubated at 37 °C (EL 202; Eletrolab, São Paulo, SP, Brazil) aerobically for 24 h to count *E. coli* and anaerobically for 48 h to count *Lactobacillus* (Genova et al., 2022). The bacterial colonies were counted, and the data multiplied by the respective dilution and transformed into log (log₁₀). The results were expressed in log of colony forming units (CFU) g⁻¹.

2.10. Statistical procedures

Statistical analyses were carried out using the procedures of SAS (Statistical Analysis System, version 9.2). Before the analysis of variance (ANOVA) or covariance (ANCOVA), the data was evaluated for outliers (values greater than or equal to 3.0 standard deviations in absolute value were considered influential). The pen was considered the experimental unit, except for slaughter and blood data, in which the animal was considered the experimental unit. The data was analyzed considering treatment and BW block as fixed and random effects, respectively. Initial BW was included in the model as a covariate when $P < 0.05$, therefore, using ANCOVA. The following overall model was used for the growth performance data:

$$Y_{ijk} = \mu + T_i + b_j + \beta (X_{ijk} - \bar{X}...) + \varepsilon_{ijk}, (1)$$

in which Y_{ijk} = average observation of the dependent variable in each plot measured in the i -th treatment, in the j -th block, and in the k -th replicate; μ = overall average; T_i = fixed effect of treatment ($i = 1, 2, \text{ and } 3$); b_j = random effect of block ($j = 1 \text{ and } 2$); β = regression coefficient of Y over X ; X_{ijk} = average observation of the covariate (initial BW) in each plot, measured in the i -th treatment, in the j -th block, and in the k -th replicate; $\bar{X}...$ = overall average for the covariate X ; ε_{ijk} = random error of the plot associated with each Y_{ijk} observation. Other variables were analyzed using the model mentioned above without the covariate effect.

Bacterial count data was transformed into log₁₀ CFU g⁻¹ for analysis. The treatment effect on the dependent variables was assessed using ANOVA or ANCOVA. Duncan's post hoc test was used to compare means between dietary treatments; it was used because the present study required that type-II error be avoided and possible effects be not lost, when $P \leq 0.05$. In addition, it had lower values of the minimum significant difference, implying that it was easier to detect differences between treatments if they existed, that is, Duncan's test has higher sensitivity to detect smaller differences between group averages. All data were reported

as means with pooled standard error of the mean.

A generalized linear model was fitted for DO (binomial distribution and logit link function). Diarrhea occurrence results were presented as observed proportions (relative frequency in %). Treatment and time were considered as fixed effects and BW block as a random factor. The treatment effect was verified using type III analysis. Significant differences were defined as $P \leq 0.05$, and DO estimates were compared using a difference between averages test, by the χ^2 statistic.

3. RESULTS

3.1. Growth performance and diarrhea occurrence

The ambient temperature and relative humidity recorded during the experimental period were 24 ± 2.48 °C and $65 \pm 7.73\%$, respectively. Piglets fed MD dietary treatment had better ($P \leq 0.05$) FCR compared with those fed the CD in the pre-starter I, pre-starter II, and total period (Table 3). In addition, piglets fed the SD diet in the pre-starter I also had better ($P \leq 0.05$) FCR than those fed the CD diet, but showed intermediate results in the pre-starter II and in the total period. There was no effect of dietary treatment on growth performance in the starter phase.

Piglets fed the MD diet in pre-starter I had higher ($P \leq 0.05$) DO compared with those fed the CD diet, while the SD diet provided in intermediate results (Table 3). There was no treatment effect on DO in the pre-starter II and starter phases.

Table 3 - Effects of liquid delactosed permeate (DLP) on growth performance and diarrhea occurrence (DO) of nursery piglets¹

Item	Treatment			SEM	P-value
	CD	SD	MD		
Pre-starter I (day 0 to 7)					
IBW (kg)	7.69	7.53	7.57	0.075	-
FBW (kg)	9.42	9.53	9.34	0.113	0.173
ADG (kg)	0.23	0.28	0.25	0.011	0.172
ADFI (kg)	0.31	0.31	0.26	0.011	0.139
FCR (kg:kg)	1.35 ^a	1.10 ^b	1.06 ^b	0.042	0.004
DO (%)	16.00 ^B	14.00 ^{AB}	18.00 ^A	0.111	0.022
Pre-starter II (day 8 to 21)					
FBW (kg)	15.68	16.04	16.36	0.313	0.255
ADG (kg)	0.44	0.47	0.50	0.022	0.244
ADFI (kg)	0.60	0.59	0.60	0.016	0.981
FCR (kg:kg)	1.37 ^a	1.25 ^{ab}	1.20 ^b	0.013	0.003
DO (%)	9.29	8.57	7.14	0.051	0.237
Starter (day 22 to 35)					
FBW (kg)	24.91	25.69	25.14	0.427	0.289
ADG (kg)	0.65	0.67	0.63	0.010	0.295
ADFI (kg)	1.02	1.04	0.97	0.017	0.210
FCR (kg:kg)	1.57	1.56	1.54	0.024	0.769
DO (%)	1.33	2.00	2.67	0.038	0.176
Total period (day 0 to 35)					
ADG (kg)	0.49	0.51	0.49	0.011	0.211
ADFI (kg)	0.72	0.72	0.67	0.014	0.098
FCR (kg:kg)	1.48 ^a	1.40 ^{ab}	1.37 ^b	0.024	0.044

IBW - initial body weight; FBW - final body weight; ADFI - average daily feed intake; ADG - average daily weight gain; FCR - feed conversion ratio; SEM - pooled standard error of the mean.

¹Data are means of 10 pen replicates per treatment and three animals per pen as the experimental unit.

²Control diet (CD) supplied in dry, mash form, without DLP; supplemented diet (SD): CD + DLP provided in specific feeders and separated from the mash diet; moist diet (MD): CD moistened with DLP in a 1:1 ratio. a,b - Averages followed by different lowercase letters in the row differ according to Duncan's post hoc test ($P \leq 0.05$). A,B - Observed proportions followed by different capital letters in the same row differ using a difference between means test, by the χ^2 statistic.

3.2. Blood metabolites

Regarding the biochemical profile, the piglets in pre-starter II phase that consumed the SD diet had a higher ($P \leq 0.05$) concentration of total protein than the animals fed the MD diet, and the animals fed CD diet showed intermediate results (Table 4). There was no treatment effect on the blood biochemical profile in the starter phase.

Regarding the hematological profile, piglets fed MD diet had higher ($P \leq 0.05$) leukocyte concentrations compared with the animals fed the CD diet, and the animals that received the SD diet had intermediate results (Table 5).

Table 4 - Effects of liquid delactosed permeate (DLP) on the blood biochemical profile of piglets in the nursery phase

Item	Treatment			SEM	P-value
	CD	SD	MD		
Pre-starter II (day 21)					
Albumin (g/dL ⁻¹)	2.02	2.05	1.97	0.035	0.577
Glucose (mg/dL ⁻¹)	115.11	114.21	114.15	3.213	0.975
Total protein (g/dL ⁻¹)	4.70 ^{ab}	4.81 ^a	4.51 ^b	0.045	0.019
Total cholesterol (mg /dL ⁻¹)	73.16	69.02	66.00	1.801	0.271
Triacylglycerols (mg /dL ⁻¹)	57.88	61.42	64.21	2.988	0.694
Urea (mg /dL ⁻¹)	13.16	11.56	9.67	0.898	0.245
Starter (day 35)					
Albumin (g/dL ⁻¹)	2.24	2.35	2.39	0.046	0.150
Glucose (mg/dL ⁻¹)	114.53	114.44	114.77	2.680	0.998
Total protein (g/dL ⁻¹)	4.84	4.79	4.91	0.099	0.712
Total cholesterol (mg/dL ⁻¹)	54.94	55.67	55.09	2.418	0.991
Triacylglycerols (mg/dL ⁻¹)	38.42	39.60	35.05	1.438	0.365
Urea (mg/dL ⁻¹)	22.46	20.37	23.19	1.185	0.577

SEM - pooled standard error of the mean.

¹Data are means of 10 piglets per treatment.

²Control diet (CD) supplied in dry, mash form, without DLP; supplemented diet (SD): CD + DLP provided in specific feeders and separated from the mash diet; moist diet (MD): CD moistened with DLP in a 1:1 ratio. a,b - Averages followed by different lowercase letters in the row differ according to Duncan's post hoc test (P≤0.05)

Table 5 - Effects of liquid delactosed permeate (DLP) on the hematological profile of nursery piglets on day 35¹

Item	Treatment			SEM	P-value
	CD	SD	MD		
Red blood cells ($\times 10^6 \mu\text{L}^{-1}$)	5.93	5.90	5.92	0.073	0.991
Hemoglobin (g dL ⁻¹)	12.01	11.76	11.68	0.122	0.534
Hematocrit (%)	35.15	34.95	34.95	0.375	0.972
MCV (fL)	60.12	60.08	60.08	0.060	0.534
RDW (%)	20.22	19.95	19.25	0.646	0.704
MCHC (g dL ⁻¹)	33.33	33.31	33.29	0.012	0.377
Plasma proteins (g dL ⁻¹)	5.36	5.30	5.35	0.042	0.834
Platelets ($\times 10^3 \mu\text{L}^{-1}$)	347.444	376.750	447.750	19.803	0.153
Leukocytes ($\times 10^3 \mu\text{L}^{-1}$)	19.510 ^b	21.815 ^{ab}	23.040 ^a	700.400	0.052
Segmented (%)	45.50	45.55	44.50	1.065	0.853
Eosinophils (%)	1.10	1.15	1.40	0.184	0.793
Lymphocytes (%)	51.30	52.70	52.25	1.038	0.805
Monocytes (%)	1.60	1.45	1.10	0.182	0.527

MCV - mean corpuscular volume; RDW - red blood cell distribution width; MCHC - mean corpuscular hemoglobin concentration;

SEM - pooled standard error of the mean.

¹Data are means of 10 piglets per treatment.

²Control diet (CD) supplied in dry, mash form, without DLP; supplemented diet (SD): CD + DLP provided in specific feeders and separated from the mash diet; moist diet (MD): CD moistened with DLP in a 1:1 ratio.

a,b - Averages followed by different lowercase letters in the row differ according to Duncan's post hoc test ($P \leq 0.05$).

3.3. pH of digestive tract contents, length, and relative weight of organs

The piglets that consumed the CD or SD diets had higher ($P \leq 0.05$) pH values of the colonic content compared with animals fed the MD diet (Table 6). Piglets fed the SD or MD diets had longer ($P \leq 0.05$) large intestine lengths than the animals fed the CD diet. There was no treatment effect on relative organ weight.

Table 6 - Effects of liquid delactosed permeate (DLP) on the pH of digestive tract contents, and length and relative weight of organs in nursery piglets on day 35

Item	Treatment			SEM	P-value
	CD	SD	MD		
	pH				
Stomach	3.85	3.38	3.41	0.138	0.106
Jejunum	5.77	5.62	5.53	0.154	0.790
Ileum	5.18	5.70	5.24	0.179	0.503
Cecum	5.38	5.44	5.35	0.035	0.370
Colon	5.80 ^a	5.89 ^a	5.55 ^b	0.063	0.001
	Length (cm)				
Small intestine	52.35	50.56	52.23	1.508	0.938
Large intestine	12.98 ^b	14.02 ^a	14.17 ^a	0.249	0.057
	Relative weight (% of body weight)				
Empty stomach	0.76	0.74	0.75	0.017	0.266
Empty small intestine	3.66	3.88	3.92	0.0930	0.157
Empty cecum	9.27	0.28	0.25	0.013	0.970
Empty large intestine	1.68	1.74	1.72	0.041	0.497
Liver with gallbladder	2.78	3.07	2.87	0.067	0.158

SEM - pooled standard error of the mean.

¹Data are means of seven piglets per treatment.

²Control diet (CD) supplied in dry, mash form, without DLP; supplemented diet (SD): CD + DLP provided in specific feeders and separated from the mash diet; moist diet (MD): CD moistened with DLP in a 1:1 ratio. a,b

- Averages followed by different lowercase letters in the row differ according to Duncan's post hoc test ($P \leq 0.05$).

3.4. Intestinal morphometry

There was no difference between treatments in intestinal morphometry on day 35 of the experiment (Table 7).

Table 7 - Effects of liquid delactosed permeate (DLP) on the intestinal morphometry of nursery piglets on day 351

Item	Treatment			SEM	P-value
	CD	SD	MD		
Jejunum					
Villus height (μm)	468	509	499	18.87	0.717
Crypt depth (μm)	200	204	214	11.37	0.785
Villus height:crypt depth	2.83	2.66	2.49	0.141	0.582
Ileum					
Villus height (μm)	416	477	498	17.77	0.144
Crypt depth (μm)	211	195	205	12.16	0.845
Villus height:crypt depth	2.30	2.78	2.75	0.177	0.526

SEM - pooled standard error of the mean.

¹Data are means of seven piglets per treatment.

²Control diet (CD) supplied in dry, mash form, without DLP; supplemented diet (SD): CD + DLP provided in specific feeders and separated from the mash diet; moist diet (MD): CD moistened with DLP in a 1:1 ratio.

3.5. Bacterial population counts

There were no differences between treatments in bacterial populations or in the *Lactobacillus:E. coli* ratio in the jejunum and ileum (Table 8).

Table 8 - Effects of liquid delactosed permeate (DLP) on the count of the number of colony-forming units ($\log_{10} \text{g}^{-1}$) of *Escherichia coli*, *Lactobacillus* and their ratios in the small intestine of nursery piglets on day 35¹

Item	Treatment			SEM	P-value
	CD	SD	MD		
	Jejunum				
<i>Escherichia coli</i>	4.93	5.50	4.67	0.272	0.499
<i>Lactobacillus</i>	7.83	8.13	8.46	0.187	0.459
<i>Lactobacillus:Escherichia coli</i> ratio	1.54	1.50	1.94	0.094	0.134
	Ileum				
<i>Escherichia coli</i>	5.49	6.36	4.76	0.351	0.120
<i>Lactobacillus</i>	8.31	8.31	8.49	0.124	0.742
<i>Lactobacillus:Escherichia coli</i> ratio	1.63	1.34	1.93	0.114	0.064

SEM - pooled standard error of the mean.

¹Data are means of seven piglets per treatment.

²Control diet (CD) supplied in dry, mash form, without DLP; supplemented diet (SD): CD + DLP provided in specific feeders and separated from the mash diet; moist diet (MD): CD moistened with DLP in a 1:1 ratio.

4. DISCUSSION

In the current study, we tested DLP (195% lactose, and 2.4% crude protein) fed separately in specific feeders (in liquid form) or mixed with a diet containing lactose (11.0 and 8.0% lactose in pre-starter I and II phases, respectively) in the form of baby feed, with the hypothesis that DLP supplementation would improve growth performance by positively influencing the intestinal health of piglets weaned at 25 days old. Delactosed permeate is a coproduct with a low crude protein concentration (Oliveira et al., 2019), but it is a source of oligosaccharides, minerals (Barile et al., 2009), and lactose (although with a lower content than conventional permeate) (Oliveira et al., 2019). In the present study, positive effects were observed on feed conversion in piglets in the pre-starter phases fed DLP in both forms of supply. This is attributed to the fact that the nutrients and energy present in DLP help with growth rate, stimulation of immunity, and development of the intestinal mucosa (Jang et al., 2021a), positively affecting the absorption cells (e.g. enterocytes) that line the intestinal mucosa (Li et al., 2018).

On the other hand, piglets weaned at 23 days old and 5.8 kg BW, fed a diet containing 35.5% lactose, had lower ADG compared with those given a diet with 22.5% lactose in the first two weeks post-weaning (Mahan and Newton, 1993). This finding suggested that weaning age alters the growth response to dairy co-product intake and influences dietary lactose requirements for optimal growth performance, as also observed in a study conducted by Jang et al. (2021b), who evaluated weaning age groups (21 and 25 days old) and whey permeate contents (7.5, 11.2, 15.0, and 18.7%, that is, 5.3, 7.9, 11.1, and 13.6% of lactose, respectively) in piglet diets. According to the same authors, the daily intake of 14.4% whey permeate (10.3% of lactose, 43.1 g of lactose day⁻¹) increased the feed efficiency of piglets weaned at 25 days old, weighing from 7 kg to 11 kg BW.

The lactose present in DLP plays an important role in post-weaning piglet performance due to its sweetness, which improves the palatability of the diet (Zhao et al., 2021), although this characteristic did not alter the animals' feed intake in the present study. As it is also a carbohydrate that is easily digested by lactase, lactose is converted into glucose and galactose, providing readily available energy (Vente-Spreuwenberg et al., 2003). This additional energy content provided by the supply of DLP may reflect in improvements in the growth of nursery piglets due to a satiety effect, as also reported by Jang et al. (2021a,b). However, there is a decline in lactase activity in piglets between the third and fifth week of

birth, reflecting a lack of response in growth performance in piglets in the starter phase (seven to nine weeks) (Pluske et al., 2003), as observed in the present study. In addition, the weaning period can cause stress to the animal due to environmental changes and abrupt dietary changes, modifying the environment and intestinal morphometry (Hu et al., 2014). Diets moistened with DLP can promote improvements in nutrient and energy digestibility (although not evaluated in this study) and positively influence piglet performance responses (e.g., better feed conversion in animals fed moist diet). Liquid delactosed permeate can provide a greater supply of energy, nutrients, and water to the enterocytes, and this promotes greater development of the intestinal villi (although we did not observe such alterations), improving digestive and absorption capacity (Vasa et al., 2024).

The digestibility of dietary lactose can provide more digestible energy or a suitable microenvironment for the development of intestinal epithelial cells, favoring the ability to absorb nutrients in weaned piglets (Zhao et al., 2021) and, therefore, DLP can contribute to better nutrient absorption. Zhang et al. (2020) obtained a trend of better energy digestibility in the group of piglets fed diet moistened with water (1:2.5 ratio). Therefore, the composition of water present in DLP (60.0 to 75.0%), a nutrient that plays a crucial role in muscle growth and allows for a better transition from milk feeding (Zhang et al., 2020) associated with the type of diet supply (e.g., moistened or liquid), may be an alternative to reduce cases of dehydration in the post-weaning period that compromise the performance of nursery piglets (Russell et al., 1996).

However, by stimulating feed intake using a diet supplemented with DLP, there is also an increase in lactose content, and the increase in this carbohydrate can cause disorders in the GIT (Zhao et al., 2021). Excessive lactose fermentation results in diarrhea because it leads to intestinal barrier dysfunction (Venema, 2012), as observed in the present study in pre-starter I phase (days 0 to 7) in piglets given the MD diet. Similar results were obtained by Pierce et al. (2005), in which piglets fed a diet consisting of 29.5% lactose had a higher DO in the second and third weeks post-weaning compared with piglets fed a diet containing 17.5% lactose.

This supports the results observed in the present study because excess lactose also causes an osmotic imbalance in the intestinal epithelium (Zhao et al., 2021), signs similar to those of lactose intolerance in humans (Forsgård, 2019). This hypothesis needs to be further investigated because it has previously been shown that lactase activity decreased by 60.0 to 80.0% throughout the small intestine in nursery piglets during the first eight days post-weaning (Pié et al., 2004). This fact might explain the occurrence of post-weaning diarrhea in piglets fed higher lactose content due to lower lactase enzyme activity, causing damage to intestinal

barrier functions (e.g., inflammation) (Peace et al., 2011). The reduction in lactase activity with DLP supplementation or excessive intake of lactose may be associated with a reduction in pH in the intestine due to microbial fermentation that produces short-chain fatty acids (SCFA) from lactose and milk oligosaccharides (Montalto et al., 2006). For the results of better FCR but higher DO, we hypothesized that: first, although piglets can digest lactose, excessive lactose intake may cause undigested lactose to reach the large intestine. As a result, it is fermented by intestinal bacteria, which produce gas and organic acids. This can lead to osmotic diarrhea, as water is fluid into the intestine. Second, a sudden increase in the lactose content of DLP can cause intestinal dysbiosis. For example, some pathogenic bacteria other than *E. coli* may show increased growth, causing digestive disorders, which can manifest as diarrhea. Third, due to its high sugar and mineral content, DLP can increase the osmolarity of the intestinal content, promoting water retention in the intestine and contributing to diarrhea, without negatively affecting the animals' performance.

With regard to the blood profile, the use of dairy co-products in diets can promote changes in blood concentration, immune system (D'Alessandro et al., 2022), and GIT in newly weaned piglets (Xiao et al., 2016). Total protein (e.g., globulin, albumin) show marked changes in piglets as they age (Tóthová et al., 2021). Marder et al. (1990) reported that total protein concentrations are affected by factors such as dehydration, disease, age, and even the diet provided (e.g., crude protein content and intake). Although a change in total protein concentration was observed in the pre-starter phase (on day 21 of the experiment), these results were within the average (values of 5.5 g dL⁻¹ in piglets) reported by Ekert Kabalin et al. (2017). It has been previously shown that total protein concentrations are affected by the enhanced activity of metabolic changes (e.g., high growth rate of pigs) (ChmielowiecKorzeniowska et al., 2012) and are associated with the physiological maturation of the liver (ParísOller et al., 2022)

When the complete blood count was evaluated, there was a change only in the concentration of leukocytes, although this increase was kept within the limits considered normal for the species (between 8.7 and 37.7×10⁹) (Friendship and Henry, 1992). Hematological changes (e.g., number of leukocytes) in newborn piglets may be associated with the maturation and development of organs (e.g., large intestine) (Ekert Kabalin et al., 2008), as well as with protection of the organism against infections (Czech et al., 2017), because the immune system tries to compensate for a functional immaturity through an increased number of leukocytes (Sipos et al., 2011). Therefore, the blood profile results found were positive for the health and development of the animals in this study. However, more studies are needed to investigate the

interference of co-products such as DLP on the blood metabolites of piglets in the nursery phase.

Lactose can be fermented by intestinal bacterial populations (e.g., *Lactobacillus* and Bifidobacteria), which produce lactic and acetic acids (Kim et al., 2005). These compounds reduce intestinal pH (although in the present study, there was no effect on the small intestine) and inhibit the growth of pathogenic bacteria such as *E. coli* (Zhao et al., 2021). In the present study, the animals fed the moist diet treatment had lower pH values of the colonic contents. The fermentative processes of lactose or its monosaccharides to generate lactic acid or SCFA (Zhao et al., 2021) promote colonic acidity (Bach Knudsen, 2012). However, piglets fed DLP alone had higher pH values of the colonic content compared with those given the moist diet, which could be related to the form in which the diets were given and the fermentation process of the nutrients.

Short-chain fatty acids can also influence intestinal morphometry, because the colon easily absorbs these compounds and uses them as a source of energy for the growth of colonocytes (Comalada et al., 2006). This fact explains the greater length of the large intestine in piglets from both dietary treatments containing DLP. In a previous study, Zhou et al. (2019) evaluated the concentration of SCFA and the development of the GIT in piglets weaned at 21 days old up to 14 days post-weaning, and observed an increase in the concentration of SCFA and, consequently, a greater length of the large intestine, especially in the colonic portion.

Although no significant results were obtained in the count of intestinal bacterial populations, a previous study (Tran et al., 2012) observed a minor change in the rectal microbial composition of some *Lactobacillus* species in post-weaning piglets fed diets containing 20.0, 15.0, and 5.0% lactose in the pre-starter I and II and starter phases, respectively. In the current study, the *E. coli* count was markedly lower in the ileum of piglets fed the moist diet, influencing the *Lactobacillus*:*E. coli* ratio. The oligosaccharides present in DLP are responsible for improvements in GIT health because these compounds are complex glycans that act as prebiotics and antimicrobials responsible for immune responses in the intestine (Bode, 2012) and for inhibiting the binding of pathogens to intestinal epithelial cells (Smilowitz et al., 2014). Although we did not observe changes in intestinal morphometry in any of the dietary treatments tested, Hu et al. (2013) reported that the adverse impacts of weaning on intestinal morphometry tend to intensify during the first week post-weaning, reaching a complete recovery by the fourteenth day post-weaning. In the present study, the intestinal morphometry of piglets was evaluated at 35 days post-weaning, when the control diet did not contain lactose, except in animals fed DLP. This may explain the lack of significant results because the animals'

intestines had possibly already recovered from injuries or were not affected as expected. Intestinal villi have a quick recovery rate (Amdi et al., 2021), and it is plausible that the piglets had in fact regenerated their intestinal villi, as well as the lactose content ingested at this phase was insufficient to significantly alter the jejunal and ileal morphometry in starter piglets. Further studies in the nursery phase on the implications of this dairy co-product in the feeding of nursery piglets are crucial.

5. CONCLUSIONS

The use of liquid delactosed permeate given alone or mixed as baby feed with a conventional diet containing lactose improves feed conversion in piglets in the pre-starter phases, although a greater DO is observed in these animals in the first few days post-weaning. In addition, the use of DLP in both forms of supply promotes minor changes in other variables (blood profile and GIT traits) in the study, but without negatively affecting the growth rate of piglets in the nursery phase.

REFERENCES

- ADPI - American Dairy Products Institute. **Dairy milk permeate**. 2015. Disponível: <<https://www.adpi.org/ingredient-resource-center/dairy-milk-permeate/>> Acesso: 01 de novembro de 2023
- AMDI, C.; PEDERSEN, M. L. M.; KLAABORG, J.; MYHILL, L. J.; ENGELSMAN, M. N.; WILLIAMS, A. R. and THYMANN, T. Pre-weaning adaptation responses in piglets fed milk replacer with gradually increasing amounts of wheat. **British Journal of Nutrition**, v. 126, n. 3, p. 375-382, 2021. Doi: 10.1017/S0007114520004225
- BACH KNUDSEN, K. E. Lactose in diet influences the degradation of mixed linked β (1-3; 1-4)-D-glucan in the small intestine of pigs. **Journal of Animal Science**, v. 90, n. suppl_4, p. 125- 127, 2012. Doi: 10.2527/jas.53788
- BARILE, D.; TAO, N.; LEBRILLA, C. B.; COISSON, J.; ARLORIO, M. and GERMAN, J. B. Permeate from cheese whey ultrafiltration is a source of milk oligosaccharides. **International Dairy Journal**, v. 19, n. 9, p. 524-530, 2009. Doi: 10.1016/j.idairyj.2009.03.008
- BODE, L. Human milk oligosaccharides: every baby needs a sugar mama. **Glycobiology**, v. 22, n. 9, p. 1147-1162, 2012. Doi: 10.1093/glycob/cws074
- CHMIELOWIEC-KORZENIOWSKA, A.; TYMCZYNA, L. and BABICZ, M. Assessment of selected parameters of biochemistry, hematology, immunology and production of pigs fattened in different seasons. **Archives Animal Breeding**, v. 55, n. 5, p. 469-479, 2012.
- COMALADA, M.; BAILÓN, E.; HARO, O.; LARA-VILLOSLADA, F.; XAUS, J.; ZARZUELO, A. and GÁLVEZ, J. The effects of short-chain fatty acids on colon epithelial proliferation and survival depend on the cellular phenotype. **Journal of Cancer Research and Clinical Oncology**, v. 132, p. 487-497, 2006. Doi: 10.1007/s00432-006-0092-x
- CZECH, A.; KLEBANIUK, R.; GRELA, E. R.; SAMOLIŃSKA, W. and OGNIK, K. Polish crossbred pigs' blood haematological parameters depending on their age and physiological state. **Annals of Warsaw University of Life Sciences–SGGW Animal Science**, n. 56, p. 2, 2017. Doi: 10.22630/AAS.2017.56.2.20
- DALLAS, D. C.; WEINBORN, V.; MOURA BELL, J. M. L. N.; WANG, M.; PARKER, E. A.; GUERRERO, A.; HETTINGA, K. A.; LEBRILLA, C. B.; GERMAN, J. B. and Barile, D. Comprehensive peptidomic and glycomic evaluation reveals that sweet whey permeate from colostrum is a source of milk protein-derived peptides and oligosaccharides. **Food Research International**, v. 63, p. 203-209, 2014. Doi: 10.1016/j.foodres.2014.03.021
- D'ALESSANDRO, E.; ARFUSO, F.; FLORIDIA, V. TARDIOLO G.; FAZIO, F.; GIANNETO, C.; PICCIONE, G. and ZUMBO, A. Different genotype and a liquid whey-supplemented diet influence the resilience of pigs through immune-modulation and anti-inflammatory response. **Frontiers in Veterinary Science**, v. 9, p. 1046101, 2022. Doi: 10.3389/fvets.2022.1046101

EKERT KABALIN, A.; BALENOVIĆ, T.; VALPOTIĆ, I.; PAVIČIĆ, Ž. and VALPOTIĆ, H. The influence of birth mass and age of suckling piglets on erythrocyte parameters. **Veterinarski Arhiv**, v. 78, n. 4, p. 307-319, 2008.

EKERT KABALIN, A.; BALENOVIĆ, T.; ŠPERANDA, M.; MILINKOVIĆ-TUR, S., MENČIK, S.; MAURIĆ, M. and PAVIČIĆ, Ž. Serum biochemical parameters in suckling piglets with low and average birth mass. **Veterinarski Arhiv**, v. 87, n. 2, p. 171-184, 2017.

FONDEVILA, G.; SALDAÑA, B.; CÁMARA, L.; AGUIRRE, L. and MATEOS, G. G. Use of recycled co-products from the food industry: effects on nutrient digestibility and growth performance in pigs from 7 to 23 kg. **Animal Feed Science and Technology**, v. 276, p. 114932, 2021. Doi: 10.1016/j.anifeedsci.2021.114932

FORSGÅRD, R. A. Lactose digestion in humans: intestinal lactase appears to be constitutive whereas the colonic microbiome is adaptable. **The American journal of Clinical Nutrition**, v. 110, n. 2, p. 273-279, 2019. Doi: 10.1093/ajcn/nqz104

FRIENDSHIP, R. M. and HENRY, S. C. 1992. **Cardiovascular system, haematology and clinical chemistry**. p.3-11. In: Diseases of swine. 7th ed. Leman, A. D.; Straw, B. E.; Mengeling, W. L.; D'Allaire, S. and Taylor, D. J., eds. Iowa State University Press, Ames.

GENOVA, J. L.; RUPOLO, P. E.; MELO, A. D. B.; SANTOS, L. B. A.; WENDT, G. N.; BARBOSA, K. A.; CARVALHO, S. T.; OLIVEIRA, N. T. E.; COSTA, L. B. and CARVALHO, P. L. O. 2021. Biological response of piglets challenged with *Escherichia coli* F4 (K88) when fed diets containing intestinal alkaline phosphatase. **Czech Journal of Animal Science**, v. 66, p. 391-402. Doi: 10.17221/82/2021-CJAS

GENOVA, J. L.; MELO, A. D. B.; RUPOLO, P. E.; MACEDO, R. E. F.; ENGRACIA FILHO, J. R.; CARVALHO, S. T.; FAUCITANO, L.; COSTA, L. B. and CARVALHO, P. L.O. 2022. New findings of intestinal alkaline phosphatase: effects on intestinal and organ health of piglets challenged with ETEC F4 (K88). **Revista Brasileira de Zootecnia**, v. 51, p. 20210144. Doi: 10.37496/rbz5120210144

HU, C. H.; XIAO, K.; LUAN, Z. S.; SONG, J. Early weaning increases intestinal permeability, alters expression of cytokine and tight junction proteins, and activates mitogen-activated protein kinases in pigs. **Journal of animal science**, v. 91, n. 3, p. 1094-1101, 2013. Doi: 10.2527/jas.2012-5796

HU, Y.; DUN, Y.; LI, S.; ZHAO, S.; PENG, N.; LIANG, Y. Effects of *Bacillus subtilis* KN-42 on growth performance, diarrhea and faecal bacterial flora of weaned piglets. **Asian-Australasian Journal of Animal Sciences**, v. 27, n. 8, p. 1131, 2014. Doi: 10.5713/ajas.2013.13737

ISO - International Organization for Standardization. 2006. **ISO 4832: Microbiology of food and animal feeding stuffs—horizontal method for the enumeration of coliforms—colony-count technique**. International Organization for Standardization, Geneva. Available at: <<https://www.iso.org/standard/38282.html>>. Accessed on: Jan. 01, 2023

JANG, K. B.; PURVIS, J. M. and KIM, S. W. Dose-response and functional role of whey permeate as a source of lactose and milk oligosaccharides on intestinal health and growth of nursery pigs. **Journal of Animal Sciences**, v. 99, n. 1, p. skab008, 2021a. Doi:10.1093/jas/skab008

JANG, K. B., DUARTE, M. E.; PURVIS, J. M. and KIM, S. W. Impacts of weaning age on dietary needs of whey permeate for pigs at 7 to 11 kg body weight. **Journal of Animal Science and Biotechnology**, v. 12, p. 1-8, 2021b. Doi: 10.1186/s40104-021-00637-4.

KIM, Y. Y.; KIL, D. Y.; OH, H. K and HAN, In K. Acidifier as an alternative material to antibiotics in animal feed. **Asian-Australasian Journal of Animal Sciences**, v. 18, n. 7, p. 1048- 1060, 2005. Doi:10.5713/ajas.2005.1048

LEPINE, A. J.; MAHAN, D. C.; CHUNG, Y. K. Growth performance of weanling pigs fed corn-soybean meal diets with or without dried whey at various L-lysine. HCl levels. **Journal of Animal Science**, v. 69, n. 5, p. 2026-2032, 1991. Doi:10.2527/1991.6952026x

LI, Y.; NGUYEN, D. N.; OBELITZ-RYOM, K.; ANDERSEN, A. D.; THYMANN, T.; CHATTERTON, D. E. and SANGILD, P. T. Bioactive whey protein concentrate and lactose stimulate gut function in formula-fed preterm pigs. **Journal of Pediatric Gastroenterology and Nutrition**, v. 66, n. 1, p. 128-134, 2018. Doi 10.1097/MPG.0000000000001699

MAHAN, D. C. and NEWTON, E. A. Evaluation of feed grains with dried skim milk and added carbohydrate sources on weanling pig performance. **Journal of Animal Science**, v. 71, n. 12, p. 3376-3382, 1993. Doi:10.2527/1993.71123376x

MARDER, J.; EYLATH, U.; MOSKOVITZ, E. and SHARIR, R. The effect of heat exposure on blood chemistry of the hyperthermic rabbit. *Comparative Biochemistry and Physiology. A, Comparative Physiology*, v. 97, n. 2, p. 245-247, 1990. Doi: 10.1016/0300- 9629(90)90179-v

MONTALTO, M.; CURIGLIANO, V.; SANTORO, L.; VASTOLA, M.; CAMMAROTA, G.; MANNA, R.; GASBARRINI, A. and GASBARRINI, G. Management and treatment of lactose malabsorption. **World Journal of Gastroenterology**, v. 12, p 187-191, 2006. Doi: 10.3748/wjg.v12.i2.187

OLIVEIRA, D.; PURI, R.; FENELON, M. A. and O'MAHONY, J. A. Delactosed permeate as a dairy processing co-product with major potential value: a review. **International Journal of Food Science & Technology**, v. 54, n. 4, p. 999-1008, 2019. Doi: 10.1111/ijfs.14064

PARÍS-OLLER, E.; MATÁS, C.; ROMAR, R.; LOPES, J. S.; GADEA, J.; CÁNOVAS, S. and COY, P. Growth analysis and blood profile in piglets born by embryo transfer. **Research in Veterinary Science**, v. 142, p. 43-53, 2022. Doi: 10.1016/j.rvsc.2021.11.010

PEACE, R. M.; CAMPBELL, J.; POLO, J.; CRENSHAW, J.; RUSSELL, L. and MOESER, A. Spray-dried porcine plasma influences intestinal barrier function, inflammation, and diarrhea in weaned pigs. **The Journal of Nutrition**, v. 141, n. 7, p. 1312-1317, 2011. Doi: 10.3945/jn.110.136796.

PÉREZ-CALVO, E.; WICAKSONO, A. N.; CANET E.; DAULTON E. ENS W.; HOELLER, U.; VERLHAC, V.; CELI, P. and COVINGTON, J. A. The measurement of volatile organic compounds in faeces of piglets as a tool to assess gastrointestinal functionality. **Biosystems Engineering**, v. 184, p. 122-129, 2019.

PIÉ, S.; LALLÈS, J. P.; BLAZY, F.; LAFFITTE, J.; SÈVE, B. and OSWALD, I. P. Weaning is associated with an upregulation of expression of inflammatory cytokines in the intestine of piglets. **The Journal of nutrition**, v. 134, n. 3, p. 641-647, 2004. Doi: 10.1093/jn/134.3.641.

PIERCE, K. M.; SWEENEY, T.; BROPHY, P. O.; CALLAN, J. J.; MCCARTHY, P. and O'DOHERTY, J. V. Dietary manipulation post weaning to improve piglet performance and gastro-intestinal health. **Animal Science**, v. 81, n. 3, p. 347-356, 2005. Doi: 10.1079/ASC50400347

PIERCE, K. M.; SWEENEY, T.; BROPHY, P. O.; CALLAN, J. J.; FITZPATRICK, E.; MCCARTHY, P. and O'DOHERTY, J. V. The effect of lactose and inulin on intestinal morphology, selected microbial populations and volatile fatty acid concentrations in the gastro-intestinal tract of the weanling pig. **Animal Science**, v. 82, n. 3, p. 311-318, 2006. Doi: 10.1079/ASC200634

PLUSKE, J. R.; KERTON, D. K.; CRANWELL, P. D.; CAMPBELL R. G.; MULLAN, P. B.; KING, R. H.; POWER, G. N.; PIERZYNOWSKI, S. G.; WESTROM, B.; RIPPE, C.; PEULEN, O. and DUNSHEA, F. R. Age, sex, and weight at weaning influence organ weight and gastrointestinal development of weanling pigs. **Australian Journal of Agricultural Research**, v. 54, n. 5, p. 515-527, 2003. Doi: 10.1071/AR02156

ROSTAGNO, H. S.; ALBINO, L. F. T.; HANNAS, M. I.; DONZELE, J. L.; SAKOMURA, N. K.; PERAZZO, F. G.; SARAIVA, A.; ABREU, M. L. T.; RODRIGUES, P. B.; OLIVEIRA, R.F.; BARRETO, S. L. T. and BRITO, C. O. 2017. **Tabelas brasileiras para aves e suínos: composição de alimentos e exigências nutricionais**. 4.ed. Universidade Federal Viçosa, Departamento de Zootecnia, Viçosa, MG.

RUSSELL, P. J.; GEARY, T. M.; BROOKS, P. H. and CAMPBELL, A. Performance, water use and effluent output of weaner pigs fed ad libitum with either dry pellets or liquid feed and the role of microbial activity in the liquid feed. **Journal of the Science of Food and Agriculture**, v. 72, n. 1, p. 8-16, 1996. Doi: 10.1002/(SICI)1097-0010(199609)72:1<8::AID-JSFA646>3.0.CO;2-K

SIPOS, W.; DUVIGNEAU, C. J.; HARTL, R. T. and SCHWENDENWEIN, I. Exploratory reference intervals on hematology and cellular immune system of multiparous Large White sows. **Veterinary Immunology and Immunopathology**, v. 141, n. 3-4, p. 307-311, 2011. Doi: 10.1016/j.vetimm.2011.03.007

SMILOWITZ, J. T.; LEBRILLA, C. B.; MILLS, D. A. J.; GERMAN, B. and FREEMAN, S. L. Breast milk oligosaccharides: structure-function relationships in the neonate. **Annual Review of Nutrition**, v. 34, p. 143-169, 2014. Doi: 10.1146/annurev-nutr-071813-105721

SUTERA, A. M.; ARFUSO, F.; TARDIOLO, G.; RIGGIO, V.; FAZIO, F.; CIGLIANO, R. A.; PAYTUVI, A.; PICCIONE, G. and ZUMBO, A. Effect of a co-feed liquid whey-integrated diet on crossbred pigs' fecal microbiota. **Animals**, v. 13, n. 11, p. 1750, 2023. Doi: 10.3390/ani13111750

TÓTHOVÁ, C.; LINK, R.; KYZEKOVÁ, P. and NAGY, O. Serum protein electrophoretic pattern in piglets during the early postnatal period. **Scientific Reports**, v. 11, n. 1, p. 17539, 2021. Doi: 10.1038/s41598-021-96957-6

TRAN, H.; MORENO, E. E.; HINKLE, J. W.; BUNDY, J.; WALTER, T. E.; BURKEY, P. S. Effects of lactose and yeast-dried milk on growth performance, fecal microbiota, and immune parameters of nursery pigs. **Journal of Animal Science**, v. 90, n. 9, p. 3049-3059, 2012. Doi: 10.2527/jas.2011-4544

VASA, S. R.; GARDINER, G. E.; ARNAUD, E. A.; O'DRISCOLL, K.; BEE, G. and LAWLOR, P. G.. Effect of supplemental milk replacer and liquid starter diet for 4 and 11 days postweaning on intestinal parameters of weaned piglets and growth to slaughter. **Animals**, v 18, p 101271, 2024. Doi: 10.1016/j.animal.2024.101271

VENEMA, K. Intestinal fermentation of lactose and prebiotic lactose derivatives, including human milk oligosaccharides. **International Dairy Journal**, v. 22, n. 2, p. 123-140, 2012. Doi:10.1016/j.idairyj.2011.10.011

VENTE-SPREEUWENBERG, M A. M.; VERDONK, J. M. A. J.; VERSTEGEN, M. W. A. and BEYNEN, A. C. Villus height and gut development in weaned piglets receiving diets containing either glucose, lactose or starch. **British Journal of Nutrition**, v. 90, n. 5, p. 907-913, 2003. Doi: 10.1079/BJN2003981

XIAO, K.; JIAO, L.; CAO, S.; SONG, Z.; HU, C.; HAN, X. Whey protein concentrate enhances intestinal integrity and influences transforming growth factor- β 1 and mitogen-activated protein kinase signaling pathways in piglets after lipopolysaccharide challenge. **British Journal of Nutrition**, v. 115, n. 6, p. 984-993, 2016. Doi:10.1017/S0007114515005085

YOO, S. H.; HONG, J. S.; YOO, H. B.; HAN, T. H.; JEONG, J. H. and KIM Y. Y. Influence of various levels of milk by-products in weaner diets on growth performance, blood urea nitrogen, diarrhea incidence, and pork quality of weaning to finishing pigs. **Asian-Australasian Journal of Animal Sciences**, v 31, p 696-704, 2018.

YU, Z.; CHEN, C. and NEWBURG, D. S. Utilization of major fucosylated and sialylated human milk oligosaccharides by isolated human gut microbes. **Glycobiology**, v. 23, n. 11, p. 1281- 1292, 2013. Doi:10.1093/glycob/cwt065

ZHANG, S.; YOO, D. H.; AO, X. and KIM, I. H. Effects of dietary probiotic, liquid feed and nutritional concentration on the growth performance, nutrient digestibility and fecal score of weaning piglets. **Asian-Australasian Journal of Animal Sciences**, v. 33, n. 10, p. 1617, 2020. Doi:10.5713/ajas.19.0473

ZHAO, J.; ZHANG, Z.; ZHANG, S.; PAGE, G. and JAWORSK, N. W. The role of lactose in weanling pig nutrition: a literature and meta-analysis review. **Journal of Animal Science and Biotechnology**, v. 12, n. 1, p. 10, 2021. Doi: 10.1186/s40104-020-00522-6

ZHOU, Z.; ZHANG, J. X.; MO, S.; TAN, X.; WANG, L.; LI, J.; LI, Y.; DING, X.; LIU, X. MA, X.; YANG, H. and YIN, Y. The production of short chain fatty acid and colonic development in weaning piglets. **Journal of Animal Physiology and Animal Nutrition**, v. 103, n. 5, p. 1530-1537, 2019. Doi: 10.1111/jpn.13164