

CAMILA GONÇALVES TEIXEIRA

**ANTIMICROBIAL PROPERTIES OF *Weissella cibaria* STRAINS ISOLATED
FROM CAMPOS DAS VERTENTES, MINAS GERAIS, BRAZIL, AND ITS
POTENTIAL USE AS PROBIOTICS AND BACTERIOCINS PRODUCERS**

Thesis submitted to the Food Science and
Technology Graduate Program of the
Universidade Federal de Viçosa in partial
fulfillment of the requirements for the degree
of *Doctor Scientiae*.

Adviser: Antônio Fernandes de Carvalho

Co-advisers: Evandro Martins
Rosângela de Freitas

**VIÇOSA - MINAS GERAIS
2022**

**Ficha catalográfica elaborada pela Biblioteca Central da Universidade
Federal de Viçosa - Campus Viçosa**

T

T266a
2022
Teixeira, Camila Gonçalves, 1992-
Antimicrobial properties of *Weissella cibaria* strains
isolated from Campos das Vertentes, Minas Gerais, Brazil, and
its potential use as probiotics and bacteriocins producers /
Camila Gonçalves Teixeira. – Viçosa, MG, 2022.
1 tese eletrônica (150 f.): il. (algumas color.).

Texto em inglês.

Inclui anexos.

Orientador: Antônio Fernandes de Carvalho.

Tese (doutorado) - Universidade Federal de Viçosa,
Departamento de Tecnologia de Alimentos, 2022.

Inclui bibliografia.

DOI: <https://doi.org/10.47328/ufvbbt.2022.789>

Modo de acesso: World Wide Web.

1. Queijo - Microbiologia. 2. Queijo-de-Minas - Campo das
Vertentes (MG : Mesoregião). 3. Bactéria do ácido láctico.
4. Tecnologia de alimentos. 5. Probióticos. I. Carvalho, Antônio
Fernandes de, 1964-. II. Universidade Federal de Viçosa.
Departamento de Tecnologia de Alimentos. Programa de
Pós-Graduação em Ciência e Tecnologia de Alimentos.
III. Título.

CDD 22. ed. 637.35


CAMILA GONÇALVES TEIXEIRA

ANTIMICROBIAL PROPERTIES OF *Weissella cibaria* STRAINS ISOLATED FROM CAMPOS DAS VERTENTES, MINAS GERAIS, BRAZIL, AND ITS POTENTIAL USE AS PROBIOTICS AND BACTERIOCINS PRODUCERS


Thesis submitted to the Food Science and Technology Graduate Program of the Universidade Federal de Viçosa in partial fulfillment of the requirements for the degree of *Doctor Scientiae*.

APPROVED: December 22, 2022.

Assent:

Documento assinado digitalmente
 CAMILA GONCALVES TEIXEIRA
Data: 23/12/2022 11:21:25-0300
Verifique em <https://verificador.iti.br>

Camila Gonçalves Teixeira
Author

Documento assinado digitalmente
 ANTONIO FERNANDES DE CARVALHO
Data: 23/12/2022 10:59:40-0300
Verifique em <https://verificador.iti.br>

Antônio Fernandes de Carvalho
Adviser

ACKNOWLEDGMENTS

To God, for walking with me and for carrying me on during the most difficult moments of my walk in my work.

To my family, especially my mothers, Francisca and Aparecida, and my fathers, Gerônimo and Genilson, for the examples of wisdom and the incentives that have always motivated me.

To my brothers, Guilherme and Henrique, and sisters Lívia and Lucimar for the moments of distraction, love, and affection.

To my boyfriend and companion Ian, for the affection, for the patience, and for always motivating me during this journey, helping me to overcome each obstacle.

To the interns at Inovaleite, who helped me a lot with the heavy work.

To all my friends at Inovaleite, especially Frida, Luana, Sidney, Raiane, and Andressa who shared and helped in every experiment and always cheered for me.

To all the friends of the Djamel laboratory group, for the company, collaboration, and moments of fun I would like to thank specially Yanath, Rabia, Désiré, Adrian, and Ali.

To friends Luiza and Gustavo, for the cheers and the words of encouragement. For not letting me down and for always being willing to listen to me.

To my friends Lauréline, Catarina, Sakhr, Luca, and Alessandra, for making my stay in France so much easier.

To Dr. Evandro, for becoming my friend during these four years and for all the uncountable time spend helping me.

To my advisors, Professor Antônio Fernandes and Djamel Drider, for the opportunity to work on their teams and be oriented by them, for the concern, friendship, and shared wisdom.

To the professors Anca Lucau-Danila and Luis Augusto Nero for being my co-advisers, and for all the help in the accomplishment of the analyses and writing the papers.

To the Federal University of Viçosa, for the opportunity to complete the postgraduate course.

This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES) – Finance Code 001.

To the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), for granting the scholarship.

To the Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG), for granting the scholarship.

To the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), to granting the scholarship.

ABSTRACT

TEIXEIRA, Camila Gonçalves, D.Sc., Universidade Federal de Viçosa, December, 2022. **Antimicrobial Properties of *Weissella cibaria* Strains Isolated from Campos Das Vertentes, Minas Gerais, Brazil and its Potential use as Probiotics and Bacteriocins Producers**. Adviser: Antônio Fernandes de Carvalho. Co-advisers: Evandro Martins and Rosângela de Freitas.

The genus *Weissella* is composed of bacteria belonging to the group known as lactic acid bacteria (LAB), are autochthonous to both dairy and non-dairy environments and many strains can produce various compounds of interest to the food industry. This work began with deep research regarding the use of *Weissella* strains for the food and pharmaceutical industry and the probiotic potential of this genus. A large number of scientific studies reported and support the use of *Weissella* strains in this field due to their health-promoting properties and technological features for the food industry. This work continues with the study of the biodiversity of *Weissella* strains isolated in the regions producing artisanal cheeses in Brazil. For this, the analysis of PFGE and Rep-PCR was carried out and it was observed that each study region represents a specific ecological niche. After, the antimicrobial potential of some representative strains was analyzed against the most common foodborne pathogens. Most of the isolates tested showed good inhibition of the pathogens. Also, the technological potential of thirteen strains was evaluated. Some strains showed positive results for most of the tests performed. After, three strains with good inhibitory activity were selected. The genomes were sequenced and assembled. With the first strain, *Weissella cibaria* W25 has carried out a comprehensive and comparative genomic analysis with *W. cibaria* 110, known to produce the weissellicin 110 bacteriocin, and four other non-bacteriocin-producing *W. cibaria* strains. The analyses showed that the strain W25 has its unique protein cluster genes which can be related to bacteriocins genes indicated by AntiSmash, confirming the possibility of producing two different bacteriocins. Based on these results the genome of the three selected strains was deeply analyzed. The presence of genes for secondary metabolites was investigated and they have a common region in the genome with a putative bacteriocin identified as Bacteriocin_IIc. This putative bacteriocin has no similarity with other bacteriocins from the genus *Weissella* so far described. Finally, the safety of using these three strains for human

and/or animal consumption and their capacity to be used as probiotics were analyzed. All three strains have no acquired antimicrobial resistance genes and no mobile elements in the genome. *W. cibaria* strains were resistant to vancomycin which is intrinsic to this genus, had no hemolytic activity, and no one of the known virulence factors was detected. They possess the defense system CRISPR-Cas. Besides they didn't present any inflammation or cytotoxicity capacity. *W. cibaria* W21 demonstrates excellent results of adhesion to intestinal cells and exclusion of *S. aureus* MRSA SA1. However, none of the three strains survived completely through the simulated gastrointestinal tract, but some protective ways can be further studied. In conclusion, this work allowed us to understand the diversity and distribution of *Weissella* species in different regions producing artisanal cheeses in Brazil. It also showed that some of them have a favorable technological and safety potential for future uses in the food industry or as a probiotic culture.

Keywords: *Weissella*. Technological properties. Antimicrobial activity. Bacteriocins.

RESUMO

TEIXEIRA, Camila Gonçalves, D.Sc., Universidade Federal de Viçosa, dezembro de 2022. **Propriedades Antimicrobianas de Estirpes de *Weissella cibaria* Isoladas dos Campos das Vertentes, Minas Gerais, Brasil e seu Potencial uso como Probióticos e Produtoras de Bacteriocinas.** Orientador: Antônio Fernandes de Carvalho. Coorientadores: Evandro Martins e Rosângela de Freitas.

O gênero *Weissella* é composto por bactérias pertencentes ao grupo conhecido como bactérias lácticas, são autóctones tanto de ambientes lácteos quanto não lácteos e muitas cepas podem produzir vários compostos de interesse para a indústria alimentícia. Este trabalho começou com uma pesquisa profunda sobre o uso de cepas de *Weissella* para a indústria alimentícia e farmacêutica e o potencial probiótico deste gênero. Um grande número de estudos científicos relatou e apoia o uso de cepas de *Weissella* neste campo devido às suas propriedades de melhoria da saúde e potencial tecnológico para a indústria de alimentos. Este trabalho continua com o estudo da biodiversidade de cepas de *Weissella* isoladas de regiões produtoras de queijos artesanais no Brasil. Para isso, foram realizadas análises de PFGE e Rep-PCR e observou-se que cada região de estudo representa um nicho ecológico específico. Depois, o potencial antimicrobiano de algumas cepas representativas foi analisado contra os patógenos de origem alimentar mais comuns. A maioria dos isolados testados apresentou boa inibição dos patógenos. Também foi avaliado o potencial tecnológico de treze cepas. Algumas cepas apresentaram resultados positivos para a maioria dos testes realizados. Após, foram selecionadas três cepas com atividade antimicrobiana e seus genomas foram sequenciados e montados. Com a primeira cepa, *Weissella cibaria* W25 realizou-se uma análise genômica abrangente e comparativa com *W. cibaria* 110, conhecida por produzir a bacteriocina weissellicina 110, e quatro outras cepas de *W. cibaria* não produtoras de bacteriocina. As análises mostraram que a cepa W25 possui seus próprios genes de proteínas que podem ser relacionados às bacteriocinas identificadas pelo programa AntiSmash, confirmando a possibilidade de produzir duas bacteriocinas diferentes. Com base nesses resultados, os genomas das três estirpes selecionadas foram profundamente analisados. A presença de genes para metabólitos secundários foi investigada e eles têm uma região comum no genoma com uma bacteriocina putativa identificada como

Bacteriocin_IIc. Esta bacteriocina putativa não tem semelhança com outras bacteriocinas do gênero *Weissella* até agora descritas. Por fim, analisou-se a segurança do uso das três estirpes para consumo humano e sua capacidade de serem usadas como probióticos. Todas as três cepas não possuem genes de resistência antimicrobiana adquiridos e nenhum elemento móvel no genoma. As estirpes selecionadas são resistentes à vancomicina, característica intrínseca a esse gênero, não apresentavam atividade hemolítica e nenhum dos fatores de virulência conhecidos foi detectado. Possuem o sistema de defesa CRISPR-Cas. Além disso, não apresentaram nenhuma capacidade inflamatória ou citotóxica. *W. cibaria* W21 demonstrou excelentes resultados de adesão às células intestinais e de exclusão de *S. aureus* MRSA SA1. No entanto, nenhuma das três estirpes sobreviveu completamente à simulação da passagem pelo trato gastrointestinal. Em conclusão, este trabalho permitiu compreender a diversidade e distribuição das espécies de *Weissella* em diferentes regiões produtoras de queijos artesanais no Brasil. Também mostrou que alguns estirpes têm potencial tecnológico e de segurança favorável para usos futuros na indústria de alimentos ou como cultura probiótica.

Palavras-chave: *Weissella*. Propriedades tecnológicas. Atividade antimicrobiana. Bacteriocinas.

SCIENTIFIC VALORIZATION

Published papers

Teixeira, Camila Gonçalves, Andressa Fusieger, Evandro Martins, et al. 2021. “Biodiversity and Technological Features of *Weissella* Isolates Obtained from Brazilian Artisanal Cheese-Producing Regions.” *LWT - Food Science and Technology* 147(April): 111474.

Teixeira, Camila Gonçalves, Raiane Rodrigues da Silva, et al. 2021. “The *Weissella* Genus in the Food Industry: A Review.” *Research, Society and Development* 10(5): 1–15.

Teixeira, Camila Gonçalves, Andressa Fusieger, Gustavo Leite Milião, et al. 2021. “*Weissella*: An Emerging Bacterium with Promising Health Benefits.” *Probiotics and Antimicrobial Proteins* 13(4): 915–25.

Teixeira, Camila Gonçalves et al. 2022. “Genomic Analyses of *Weissella Cibaria* W25, a Potential Bacteriocin-Producing Strain Isolated from Pasture in Campos Das Vertentes, Minas Gerais, Brazil.” *Microorganisms* 10(2): 314.

Conference abstracts

Camila Gonçalves Teixeira, Evandro Martins, Andressa Fusieger, Rosângela De Freitas, Tatiana Santos Lima, Júlia Oliveira Lopes, Luis Gustavo Lima Nascimento, Antônio Fernandes De Carvalho. “Technological Potential of *Weissella* Strains Isolated from Artisanal Cheeses from Different Regions of Brazil” XXI Encontro Nacional e VII Congresso Latino Americano de Analistas de Alimentos, May, 26 to 30, 2019, Florianópolis/SC (Brasil).

Camila Gonçalves Teixeira, Antônio Fernandes de Carvalho, Djamel Drider. “Brazilian *Weissella* strains with antimicrobial activity against Gram-positive and Gram-negative foodborne pathogens.” Congrès de la Société Française de Microbiologie, September, 22 to 24, 2021, Nantes (France).

Papers to be published

Camila Gonçalves Teixeira, Rafaela da Silva Rodrigues, Ricardo Seiti Yamatogi, Anca Lucau-Danila, Djamel Drider, Luís Augusto Nero, Antônio Fernandes de Carvalho. “Genome characterization of *Weissella* strains isolated from Campos das Vertentes, Minas Gerais, Brazil focused revealed new bacteriocins with a large spectrum of activity”

Camila Gonçalves Teixeira, Yanath Belguesmia, Rafaela da Silva Rodrigues, Anca Lucau-Danila, Luís Augusto Nero, Antônio Fernandes de Carvalho, Djamel Drider. “Biosecurity of *Weissella* strains isolated from Campos das Vertentes, Minas Gerais, Brazil focused on its potential as probiotic”.

SUMMARY

GENERAL INTRODUCTION	14
1. Context	15
2. Objectives.....	15
3. Manuscript organization	15
CHAPTER I: Literature review.....	15
CHAPTER II: Biodiversity and technological features of <i>Weissella</i> isolates obtained from Brazilian artisanal cheese-producing regions.....	16
CHAPTER III: <i>Weissella cibaria</i> W25: Comparative genomic analyses	16
CHAPTER IV: <i>Weissella cibaria</i> strains isolated from Campo das Vertentes (MG), a potential bacteriocin producers	16
CHAPTER V: <i>Weissella cibaria</i> strains isolated from Campo das Vertentes (MG), biosecurity and probiotics analyses.....	17
4. General conclusions and perspectives	17
CHAPTER I: Literature Review	18
PART A: The <i>Weissella</i> genus in the food industry: a review.....	19
1. Introduction	22
2. Methodology.....	24
3. Physiological and metabolic characteristics	25
4. Ecology	27
5. Some applications of bacteria from genus <i>Weissella</i> in food	29
6. Conclusion	35
7. References.....	35
PART B: <i>Weissella</i>: An Emerging Bacterium with Promising Health Benefits	43
1. Introduction	44
2. <i>Weissella</i> Species Identification and Characterization.....	45
3. Beneficial Properties of <i>Weissella</i>	47
4. Health Benefits of <i>Weissella</i>	47
5. EPS Production by <i>Weissella</i>	51

6. <i>Weissella</i> as an Emerging Pathogen	54
7. Conclusion	56
8. References.....	56
CHAPTER II: Biodiversity and technological features of <i>Weissella</i> isolates obtained from Brazilian artisanal cheese-producing regions	65
Biodiversity and technological features of <i>Weissella</i> isolates obtained from Brazilian artisanal cheese-producing regions.....	66
1. Introduction	67
2. Material and methods.....	68
2. Results	74
3. Discussion.....	80
4. Conclusion	83
5. Appendix A. Supplementary data.....	83
6. References.....	84
CHAPTER III: <i>Weissella cibaria</i> W25: Comparative genomic analyses	91
Genomic analyses of <i>Weissella cibaria</i> W25, a potential bacteriocin-producing strain isolated from pasture in Campos das Vertentes, Minas Gerais, Brazil	92
1. Introduction	93
2. Materials and Methods.....	93
3. Results and Discussion.....	95
4. Conclusions.....	101
5. References.....	102
CHAPTER IV: <i>Weissella cibaria</i> strains isolated from Campo das Vertentes (MG), a potential bacteriocin producers	105
Genome characterization of <i>Weissella</i> strains isolated from Campos das Vertentes, Minas Gerais, Brazil focused revealed new bacteriocins with a large spectrum of activity	106
1. Introduction	107
2. Material and Methods.....	108
3. Results and Discussion.....	111
4. Conclusions.....	116

5. Data Availability.....	117
6. Conflict of Interest	117
7. Acknowledgements.....	117
8. References.....	117

CHAPTER V: *Weissella cibaria* strains isolated from Campo das Vertentes

(MG), biosecurity and probiotics analyses	123
--	------------

Biosecurity of *Weissella* strains isolated from Campos das Vertentes, Minas

Gerais, Brazil focused on its potential as probiotic.....	124
--	------------

1. Introduction	125
2. Material and Methods.....	126
3. Results and Discussion.....	131
4. Conclusion	139
5. References.....	140

GENERAL CONCLUSION.....	145
--------------------------------	------------

GENERAL INTRODUCTION

1. Context

The study of microbial diversity in dairy and non-dairy environments has great importance for the understanding of the presence of different microorganisms in these environments, as well as the impact of these microorganisms on the final product when we refer to artisanal dairy products. Each environment has unique characteristics that favor and allow the development of different bacterial species.

Handmade cheeses and raw milk are considered as potential sources of new strains of lactic acid bacteria. The way of making these cheeses causes the fermentation to be conducted by bacteria contaminating the grazing, animal skin, utensils, surfaces and other environments that may come into contact with the cheese during manufacture. The study of the bacterial community present in artisanal cheeses has revealed the presence of species that had not yet been related to cheeses and a great diversity of lactic bacteria with differentiated technological characteristics. In addition, non-dairy environments such as grass, different types of silage and even animal skin have also been an important source of new strains that have adapted and therefore, can provide interesting features to be explored.

The diversity of *Weissella* isolated from dairy and non-dairy environments is of great interest for the understanding of this microorganism in the final products and can be studied by molecular techniques such as PFGE and Rep-PCR that are able to phylogenetically relate the strains. In addition, phenotypic characterization is necessary to understand the characteristics and potential of each of them.

2. Objectives

This study aims to understand the microbial diversity in different environments (dairy and non-dairy) seeking the application of the strains with potential to be used as a bacteriocin producer for the food industry or for the pharmaceutical industry as a probiotic.

3. Manuscript organization

CHAPTER I: Literature review

This chapter is divided into two parts, the first part presents overall information about the *Weissella* genus and its potential application for the food and pharmaceutical industry. The physiological and metabolic characteristics, ecology and application in

the food industry is the part A. In the second part (part B), the *Weissella* strains are approached for their species identification and characterization, beneficial properties, health benefits and exopolysaccharides production, and possibility for being used as probiotic culture. The first part of this review is published as “*The Weissella Genus in the Food Industry: A Review*” in the *Research, Society and Development*. The second part of this review is published as “*Weissella: An Emerging Bacterium with Promising Health Benefits*” in the *Probiotics and Antimicrobial Proteins*.

CHAPTER II: Biodiversity and technological features of *Weissella* isolates obtained from Brazilian artisanal cheese-producing regions

This part aimed to identify the microbial biodiversity and evaluate the technological potential of *Weissella* spp. isolates obtained from different Brazilian regions that produce artisanal cheeses, such as Campo das Vertentes (Minas Gerais), Southern Pará, Guanambi (Bahia) and Marajó Island. The contents of this chapter are published as “Biodiversity and Technological Features of *Weissella* Isolates Obtained from Brazilian Artisanal Cheese-Producing Regions” in *LWT - Food Science and Technology*.

CHAPTER III: *Weissella cibaria* W25: Comparative genomic analyses

This part of the thesis aimed to announce and analyze the sequencing and annotation of the *Weissella cibaria* W25 genome and carry out a comprehensive comparative genomic analysis with *W. cibaria* 110, known to produce the weissellicin 110 bacteriocin, and four other non-bacteriocin-producing *W. cibaria* strains. The contents of this chapter are published as “Genomic Analyses of *Weissella Cibaria* W25, a Potential Bacteriocin-Producing Strain Isolated from Pasture in Campos Das Vertentes, Minas Gerais, Brazil.” In *Microorganisms*.

CHAPTER IV: *Weissella cibaria* strains isolated from Campo das Vertentes (MG), a potential bacteriocin producers

The aim of this part is to identify the antimicrobial compounds produced by *Weissella* strains isolated from Campos das Vertentes, Minas Gerais, Brazil, scheduling their future application as pathogens inhibitors in the food industry and preservation. The content of this chapter will be submitted as “*Genome characterization of Weissella*

strains isolated from Campos das Vertentes, Minas Gerais, Brazil focused revealed new bacteriocins with a large spectrum of activity”.

CHAPTER V: *Weissella cibaria* strains isolated from Campo das Vertentes (MG), biosecurity and probiotics analyses

In this last chapter we evaluated the safety capacity of three strains of *Weissella cibaria*, isolated from different producing artisanal cheese regions of Campos das Vertentes, in Minas Gerais, Brazil to be use as probiotic for human consumption or animal feed. This chapter will be valorized as “Biosecurity of *Weissella* strains isolated from Campos das Vertentes, Minas Gerais, Brazil focused on its potential as probiotic”

4. General conclusions and perspectives

The genus *Weissella* has many interesting properties and can be explored to aggregate more to the food and pharmaceutical industries. The main conclusions derived from the study are lined out with the future research needed to increment the knowledge about the strains regarding their use as probiotics or as an ingredient in the food industry.

CHAPTER I: Literature Review

PART A: The *Weissella* genus in the food industry: a review
O gênero *Weissella* na indústria de alimentos: uma revisão
El género *Weissella* en la industria alimentaria: una revisión

The content of this part has been published in:

Research Society and Development

Camila Gonçalves Teixeira, Raiane Rodrigues da Silva, Andressa Fusieger,
Evandro Martins, Rosângela de Freitas, Antônio Fernandes de Carvalho

License to reuse this content in thesis is in Annex 1

Abstract

The genus *Weissella* is composed of bacteria classified as Gram-positive, catalase negative, non-spore forming, coccoid morphology or short bacilli. They belong to the group of lactic acid bacteria (LAB), mainly by production of lactic acid from the fermentation of carbohydrates. *Weissella* species are distributed in different habitats, such as soils, milking machines, sugar cane and some strains with interesting technological features can be isolated from fermented foods, such as cheeses made from raw milk, fermented vegetables and fermented milk. From the point of view of food technology, some strains have potential in the production of exopolysaccharides, non-digestible oligosaccharides, which is beyond their probiotic potential. Therefore, the bacteria belonging to the genus *Weissella* might have great technological importance, being also involved in the control of foodborne diseases by production of bacteriocins and hydrogen peroxide. This genus has great potential for use in the food industry.

Keywords: *Weissella*; Food industry; Metabolism; Ecology; Technological potential; Bacteriocins.

Resumo

O gênero *Weissella* é composto por bactérias classificadas como Gram-positivas, catalase negativas, não formadoras de esporos, morfologia cocóide ou bacilos curtos. Pertencem ao grupo das bactérias lácticas (BAL), principalmente pela produção de ácido láctico a partir da fermentação de carboidratos. As espécies de *Weissella* estão distribuídas em diferentes habitats, como solos, ordenhadeiras, cana-de-açúcar e algumas cepas com características tecnológicas interessantes podem ser isoladas de alimentos fermentados, como queijos de leite cru, vegetais fermentados e leite fermentado. Do ponto de vista da tecnologia de alimentos, algumas cepas apresentam potencial na produção de exopolissacarídeos, oligossacarídeos não digeríveis, que está além de seu potencial probiótico. Portanto, as bactérias pertencentes ao gênero *Weissella* podem ter grande importância tecnológica, estando também envolvidas no controle de doenças de origem alimentar pela produção de bacteriocinas e peróxido de hidrogênio. Este gênero possui grande potencial para uso na indústria alimentícia.

Palavras-chave: *Weissella*; Indústria alimentícia; Metabolismo; Ecologia; Potencial tecnológico; Bacteriocinas.

Resumen

El género *Weissella* está compuesto por bacterias clasificadas como gram-positivas, catalasa negativas, no formadoras de esporas, de morfología cocoide o bacilos cortos. Pertenecen al grupo de las bacterias del ácido láctico (BAL), principalmente por la producción de ácido láctico a partir de la fermentación de carbohidratos. Las especies de *Weissella* se encuentran distribuidas en distintos hábitats, como suelos, ordeñadoras, caña de azúcar y algunas cepas con interesantes características tecnológicas pueden aislarse de alimentos fermentados, como quesos elaborados a partir de leche cruda, vegetales fermentados y leche fermentada. Desde el punto de vista de la tecnología de los alimentos, algunas cepas tienen potencial en la producción de exopolisacáridos, oligosacáridos no digeribles, que está más allá de su potencial probiótico. Por tanto, las bacterias pertenecientes al género *Weissella* podrían tener una gran importancia tecnológica, estando también implicadas en el control de enfermedades transmitidas por alimentos mediante la producción de bacteriocinas y peróxido de hidrógeno. Este género tiene un gran potencial para su uso en la industria alimentaria.

Palabras clave: *Weissella*; Industria de alimentos; Metabolismo; Ecología; Potencial tecnológico; Bacteriocinas.


1. Introduction

At first, microbiology techniques for identification of genus and species were based only on phenotypic tests such as morphological and biochemical tests for typing, subtyping and identification of microbial genera, species and subspecies. However, despite presenting atypical characteristics, many microorganisms were mistakenly grouped due to the low discriminatory power of these techniques when it comes to microorganisms with little genetic variability. Nevertheless, since the 80's, the techniques of manipulation of the genetic material, allow the evaluation of the bacterial genome. Because of the advancement in the studies of molecular biology, it is now possible to construct more and more precise phylogenetic techniques, such as the evaluation of rDNA (Fakruddin & Mannan, 2013, Gandra, Gandra, de Mello & da Godoi, 2008).

In 1993, using the rDNA technique, Collins found that strains with very close phenotypic characteristics, previously classified as *Leuconostoc*, presented profiles of bands very different from others belonging to the same genus. This was discovered after the study of atypical *Leuconostoc* cultures present in fermented and dried sausages produced in Greece. These isolated microorganisms resembled the bacterium *Leuconostoc* sp. for producing D (-) lactic acid but differed in several biochemical tests. The name of this genus was proposed by Collins *et al.* (Collins, Samelis, Metaxopoulos & Wallbanks, 1993) in honor of the German microbiologist Norbert Weiss due to his many contributions in the field of research related to LAB.

Collins *et al.* (1993) used genetic sequencing of the 16S rDNA to investigate the relationship between isolated bacteria and recognized strains of the genus *Leuconostoc*. They showed that the strains analyzed were phylogenetically closer to the *Leuconostoc paramesenteroides* than to the other *Leuconostoc* species (Figure 1). They proposed that *L. paramesenteroides*, the new species isolated from the fermented sausages, and some heterofermentative *Lactobacillus*, should belong to the new genus, named *Weissella*. The name *Weissella hellenica* was proposed to name the new species.

Figure 1. Phylogenetic tree showing the proximity of the genus *Weissella* with other genera of lactic acid bacteria.



Source: Adapted from Collins (1993).

Bacteria belonging to *Weissella* genus are hardly differentiated from *Leuconostoc* strains and heterofermentative *Lactobacillus* by phenotypic characteristics; thus, the description was only possible through molecular taxonomic analyzes. Collins *et al.* (1993) proposed the reclassification of *L. paramesenteroides*, *Lb. confusus*, *Lb. halotolerans*, *Lb. kandleri*, *Lb. minor*, and *Lb. viridescens* to *W. paramesenteroides*, *W. confusa*, *W. halotolerans*, *W. kandleri*, *W. minor* and *W. viridescens*, respectively. Subsequently, other studies have identified new species, and currently 25 species are validated: *W. viridescens*, (Niven & Evans, 1956), *W. paramesenteroides* (Garvie, 1967), *W. confusa* (Holzapfel e Kandler, 1969 apud Collins *et al.*, 1993), *W. kandleri* (Holzapfel & Van Wyk, 1982), *W. halotolerans* (Kandler, Schillinger & Weiss, 1983), *W. minor* (Kandler *et al.*, 1983), *W. hellenica*

(Collins *et al.*, 1993), *W. thailandensis* (Tanasupawat, Shida, Okada, & Komagata, 2000), *W. soli* (Magnusson, Jonsson, Schnurer & Roos, 2002), *W. cibaria* (Björkroth, K. J. *et al.*, 2002), *W. koreensis* (Lee, *et al.*, 2002), *W. ghanensis* (De Bruyne, Camu, Lefebvre, De Vuyst & Vandamme, 2008), *W. beninenses* (Padonou, *et al.* 2010), *W. fabaria* (De Bruyne, Camu, De Vuyst, & Vandamme, 2010), *W. ceti* (Vela *et al.*, 2011), *W. fabalis* (Snauwaert, Papalexandratou, De Vuyst, & Vandamme, 2013), *W. oryzae* (Tohno *et al.*, 2013), *W. diestrammenae* (Oh, *et al.*, 2013), *W. uvarum* (Nisiotou, Dourou, Filippousi, Banilas, & Tassou, 2014), *W. cryptocerci* (Heo, *et al.* 2019), *W. bombi* (Praet, *et al.*, 2015), *W. jogaejeotgali* (Lee *et al.* 2015), *W. kimchi* (Choi *et al.*, 2002), *W. muntiaci* (Lin *et al.*, 2020) and *W. sagaensis* (Li, Tian & Gu, 2020).

The genus *Weissella* belongs to the phylum *Firmicutes*, class *Bacilli*, order *Lactobacillales* and family *Leuconostocaceae*. The bacteria belonging to this genus may have different morphologies, being short rods with rounded to tapered ends or coccoid in shape, which is the morphology of microorganisms belonging to the genus *Leuconostocs*, *Oenococcus* and *Streptococci* (Collins *et al.*, 1993). In some species a tendency to pleomorphism occurs depending on the stress condition to which the bacterium is submitted. In relation to the organization, they can also be found in pairs, alone or in small chains (Gandra, *et al.*, 2008, Collins *et al.*, 1993). *Weissella* is a relatively new genus, which is not yet used as a starter or adjunct culture by the food industry, as there are still few studies on its use as such. The purpose of this review is to bring the most relevant research in this genus to the food industry, presenting the potential application of the strains of the genus *Weissella* in food.

2. Methodology

This paper is a bibliographic research (Pereira, Shitsuka, Parreira & Shitsuka, 2018). Therefore, the objective is the verification of scientific data and the discussion about them on the proposed theme. This article was done in a descriptive and qualitative way in order to present how *Weissella* sp. can be applied in the food industry.

To obtain the theoretical foundation it was use the database of the websites: *Google academic*, *Scielo*, *PubMed* and *Science direct* and for the research was used keywords as *Weissella*, bacteriocin, exopolysaccharides and lactic acid bacteria. The time period used was from the year 1993, when the species was recognized, until the year 2021 as a form to compare the evolution of researches and application of the

Weissella genus and their compounds. Also for this research there was no limitation about language as a way to gather as much information as possible. By the end, was select 90 sources of information.

3. Physiological and metabolic characteristics

Bacteria from the genus *Weissella* are chemoorganotrophic, facultative anaerobic, Gram-positive, non-spore forming, catalase negative (Collins *et al.*, 1993) and have no motility with except to *W. beninenses* that has peritrichous flagella (Padonou, *et al.* 2010). All the microorganisms of this genus are compulsory heterofermenters producing lactic acid, carbon dioxide, ethanol and/or acetic acid from carbohydrate fermentation. They use the phospho-ketolase and the hexose monophosphate pathways to perform the carbohydrate fermentation (Garvie, 1967, Holzapfel & Van Wyk, 1982).

Weissella sp. have very complex nutritional needs requiring peptides, amino acids (arginine, aspartic acid, cystine, glutamic acid, histidine, isoleucine, phenylalanine, serine, threonine, tryptophan and valine), fatty acids, nucleic acids, fermentable carbohydrates (glucose, fructose, mannose, maltose, sucrose, trehalose) and vitamins (riboflavin, pyridoxal, folic acid, biotin, nicotine, thiamine, panthotenic acid) for their development (Gandra, *et al.*, 2008, Björkroth, Dicks, & Endo, 2014, Kandler *et al.*, 1983). However, culture media used as Man Rogosa and Sharp (MRS) and M17 for lactic acid bacteria are easily employed for the multiplication of these microorganisms. Moreover, all the nutritional requirements can be found in many raw materials used in the food industry such as milk, meat and vegetables, providing the use of such cultures for fermentation thereof.


As part of the LAB group, the species of the genus *Weissella* are chemoorganotrophic microorganisms with strictly fermentative metabolism being unable to synthesize porphyrinic groups (e.g. heme). All species can multiply at 15 °C while some multiply at temperatures between 42 °C and 45 °C, having an optimum temperature of multiplication between 20 °C and 30 °C (Garvie, 1967). These bacteria can be applied in fermented dairy products like fermented milk which has the temperature of incubation of the microorganisms around 40 °C and fermented meat sausages with an incubation temperature of 25 °C.

Different species produces dextran, hydrolyzed esculin and produces ammonia from arginine. Dextran is an exopolysaccharide, and these are important for the

manufacture of products such as yogurt, in which it is desired to have viscosity (Mende, Rohm, & Jaros, 2016). In addition, as these bacteria are able to ferment several sugars such as cellobiose, fructose, galactose, maltose, raffinose, ribose, trehalose and xylose (Fusco *et al.*, 2015). they can be used as starter cultures for the manufacture of products with different sensory characteristics.

W. cibaria MG1 possesses all the necessary genes for the use of the phosphoketolase pathway and metabolize the galactose via the Leloir pathway and in the figure 2 is possible to understand how it is the metabolism. It can also use maltose, fructose, ribose, xylose, sucrose and gluconate as carbon sources. In addition, a β -galactosidase has been known to indicate that lactose can also be metabolized (Lynch *et al.*, 2015). which suggests that this strain can be used in the fermentation of dairy products that have lactose as their main sugar.

Figure 2. Metabolic pathways used by bacteria of the genus *Weissella*.



Source: Own authorship (2021).

Most EPS produced by strains of the genus *Weissella* are homopolysaccharides (HoPS) such as dextran, fructan, galactan, levan and inulin for example (Björkroth, *et al.*, 2002, Lee, *et al.*, 2002). Strains of the genus *Weissella* use sucrose as the

obligatory substrate for the extracellular synthesis of HoPS. The enzymes used to hydrolyze sucrose are glycosyltransferase, glucan-sucrase (GS) or fructan-sucrase (FS), these being highly specific (Zeidan, *et al.*, 2017). The GS and FS enzymes cleave the glycosidic linkage of sucrose and couple a glucose (GS)/fructose (FS) unit to a glucan (GS)/fructan (FS) chain, water, sucrose or other acceptor (Meng, *et al.*, 2016).

The *Weissella* strains are dependent on the proteolytic system to obtain essential amino acids, which are precursors of peptides, proteins and aromatic compounds. Lynch *et al.* (2015) studied the metabolic traits of *W. cibaria* and noted that the secretome of the genus *Weissella* includes both extracellular proteins, and cell wall proteins, that are secreted. It has been found that many of the secretome proteins were large, containing multiple domains, and greater than 1000 amino acids. They also showed that *W. cibaria* and *W. confusa* have a similar number of secretory proteins as *Lb. rhamnosus* GG, which is a known probiotic strain.

The strain *W. cibaria* MG1 has the production capacity of acetate, via the pyruvate oxidase pathway in the presence of oxygen, and the production of lactic acid, diacetyl or acetoin via the diacetyl/acetoin pathway in anaerobiosis (Lynch *et al.*, 2015). Some strains of *W. confusa*, *W. paramesenteroides* and *W. cibaria* have the ability to metabolize D-xylose, glucose, D-fructose, D-mannose, sucrose, D-maltose and cellobiose and also showed β -glucosidase and β -galactosidase activity (López-Hernández, Rodríguez-Alegría, López-Munguía, & Wachter, 2017).

4. Ecology

Although *Weissella* sp. has a very complex nutritional requirement, it is found as autochthonous bacteria in different ecosystems. Due to this fact, they may contaminate food since it can spread easily in the processing environment. Different strains were isolated from soil (Padonou, *et al.* 2010), sediments of swamps (De Bruyne, *et al.*, 2010), lake water (Vela *et al.*, 2011), being identified, mainly, in fermented foods as cheese made with raw milk (Snauwaert *et al.*, 2013, Tohno *et al.*, 2013), fermented milk (Oh, *et al.*, 2013, Nisiotou, *et al.*, 2014, 85], vegetables (Wouters, Grosu-Tudor, Zamfir, & De Vuyst, 2013). and sausages (Huys, Leisner, & Björkroth, 2012, Mende, Rohm, & Jaros, 2016) in Table 1 those species and their habitat are better explained.

Table 1. Occurrence of *Weissella* species in different ecosystems.

Species	Habitat or source	Reference
<i>W. cibaria</i>	Orange, pineapple, banana	Endo, <i>et al.</i> , 2009
	Tomato	Di Cagno, <i>et al.</i> , 2009
	Wheat flour	Alfonzo, <i>et al.</i> , 2013
	Blackberry, papaya	Di Cagno, Minervini, Rizzello, De Angelis & Gobbetti, 2011
<i>W. confuse</i>	Rhizosphere of the olive trees and surrounding soil	Fhoula <i>et al.</i> , 2013
	Red and yellow raw pepper	Di Cagno, <i>et al.</i> , 2009
<i>W. halotolerans</i>	Rhizosphere of the olive trees and surrounding soil	Alfonzo, <i>et al.</i> , 2013
	Fermented sausage	Tenea & Lara, 2019
<i>W. hellenica</i>	Vegetable forage crops	Tohno, Kobayashi, Nomura, Uegaki & Cai, 2012
	Croatian cheese fermented from raw milk	Fuka, <i>et al.</i> , 2013
<i>W. kandleri</i>	Desert plants	Holzappel & Van Wyk, 1982
<i>W. uvarum</i>	Wine grapes	Nisiotou, <i>et al.</i> , 2014
<i>W. sagaensis</i>	Chinese yogurt	Li, <i>et al.</i> , 2020
<i>W. paramesenteroides</i>	Raw milk cheeses	Masoud, <i>et al.</i> , 2012
	Fermented sausage	Juárez-Castelán, <i>et al.</i> , 2019

Source: Own authorship (2021).

Lynch and colleagues investigated the genome of the *W. cibaria* species and according to the study of pan-proteome and core-proteome at the species level. Pan-proteome are all the proteins which are present in a given condition for all the species of a given life branch and the core-proteome are the proteins which are conserved in all the species of a given life branch and produced for a given condition (Trapp, *et al.*, 2016). Lynch and colleagues noted that the pan-proteome was much smaller and the core-proteome much larger in a level of genus. The fact that the core-proteome is much

larger (corresponding to 69% of the pan-proteome of the species) may explain the ability of *W. cibaria* to survive in several ecological niches where they were found since they have a higher number of proteins (729 proteins) that help in their adaptation (Lynch *et al.*, 2015).

Some strains of *Weissella* such as *W. cibaria* MG1 (Lynch *et al.*, 2015) and *W. cetti* (Ortega, *et al.*, 2018) can hydrolyze arginine, which favors its survival in environments where it is subjected to stress. A good example for this is providing a greater amount of ATP when the carbon source is scarce or producing ammonia by protecting from acid stress (Lynch *et al.*, 2015). A technological benefit of arginine deamination is the production of ornithine which is an important precursor of crust aroma compounds in the sourdough (De Angelis, *et al.*, 2002).

Because they are autochthonous in many places, some strains of *Weissella* can be important in the characterization of traditional products of certain regions. As an example, strains of *W. thailandensis* and *W. cibaria* have been related to Thai fermented fish (Björkroth, Dicks & Endo, 2014, Mende, Rohm, & Jaros, 2016), while the strains *W. cibaria*, *W. confusa* and *W. koreensis* were detected in fermented foods of vegetal origin (Fusco *et al.*, 2015, Lynch *et al.*, 2015). In addition, the bacterium *W. beninensis* was isolated from the submerged fermentation of cassava (Padonou, *et al.* 2010) and the bacteria *W. ghanensis* and *W. fabaria* were detected in piles of fermented Ghana cocoa beans (De Bruyne, *et al.*, 2010). These products may not have the same characteristics if they were manufactured without the presence of these strains.

5. Some applications of bacteria from genus *Weissella* in food

Weissella sp. has great potential in its application in food and for this reason it has been studied. Some strains show antagonistic activity against pathogens due to the production of several compounds like bacteriocins, organic acids, hydrogen peroxide, among others (Fusco *et al.*, 2015, Meng, *et al.*, 2016, López-Hernández, Rodríguez-Alegría, López-Munguía, & Wachter, 2017, Goh, & Philip, 2015, Yu *et al.*, 2019, Trias, Bañeras, Montesinos & Badosa, 2008). In table 2, the types of bacteriocin and what species are able to produce these compounds are demonstrated. *W. cibaria* TM 128 presented the production of organic acids and hydrogen peroxide, acting as inhibitors of the growth of phytopathogenic and deteriorating fungi and bacteria of fruits and vegetables (Trias, *et al.*, 2008). Some research demonstrates the antimicrobial

capacity of the compounds produced by *Weissella* against Gram-positive and Gram-negative bacteria growth (Trias, *et al.*, 2008, Kariyawasam, Jeewanthi, Lee & Paik, 2019).

Table 2. Bacteriocins produced by *Weissella* strains.

Bacteriocin	Producing species	Reference
Weissellicin 110	<i>W. cibaria</i> 110	Srionnual, Yanagida, Lin, Hsiao, & Chen, 2007
Weissellin A	<i>W. paramesenteroides</i> DX	Di Cagno, Minervini, Rizzello, De Angelis, M. & Gobbetti, 2011
Weissellicin L	<i>W. hellenica</i> 4-7	Leong, K. H. <i>et al.</i> , 2013
Weissellicin D	<i>W. hellenica</i> D1501	Chen, <i>et al.</i> , 2014
Weissellicin M	<i>W. hellenica</i> QU 13	Masuda, <i>et al.</i> , 2012
Weissellicin Y	<i>W. hellenica</i> QU 13	Kariyawasam, <i>et al.</i> , 2019

Source: Own authorship (2021).

Kariyawasam *et al.* (2019), used the strain *W. cibaria* D30 as an adjunct culture in cottage cheese manufacture and the strain increases the prevention of the growth of *Listeria monocytogenes* and ensure the microbial safety of ready-to-eat soft cheeses. Nam, Ha, Bae & Lee (2014), showed that *W. confusa* has antagonistic activity against the pathogen *Helicobacter pylori* a gram-negative microorganism that causes gastritis and gastric carcinoma, infects through the intake of food, and attaches to gastric and duodenal mucous membranes. *W. confusa* strain PL9001 inhibited the binding of *H. pylori* to human gastric-cell line MKN-45 cells by more than 90%. The results suggest that *Weissella* strains can be used as probiotics added in fermented milk, for example with the aim of fighting *H. pylori*. Besides that, *W. confusa* DD-A7 has antagonistic activity against the multidrug-resistant *Escherichia coli* which is resistant against almost all antibiotics used for its treatment. The strain *W. confusa* DD-A7 was capable to trigger an oxidative attack and limits the growth of the pathogen (Dey, Khan & Kang, 2019).

The first new bacteriocin produced by *Weissella* strains to be discovered was Weissellicin 110 in the year 2007. This compound is produced by the strain *W. cibaria*

110 isolated from the Thai fermented fish product plaa-som. This bacteriocin has antimicrobial activity against some Gram-positive microorganisms and it is resistant to high temperatures and catalase, but loses its activity when exposed to proteinase K and trypsin (Srionnual *et al.*, 2007).

In the year 2014, Chen and colleagues (Chen, *et al.*, 2014) discovered a new bacteriocin called Weissellicin D produced by the strain *W. hellenica* D1501 associated with Chinese Dong fermented meat. This bacteriocin has antimicrobial activity against the pathogenic bacteria *Staphylococcus aureus*, *L. monocytogenes* and *E. coli*. This same strain has already been tested for its antagonistic capacity against the pathogens *Kurthia gibsonii*, *S. aureus* and *E. coli* in soybean milk and was subsequently used in the manufacture of a new type of Tofu with increased shelf life due to presence of volatile antimicrobial compounds and bacteriocins (Chen, Rui, Lu, Li & Dong, 2014).

Besides of Weissellicin D, others isolate of *W. hellenica* demonstrated the production of Weissellicins L, M and Y which showed antagonist activity against *L. monocytogenes* and *Bacillus coagulans* (Ayeni, *et al.*, 2011, Masoud, *et al.*, 2012). Bacteriocins are natural antimicrobial compounds, there is an interest in their use by the food industry for the purpose of application as bioconservants, i.e. natural preservatives and possible substitutes for chemical preservatives (O'Connor, *et al.*, 2012). Some characteristics that make bacteriocins that are produced by lactic acid bacteria safe when used at industrial level are their non-toxicity to eukaryotic cells inactivation by digestive proteases, little influence on the intestinal microbiota (Jawan *et al.*, 2019, Wouters *et al.*, 2013) tolerance to different temperatures and pHs, action against pathogens and microorganisms spoilage of food, and do not generating cross-resistance to antibiotics (Wouters *et al.*, 2013, Juárez-Castelán, *et al.*, 2019, Tenea & Lara, 2019).

Bacteriocins are also of great importance in the food industry, for example in the production of biodegradable food packaging with antimicrobial properties (Teixeira, *et al.*, 2021). By incorporating the Bacteriocin 7293 produced by the strain *W. hellenica* BCC 7293, it was possible to control pathogenic bacteria in fillets of pangasius fish. The film produced inhibited the multiplication of both Gram-positive bacteria such as *L. monocytogenes* and *S. aureus* as well as Gram-negative bacteria such as *Pseudomonas aeruginosa*, *Aeromonas hydrophila*, *E. coli* and *Salmonella Typhimurium* (Woraprayote, *et al.* 2018).

Weissella can also be useful for the food industry due to its capacity of some strains to produce EPS in sucrose culture medium (Woraprayote, *et al.* 2018). EPS are structures of high molecular mass, composed of carbohydrates, which when added to the food, behave mainly as thickening and emulsifying agents. The polysaccharides that adds thickening and gelling properties are irreplaceable in the food industry formulations. In addition to its ability as probiotics, EPS imposes highly desirable rheological changes in the food matrix, such as viscosity increase, improved texture and reduced syneresis (Lakra, Domdi, Tilwani & Arul,2020). They can be used to replace corn starch in manufacturing puddings, for example, where these characteristics are desired. Some studies have been tested strategies to improve the EPS characteristics in order to enlarge its application in the food industry (Kavitake, *et al.*, 2019). Most of the strains of *Weissella* have the capacity to produce the EPS dextran and can also produce other EPS as shown in Table 3.

Table 3. Exopolysaccharides produced by *Weissella* strains.

Species	Product	Reference
<i>Weissella cibaria</i> RBA12	Dextran	Baruah, Maina, Katina, Juvonen & Goyal, 2017
<i>W. confusa</i>	Dextran	Kajala I. <i>et al.</i> , 2016
<i>Weissella confusa</i> KR780676	Linear exopolysaccharide galactan	Devi, Kavitake & Shetty, 2016, Kavitake, Devi & Shetty, 2016
<i>W. confusa</i> and <i>W. cibaria</i>	Dextran, fructan from sucrose, ropy capsular polysaccharide, levan and inulin	Malang, Maina, Schwab, Tenkanen & Lacroix, 2015
<i>W. confusa</i> XG-3	Dextran	Zhao, <i>et al.</i> , 2021
<i>W. cibaria</i> MD2	Fructan	Lakra,, Ramatchandirane, Kumar, Suchiang & Arul, 2021

Source: Own authorship (2021).

The EPS dextran has great efficacy as soluble dietary fiber since it has the capacity to be fermented by the probiotic intestinal microbiota due to its low digestibility when compared to commercial prebiotic inulin (Teixeira, *et al.*, 2021). The basis for

EPS production by *Weissella* strains is diverse, some studies use wheat flour and rye as the basis for fermentation. *W. confusa* presented higher rye meal production due to the higher optimum pH period for the synthesis of dextran during fermentation (Cotter, Ross & Hill, 2013, Kavitake, *et al.*, 2019).

The strains that have the capacity to produce more than one exopolysaccharide are of great interest when it comes to the food industry because of their synergistic effect on the texture and also the nutritional improvement (Malang, *et al.*, 2015). Some EPS that are being produced, such as galactan, has good emulsifying and stabilizing capacities which provides its future use in cosmetic and food emulsions (Ortega, *et al.*, 2018, De Angelis, *et al.*, 2002, Kavitake, Balvan, Devi & Shetty, 2020). The high molecular weight dextran produced by *W. confusa* QS813 can form hydrogens bonds and steric interactions with proteins. When this EPS is used in frozen dough it's maintained the structural integrity of gluten during the freeze-thaw-cycles being appropriated to use as a cryoprotectant in wheat gluten-based frozen food (Tang, *et al.*, 2019).

Because the production of exopolysaccharide, the application of *W. cibaria* as an assistant in the manufacture of cheddar cheese was performed resulting in a product with higher retention of humidity and without alteration of the proteolysis degree (Lynch, *et al.*, 2014). In addition, *W. cibaria* MG1 produces exopolysaccharides (dextran) and oligosaccharides (glucopolysaccharides) and because of that was studies with the intention of producing a new fermented drink from wort sucrose-supplemented barley-malt-derived (Zannini, *et al.*, 2013). The oligosaccharides are being applied in some foods as prebiotics, when it is impossible to use probiotics, an example of this are infant formulas. In the developing countries, infant formulas are heated before consumption, because of their dubious quality of the water. In this way, the addition of probiotics is impossible, and prebiotics are added instead. The oligosaccharides produced by *Weissella* can be a source of prebiotics for the food industry.

Rosca *et al.* (2018), verified that *W. confusa* produced a dextran with high structural stability and purity that have pharmaceutical importance due to its antifungal characteristics against the pathogenic yeast *Candida albicans*. Some strains such as the *W. cibaria* strain isolated from goat's milk also have been shown to have great potential as probiotics, such as resistance to 1% bile salt and tolerance to pH 3.0 (Elavarasi, Pugazhendhi, Poornima Priyadharsani, Valsala, & Thamaraiselvi, 2014). In

addition, *W. cibaria*, jointly with *Lb. plantarum*, showed *in vitro* tests high antioxidant capacity, survived simulated gastric and intestinal transit, and tolerated bile acids and salts (Yu *et al.*, 2019). Furthermore, the two strains were administered to male Wistar albino rats and showed an improvement in liver and kidney functions, damaged by heavy metals compared to rats that received only the heavy metals in their diet (Ojekunle, Banwo, & Sanni, 2017).

In addition, the *Weissella* species-pair *W. cibaria/confusa*, considered potentially probiotic by Immerzeel *et al.* (2014), were studied in relation to the use of xylooligosaccharides (XOS) as carbon source, since these are considered prebiotic. The study showed that strains absorbed XOS, both xylobiose and xylotriose, and gave an increase in the production of lactic acid when xylan hydrolyzed was used. Not only but some stains can also use xylooligosaccharides (XOS) to produce short-chain fatty acids a feature only observed in *Leuconostoc lactis* and a few strains of the established probiotics *Lactobacillus* (Månberger, *et al.*, 2020).

Adesulu-Dahunsi *et al.* (2018) suggest that EPS produced by *W. cibaria* GA44 may be a commercial alternative for the food industry once it presents strong properties as antioxidant when compared to commercial antioxidant ascorbic acid, especially scavenging of superoxide anions and hydroxyl radicals.

On the other hand, the EPS glucansucrase produced by *Weissella* sp. TN610 has the ability to solidify semi-skimmed milk supplemented with sucrose which shows its potential in the application as a safe additive food to improve the texture of dairy products (Bejar, *et al.*, 2013). Besides that, a novel quinoa-based yoghurt fermented with dextran producer *W. cibaria* MG1 was developed and the concentration of EPS (40 mg/L) ensured the high-water retention capacity and viscosity (0.57 mPa s) of the final product (Zaninni, Jeske, Lynch & Arendt, 2018).

The EPS dextran, levan and ropy capsular polysaccharide, produced by *W. confusa* were evaluated in breads when it comes to the delaying of the deterioration by fungus, as well as improving of the texture were observed (Tinzi-Malang, Rast, Grattepanche, Sych, & Lacroix, 2015). In addition to the production of EPS, some strains of *W. cibaria* and *W. confusa* as well as producing lactic acid also have the capacity to produce folate (vitamin B9), which allows the nutritional improvement of fermented products that use these strains in the fermentation process (Deatraksa *et al.*, 2018).

6. Conclusion

Weissella sp. may have a wide range of potential applications in food products, since they can produce a high variety of compounds from the production of EPS, bacteriocins, even vitamin such as B9. However, a few studies have been conducted and the industrial application of *Weissella* strains it is still not a reality. There are still some obstacles that prevent the use of *Weissella* as a starter culture in the food industry, such as the lack of knowledge about the pathogenicity of some strains of the genus and its antagonistic capacity against other microorganisms of industrial interest due to its production of bacteriocins. Because some strains have a potential for opportunistic infection in humans, the food industry should always be vigilant for safety testing of any strain before its technological application.

Thereby, for the future use of this potential species in food industry, is necessary more researches about how the *Weissella* sp. and their compounds can be apply and with this review article it's possible to understand that it's necessary more research about the safety use of this strains in food.

7. References

- Fakruddin, M. & Mannan, KSB. (2013). Methods for Analyzing Diversity of Microbial Communities in Natural Environments. *Ceylon Journal of Science (Biological Sciences)*, 42 (1), 19-33.
- Gandra, E. A.; Gandra, T. K. V.; de Mello, W. S. & da Godoi, H. 2008. Técnicas moleculares aplicadas à microbiologia de alimentos. *Acta Scientiarum - Technology*, 30 (1), 109–118.
- Collins M. D.; Samelis, J.; Metaxopoulos, J. & Wallbanks, S. (1993). Taxonomic studies on some leuconostoc-like organisms from fermented sausages: description of a new genus *Weissella* for the *Leuconostoc paramesenteroides* group of species. *Journal of Applied Bacteriology*, 75(6), 595–603.
- Niven, C. F. & Evans, J. B. (1956). Species That Produces a Green Discoloration of Cured Meat Pigments. *Journal Bacteriology*, 758–759.
- Garvie, E. I. (1967). The growth factor and amino acid requirements of species of the genus *Leuconostoc*, including *Leuconostoc paramesenteroides* (sp. nov.) and *Leuconostoc oenos*. *Journal of General Microbiology*, 48(1967), 439–447.
- Holzappel, W. H. & Van Wyk, E. P. (1982). *Lactobacillus kandleri* sp. nov., a New Species of the Subgenus Betabacterium, with Glycine in the Peptidoglycan," *Zentralblatt für Bakteriologie Mikrobiologie und Hygiene: I. Abt. Originale C: Allgemeine, angewandte und ökologische Mikrobiologie*, 3(4), 495–502.
- Kandler, O.; Schillinger, U. & Weiss, N. (1983). *Lactobacillus halotolerans* sp.nov., nom.rev. and *Lactobacillus minor* sp.nov., nom.rev. *Systematic and Applied Microbiology*, 4(2), 280–285.

Tanasupawat, S.; Shida, O.; Okada, S. & Komagata, K. (2000). *Lactobacillus acidipiscis* sp. nov. and *Weissella thailandensis* sp. nov., isolated from fermented fish in Thailand," *International Journal of Systematic and Evolutionary Microbiology*, 50, 1479–1485.

Magnusson, J.; Jonsson, H.; Schnurer, J. & Roos, S. (2002). *Weissella soli* sp. nov., a lactic acid bacterium isolated from soil. *International Journal of Systematic and Evolutionary Microbiology*, 52(3), 831–834.

Björkroth, K. J.; Schillinger, U.; Geisen, R.; Weiss, N.; Hoste, B.; Holzappel, H. W.; Korkeala, H. J.; Vandamme, P., (2002). Taxonomic study of *Weissella confusa* and description of *Weissella cibaria* sp. nov., detected in food and clinical samples. *International Journal of Systematic and Evolutionary Microbiology*, 52 (1), 141–148.

Lee, J.-S.; Lee, K. C.; Ahn, J.-S.; Mheen, T.-IY.; Pyun, R. & Park, Y.H. (2002). *Weissella koreensis* sp. nov., isolated from kimchi. *International Journal of Systematic and Evolutionary Microbiology*, 52, 1257–1261.

De Bruyne, K.; Camu, N.; Lefebvre, K.; De Vuyst, L., & Vandamme, P. (2008). *Weissella ghanensis* sp. nov., isolated from a Ghanaian cocoa fermentation. *International Journal of Systematic and Evolutionary Microbiology*, 58 (2008), 2721–2725.

Padonou S. W.; Schillinger, U.; Nielsen, D. S.; Franz, C. M. A. P.; Hansen, M.; Hounhouigan, J. D.; Nago, M. C.; Jakobsen, M. *Weissella beninensis* sp. nov., a motile lactic acid bacterium from submerged cassava fermentations, and emended description of the genus *Weissella*. *International Journal of Systematic and Evolutionary Microbiology*, 60 (9), 2193–2198.

De Bruyne, K.; Camu, N.; De Vuyst, L. & Vandamme, P. (2010). *Weissella fabaria* sp. nov., from a Ghanaian cocoa fermentation. *International Journal of Systematic and Evolutionary Microbiology*, 60 (2010), 1999–2005.

Vela, A. I.; Fernández, A.; de Quirós, Y. B.; Herráez, P.; Domínguez, L. & Fernández-Garayzábal, J. F. (2011). *Weissella ceti* sp. nov., isolated from beaked whales (*Mesoplodon bidens*). *International Journal of Systematic and Evolutionary Microbiology*, 61(11), 2758–2762.

Snauwaert, I.; Papalexandratou, Z.; De Vuyst, L. & Vandamme, P. (2013). Characterization of strains of *Weissella fabalis* sp. nov. and *Fructobacillus tropaeoli* from spontaneous cocoa bean fermentations," *International Journal of Systematic and Evolutionary Microbiology*, 63(PART 5), 1709–1716.

Tohno, M.; Kitahara, M.; Inoue, H.; Uegaki, R.; Irisawa, T.; Ohkuma, M.; Tajima, K. (2013). *Weissella oryzae* sp. nov., isolated from fermented rice grains, *International Journal of Systematic and Evolutionary Microbiology*, 63 (PART4), 1417–1420.

Oh, S. J.; Shin, N.; Hyun, D.; Kim, P. S.; Kim, J. Y.; Kim, M.; Yun, J.; Bae, J. (2013). *Weissella diestrammenae* sp. nov., isolated from the gut of a camel cricket (*Diestrammena coreana*). *International Journal of Systematic and Evolutionary Microbiology*, 63 (PART8), 2951–2956.

Nisiotou, A.; Dourou, D.; Filippousi, M.-E.; Banilas, G. & Tassou, C. (2014). *Weissella uvarum* sp. nov., Isolated from wine grapes, *International Journal of Systematic and Evolutionary Microbiology*, 64 (2014), 3885–3890.

Heo, J.; Hamada, M.; Cho, H.; Weon, H.; Kim, J.; Hong, S.; Kim, S.; Kwon, S., (2019). *Weissella cryptocerci* sp. nov., isolated from gut of the insect *Cryptocercus kyebangensis*.

- International Journal of Systematic and Evolutionary Microbiology*, 69 (7),10–16.
- Praet, J.; Meeus, I.; Cnockaert, M.; Houf, K.; Smagghe, G. & Vandamme, P. (2015). Novel lactic acid bacteria isolated from the bumble bee gut : *Convivina intestini* gen. nov., sp. nov., *Lactobacillus bombicola* sp. nov., and *Weissella bombi* sp. nov. *Antonie van Leeuwenhoek, Journal of Microbiology*, 107 (5), 1337–1349.
- Lee, S. H., Ku, H. J., Ahn, M. J., Hong, J. S., Lee, S. H., Shin, H., ... & Lee, J. H. (2015). *Weissella jogaejeotgali* sp. nov., isolated from jogae jeotgal, a traditional Korean fermented seafood. *International journal of systematic and evolutionary microbiology*, 65(12), 4674-4681.
- Choi, H. J., Cheigh, C. I., Kim, S. B., Lee, J. C., Lee, D. W., Choi, S. W., ... & Pyun, Y. R. (2002). *Weissella kimchii* sp. nov., a novel lactic acid bacterium from kimchi. *International Journal of Systematic and Evolutionary Microbiology*, 52(2), 507-511.
- Huys, G.; Leisner, J. & Björkroth, J. (2012). The Lesser LAB Gods: *Pediococcus*, *Leuconostoc*, *Weissella*, *Carnobacterium*, and Affiliated Genera. in *Lactic Acid Bacteria, Microbiological and Funcional Aspects*. 94–112.
- Björkroth, J.A. & Dicks, L. M. T. & Holzapfel, W. H. (2009). Genus III. *Weissella* Collins, Samelis, Metaxopoulos and Wallbanks 1994, 370VP. *Bergey's Manual of Systematic Bacteriology*. 3. 643-654.
- Björkroth, J.; Dicks, L. M. T. & Endo, A. (2014). The genus *Weissella*, in *Lactic Acid Bacteria, Biodiversity and Taxonomy*, First., W. H. Holzapfel and B. J. B. Wood, Eds. UK: Wiley Blackwell, pp. 417–421.
- Mende, S.; Rohm, H. & Jaros, D. (2016). Influence of exopolysaccharides on the structure, texture, stability and sensory properties of yoghurt and related products. *International Dairy Journal*, 52, 57–71.
- Fusco, V., Quero, G. M., Cho, G. S., Kabisch, J., Meske, D., Neve, H., ... & Franz, C. M. (2015). The genus *Weissella*: taxonomy, ecology and biotechnological potential. *Frontiers in microbiology*, 6, 1.
- Lynch, K. M.; Lucid, A.; Arendt, E. K.; Sleator, R. D.; Lucey, B. & Coffey, A. (2015). Genomics of *Weissella cibaria* with an examination of its metabolic traits. *Microbiology*, 161 (December 2014), pp. 914–930.
- Torino, M. I.; de Valdez, G. F. & Mozzi, F. (2015). Biopolymers from lactic acid bacteria. Novel applications in foods and beverages. *Frontiers in Microbiology*, 6 (JUL), 1–16.
- Zeidan, A. A., Poulsen, V. K., Janzen, T., Buldo, P., Derkx, P. M., Øregaard, G., & Neves, A. R. (2017). Polysaccharide production by lactic acid bacteria: from genes to industrial applications. *FEMS microbiology reviews*, 41(Supp_1), S168-S200.
- Meng, X., Gangoiti, J., Bai, Y., Pijning, T., Van Leeuwen, S. S., & Dijkhuizen, L. (2016). Structure–function relationships of family GH70 glucansucrase and 4, 6- α -glucanotransferase enzymes, and their evolutionary relationships with family GH13 enzymes. *Cellular and Molecular Life Sciences*, 73(14), 2681-2706.
- López-Hernández, M.; Rodríguez-Alegría, M. E.; López-Munguía, A. & Wachter, C. (2017). Evaluatin of Xylan as Carbon Source for *Weissella* spp., a Predominant Strain in Pozol Fermentation. *LWT - Food Science and Technology*, 89 (March), 192–197.

Goh, H. F. & Philip, K. (2015). Purification and Characterization of Bacteriocin Produced by *Weissella confusa* A3 of Dairy Origin. *PLoS ONE*.

Yu, H. S., Lee, N. K., Choi, A. J., Choe, J. S., Bae, C. H., & Paik, H. D. (2019). Antagonistic and antioxidant effect of probiotic *Weissella cibaria* JW15. *Food science and biotechnology*, 28(3), 851-855.

Ayeni, F. A.; Sánchez, B.; Adeniyi, B. A.; de los Reyes-Gavilán, C. G.; Margolles, A. & Ruas-Madiedo, P. (2011). Evaluation of the functional potential of *Weissella* and *Lactobacillus* isolates obtained from Nigerian traditional fermented foods and cow's intestine. *International Journal of Food Microbiology*, 147 (2), 97–104.

Masoud, W.; Vogensen, F. K.; Lillevang, S.; Abu Al-Soud, W.; Sørensen, S. J. & Jakobsen, M. (2012). The fate of indigenous microbiota, starter cultures, *Escherichia coli*, *Listeria innocua* and *Staphylococcus aureus* in Danish raw milk and cheeses determined by pyrosequencing and quantitative real time (qRT)-PCR. *International Journal of Food Microbiology*, vol. 153(1–2), 192–202.

Jawan, R., Kasimin, M.E., Jalal, S.N., Mohd Faik, A.A., Abbasiliasi, S., Ariff, A., (2019). Isolation, characterisation and in vitro evaluation of bacteriocins-producing lactic acid bacteria from fermented products of Northern Borneo for their beneficial roles in food industry. *Journal of Physics: Conference Series*. Ser. 1358.

Wouters, D.; Grosu-Tudor, S.; Zamfir, M. & De Vuyst, L. (2013). Bacterial community dynamics, lactic acid bacteria species diversity and metabolite kinetics of traditional Romanian vegetable fermentations. *Journal of the Science of Food and Agriculture*, 93 (4), 4749–760.

Juárez-Castelán, C., García-Cano, I., Escobar-Zepeda, A., Azaola-Espinosa, A., Álvarez-Cisneros, Y. & Ponce-Alquicira, E. (2019). Evaluation of the bacterial diversity of Spanish-type chorizo during the ripening process using high-throughput sequencing and physicochemical characterization. *Meat Sci.* 150, 7–13.

Tenea, G. N. & Lara, M. I. (2019). Antimicrobial compounds produced by *Weissella confusa* Cys2-2 strain inhibit Gram-negative bacteria growth, CyTA - *Journal of Food*, 17:1, 105-111.

Trapp, J., Almunia, C., Gaillard, J. C., Pible, O., Chaumot, A., Geffard, O., & Armengaud, J. (2016). Proteogenomic insights into the core-proteome of female reproductive tissues from crustacean amphipods. *Journal of proteomics*, 135, 51-61.

Castrejón-Nájera, J., Ortega, C., Fajardo, R., Irgang, R., Tapia-Cammas, D., Poblete-Morales, M., & Avendaño-Herrera, R. (2018). Isolation characterization, virulence potential of *Weissella cети* responsible for weissellosis outbreak in rainbow trout (*Oncorhynchus mykiss*) cultured in Mexico. *Transboundary and emerging diseases*, 65(6), 1401-1407.

De Angelis, M., Mariotti, L., Rossi, J., Servili, M., Fox, P. F., Rollán, G., & Gobbetti, M. (2002). Arginine catabolism by sourdough lactic acid bacteria: purification and characterization of the arginine deiminase pathway enzymes from *Lactobacillus sanfranciscensis* CB1. *Applied and Environmental Microbiology*, 68(12), 6193-6201.

Srionnual, S.; Yanagida, F.; Lin, L.-H.; Hsiao, K.-N. & Chen, Y.-S. (2007). Weissellicin 110, a newly discovered bacteriocin from *Weissella cibaria* 110, isolated from Plaasom, a fermented fish product from Thailand. *Applied and Environmental Microbiology*, 73 (7), 2247–2250.

Masuda, Y.; Zendo, T.; Sawa, N.; Perez, R. H.; Nakayama, J. & Sonomoto, K. (2012). Characterization and identification of weissellicin Y and weissellicin M, novel bacteriocins

produced by *Weissella hellenica* QU 13. *Journal of Applied Microbiology*, 112 (1), 99–108.

Leong, K. H., Chen, Y. S., Lin, Y. H., Pan, S. F., Yu, B., Wu, H. C., & Yanagida, F. (2013). Weissellicin L, a novel bacteriocin from sian-sianzih-isolated *Weissella hellenica* 4-7. *Journal of applied microbiology*, 115(1), 70-76.

Chen, C., Chen, X., Jiang, M., Rui, X.; Li, W. & Dong, M. (2014). A newly discovered bacteriocin from *Weissella hellenica* D1501 associated with Chinese Dong fermented meat (Nanx Wudl). *Food Control*, 42 (September 2013), 116–124.

Trias Mansilla, R., Bañeras Vives, L., Montesinos Seguí, E., & Badosa Romañó, E. (2008). Lactic acid bacteria from fresh fruit and vegetables as biocontrol agents of phytopathogenic bacteria and fungi. *International Microbiology*, 2008, núm. 11. p. 231-236.

Kariyawasam, K. M. G. M. M., Jeewanthi, R. K. C., Lee, N.-K. & Paik, H.-D. (2019). Characterization of cottage cheese using *Weissella cibaria* D30: Physicochemical, antioxidant, and antilisterial properties. *Journal of Dairy Science*, 102, 1–7.

Nam, H., Ha, M., Bae, O. & Lee, Y. (2002). Effect of *Weissella confusa* Strain PL9001 on the Adherence and Growth of *Helicobacter pylori*. *Applied and Environmental Microbiology*, 68(9), 4642–4645.

Dey, D. K., Khan, I. & Kang, S. C. (2019). Anti-bacterial susceptibility profiling of *Weissella confusa* DD _ A7 against the multidrug-resistant ESBL-positive *E. coli*. *Microbial Pathogenesis*, 128 (November 2018), 119–130.

Chen, C., Rui, X., Lu, Z., Li, W., & Dong, M. (2014). Enhanced shelf-life of tofu by using bacteriocinogenic *Weissella hellenica* D1501 as bioprotective cultures. *Food Control*, 46, 203-209.

O'Connor, P. M., Kuniyoshi, T. M., Oliveira, R.P., Hill, C., Ross, R.P. & Cotter, P. D. (2020). Antimicrobials for food and feed; a bacteriocin perspective. *Curr. Opin. Biotechnol.* 61, 160–167.

Cotter, P. D., Ross, R. P. & Hill, C. (2013). Bacteriocins-a viable alternative to antibiotics. *Nature Reviews Microbiology*, 11 (2), 95–105.

Woraprayote, W., Pumpuang, L., Tosukhowong, A., Zendo, T., Sonomoto, K., Benjakul, S., & Visessanguan, W. (2018). Antimicrobial biodegradable food packaging impregnated with Bacteriocin 7293 for control of pathogenic bacteria in pangasius fish fillets. *LWT - Food Science and Technology*, 89 (April 2017), 427–433.

Lakra, A. K., Domdi, L., Tilwani Y. M. & Arul, V. (2020). Physicochemical and functional characterization of mannan exopolysaccharide from *Weissella confusa* MD1 with bioactivities. *International Journal of Biological Macromolecules*, 143, 797–805.

Kavitake, D., Techi, M., Abid, U. K., Kandasamy, S., Devi, P. B. & Shetty, P. H. (2019). Effect of c-irradiation on physico-chemical and antioxidant properties of galactan exopolysaccharide from *Weissella confusa* KR780676. *Journal of Food Science and Technology*, 10 (88), 1–9.

Baruah, R., Maina, N. H., Katina, K., Juvonen, R., & Goyal, A. (2017). Functional food applications of dextran from *Weissella cibaria* RBA12 from pummelo (*Citrus maxima*). *International journal of food microbiology*, 242, 124-131.

Kajala, I., Mäkelä, J., Coda, R., Shukla, S., Shi, Q., Maina, N. H., Juvonen, R., Ekholm, P.,

Goyal, A., Tenkanen, M. & Katina, K. (2016). Rye bran as fermentation matrix boosts in situ dextran production by *Weissella confusa* compared to wheat bran. *Applied Microbiology and Biotechnology*, 100 (8), 3499–3510.

Malang, S. K.; Maina, N. H.; Schwab, C.; Tenkanen, M. & Lacroix, C. (2015). Characterization of exopolysaccharide and ropy capsular polysaccharide formation by *Weissella*. *Food Microbiology*, 46, 418–427.

Devi, P. B., Kavitate, D. & Shetty, P. H. (2016). Physico-chemical characterization of galactan exopolysaccharide produced by *Weissella confusa* KR780676. *International Journal of Biological Macromolecules*, 93 (Part A), 822–828.

Kavitate, D., Devi, P. B. & Shetty, P. H. (2016). Characterization of a novel galactan produced by *Weissella confusa* KR780676 from an acidic fermented food. *International Journal of Biological Macromolecules*, 86 (May 2016), 681–689.

Tang, X., Zhang, B., Huang, W., Ma, Z., Zhang, F., Wang, F., Zou, Q. & Zheng, J. (2019). Food Hydrocolloids Hydration, water distribution and microstructure of gluten during freeze thaw process: Role of a high molecular weight dextran produced by *Weissella confusa* QS813. *Food Hydrocolloids*, 90(October 2018), 377–384.

Lynch, K. M., McSweeney, P. L. H., Arendt, E. K., Uniacke-Lowe, T., Galle, S. & Coffey, A. (2014). Isolation and characterisation of exopolysaccharide-producing *Weissella* and *Lactobacillus* and their application as adjunct cultures in Cheddar cheese. *International Dairy Journal*, 34 (1), 125–134.

Zannini, E., Mauch, A., Galle, S., Gänzle, M., Coffey, A., Arendt, E. K., Taylor, J. P. & Waters, D. M. (2013). Barley malt wort fermentation by exopolysaccharide-forming *Weissella cibaria* MG1 for the production of a novel beverage. *Journal of Applied Microbiology*, 115 (6), 1379–1387.

Rosca, I., Petrovici, A. R., Peptanariu, D., Nicolescu, A., Dodi, G., Avadanei, M., Inavoc, I. C., Bostanaru, A. C., Mares, M. & Ciolacu, D. (2018). Biosynthesis of dextran by *Weissella confusa* and its *In vitro* functional characteristics. *International Journal of Biological Macromolecules*, 107 (July 2017), 1765–1772.

Elavarasi, V., Pugazhendhi, A., Poornima Priyadharsani, T. K., Valsala, H. & Thamaraiselvi, K. (2014). Screening and Characterization of *Weissella cibaria* Isolated from Food Source for Probiotic Properties. *International Journal of Computer Applications*, 1 (May 2014), 29–32.

Ojekunle, O., Banwo, K., & Sanni, A. I. (2017). *In vitro* and *In vivo* evaluation of *Weissella cibaria* and *Lactobacillus plantarum* for their protective effect against cadmium and lead toxicities. *Letters in Applied Microbiology*, 64 (5), 379–385.

Immerzeel, P., Falck, P., Galbe, M., Adlercreutz, P., Nordberg Karlsson, E. & Stålbrand, H. (2014). Extraction of water-soluble xylan from wheat bran and utilization of enzymatically produced xylooligosaccharides by *Lactobacillus*, *Bifidobacterium* and *Weissella* spp. *LWT - Food Science and Technology*, 56 (2), 321–327.

Adesulu-dahunsi, A. T., Sanni, A. I. & Jeyaram, K. (2018). Production, characterization and *In vitro* antioxidant activities of exopolysaccharide from *Weissella cibaria* GA44. *LWT - Food Science and Technology*, 87 (2018), 432–442.

Bejar, W., Gabriel, V., Amari, M., Morel, S., Mezghani, M., Maguin, E., Fontagné-Faucher, C., Bejar, S., & Chouayekh, H. (2013). Characterization of glucansucrase and dextran from

Weissella sp. TN610 with potential as safe food additives. *International Journal of Biological Macromolecules*, 52 (2013), 125–132.

Zannini, E., Jeske, S., Lynch, K. & Arendt, E. K. (2018). Development of novel quinoa-based yoghurt fermented with dextran producer *Weissella cibaria* MG1. *International Journal of Food Microbiology*, 268 (December 2017), 19–26.

Tinzi-Malang, S. K., Rast, P., Grattepanche, F., Sych, J. & Lacroix, C. (2015). Exopolysaccharides from co-cultures of *Weissella confusa* 11GU-1 and *Propionibacterium freudenreichii* JS15 act synergistically on wheat dough and bread texture. *International Journal of Food Microbiology*, 214 (April 2015), 91–101.

Deatraksa, J., Sunthornthummas, S., Rangsiruji, A., Sarawaneeyaruk, S., Suwannasai, N. & Pringsulaka, O. (2018). Isolation of folate-producing *Weissella* spp. from Thai fermented fish (Plaas Som Fug). *LWT - Food Science and Technology*, 89 (October 2017), 388–391.

Endo, A., Futagawa-Endo, Y., Kawasaki, S., Dicks, L. M. T., Niimura, Y. & Okada, S. (2009). Sodium acetate enhances hydrogen peroxide production in *Weissella cibaria*. *Letters in Applied Microbiology*, 49 (1), 136–141.

Di Cagno, R., Surico, R. F., Minervini, G., De Angelis, M., Rizzello, C. G. & Gobbetti, M. (2009). Use of autochthonous starters to ferment red and yellow peppers (*Capsicum annum* L.) to be stored at room temperature,” *International Journal of Food Microbiology*, 130 (2), 108–116.

Alfonzo, A., Ventimiglia, G., Corona, O., Di Gerlando, R., Gaglio, R., Francesca, N., Moschetti, G. & Settanni, L. (2013). Diversity and technological potential of lactic acid bacteria of wheat flours. *Food Microbiology*, 36 (2), 343–354.

Di Cagno, R., Minervini, G., Rizzello, C. G., De Angelis, M. & Gobbetti, M. (2011). Effect of lactic acid fermentation on antioxidant, texture, color and sensory properties of red and green smoothies. *Food Microbiology*, 28 (5), 1062–1071.

Fhoula, I., Najjari, A., Turki, Y., Jaballah, S., Boudabous, A. & Ouzari, H. (2013). Diversity and Antimicrobial Properties of Lactic Acid Bacteria Isolated from Rhizosphere of Olive Trees and Desert Truffles of Tunisia. *BioMed Research International*, 2013, 14.

Tohno, M., Kobayashi, H., Nomura, M., Uegaki, R. & Cai, Y. (2012). Identification and characterization of lactic acid bacteria isolated from mixed pasture of timothy and orchardgrass, and its badly preserved silage. *Animal Science Journal*, 83 (4), 318–330.

Fuka, M. M., Wallisch, S., Engel, M., Welzl, G., Havranek, J. & Schloter, M. (2013). Dynamics of bacterial communities during the ripening process of different Croatian cheese types derived from raw ewe’s milk cheeses. *PLoS ONE*, 8 (11), 1–10.

Lin S.T., Wang L.T., Wu Y.C., Guu J.J., Tamura T., Mori K., Huang L., Watanabe K. (2020). *Weissella muntiaci* sp. nov., isolated from faeces of Formosan barking deer (*Muntiacus reevesi*). *International Journal of Systematic and Evolutionary Microbiology*, 70(3), 1578-1584.

Li Y. Q., Tian W. L., Gu C. T. (2020). *Weissella sagaensis* sp. nov., isolated from traditional Chinese yogurt. *International Journal of Systematic and Evolutionary Microbiology*, 70(4), 2485-2492.

Månberger, A., Verbrugghe, P., Guðmundsdóttir, E.E. et al. (2020). Taxogenomic assessment and genomic characterisation of *Weissella cibaria* strain 92 able to metabolise oligosaccharides derived from dietary fibres. *Sci Rep* 10, 5853.

Teixeira, C. G., Fusieger, A., Milião, G. L., Martins, E., Drider, D., Nero, L. A., & de Carvalho, A. F. (2021). *Weissella*: An Emerging Bacterium with Promising Health Benefits. *Probiotics and Antimicrobial Proteins*, 1-11.

Kavitake, D., Balyan, S., Devi, P. B., & Shetty, P. H. (2020). Evaluation of oil-in-water (O/W) emulsifying properties of galactan exopolysaccharide from *Weissella confusa* KR780676. *Journal of food science and technology*, 57(4), 1579-1585.

Zhao, D., Jiang, J., Liu, L., Wang, S., Ping, W., & Ge, J. (2021). Characterization of exopolysaccharides produced by *Weissella confusa* XG-3 and their potential biotechnological applications. *International Journal of Biological Macromolecules*.

Lakra, A. K., Ramatchandirane, M., Kumar, S., Suchiang, K., & Arul, V. (2021). Physico-chemical characterization and aging effects of fructan exopolysaccharide produced by *Weissella cibaria* MD2 on *Caenorhabditis elegans*. *LWT*, 143, 111100.

Pereira, A. S., Shitsuka, D. M., Parreira, F. J., & Shitsuka, R. (2018). *Metodologia da pesquisa científica*. [ebook].

PART B: *Weissella*: An Emerging Bacterium with Promising Health Benefits

The content of this part has been published in:

Probiotics and Antimicrobial Proteins

Camila Gonçalves Teixeira, Andressa Fusieger, Gustavo Leite Milião,
Evandro Martins, Djamel Drider, Luís Augusto Nero, Antônio Fernandes de
Carvalho

License to reuse this content in thesis is in Annex 2

Abstract

Weissella strains have been the subject of much research over the last 5 years because of the genus' technological and probiotic potential. Certain strains have attracted the attention of the pharmaceutical, medical, and food industries because of their ability to produce antimicrobial exopolysaccharides (EPSs). Moreover, *Weissella* strains are able to keep foodborne pathogens in check because of the bacteriocins, hydrogen peroxide, and organic acids they can produce; all listed have recognized pathogen inhibitory activities. The *Weissella* genus has also shown potential for treating atopic dermatitis and certain cancers. *W. cibaria*, *W. confusa*, and *W. paramesenteroides* are particularly of note because of their probiotic potential (fermentation of prebiotic fibers) and their ability to survive in the gastrointestinal tract. It is important to note that most of the *Weissella* strains with these health-promoting properties have been shown to be safe, due to the absence or the low occurrence of virulence or antibiotic-resistant genes. A large number of scientific studies continue to report on and to support the use of *Weissella* strains in the food and pharmaceutical industries. This review provides an overview of these studies and draws conclusions for future uses of this rich and previously unexplored genus.

Keywords: *Weissella*, Technological potential, Exopolysaccharide, Probiotic

1. Introduction

Advances in molecular biology can be credited with the discovery of the *Weissella* genus. Collins et al. [1] reported that strains with very similar phenotypic characteristics that had previously been included in the *Leuconostoc* genus exhibited different DNA profiles from other bacteria in the group. The authors used 16S rRNA sequences to study *Weissella*. They found that certain strains demonstrated specific features that were phylogenetically closer to *Leuconostoc paramesenteroides* than others. Collins et al. [1] recommended that *L. paramesenteroides*, a new species isolated from fermented sausages, and certain heterofermentative lactobacilli be classified as a new genus, *Weissella*.

The *Weissella* genus belongs to the Firmicutes phylum, Bacilli class, the Lactobacillales order, and the *Leuconostocaceae* family. *Weissella* bacteria are closely related to other lactic acid bacteria (LAB) genera. Fusco et al. [2] reported significant difficulty in separating the *Weissella* genus from other LAB, particularly from *Leuconostoc* species. *Weissella* species morphology may differ within the genus. The short rods with rounded to tapered ends or coccoid shapes that typify *Weissella* species are similar to *Leuconostoc*, *Oenococcus*, and streptococci [1]. Nevertheless, some *Weissella* species can present pleomorphisms under stress conditions. Cells can be present alone, in pairs, or even in small chains [1, 3]. Currently, 25 *Weissella* species have been determined: *W. viridescens*, [4], *W. paramesenteroides* [5], *W. confusa* [1], *W. kandleri* [6], *W. halotolerans* [7], *W. minor* [7], *W. hellenica* [1], *W. thailandensis* [8], *W. soli* [9], *W. cibaria* [10], *W. koreensis* [11], *W. ghanensis* [12], *W. beninenses* [13], *W. fabaria* [14], *W. ceti* [15], *W. fabalis* [16], *W. oryzae* [17], *W. diestrammenae* [18], *W. uvarum* [19], *W. cryptocerci* [20], *W. bombi* [21], *W. jogaejeotgali* [22], *W. kimchi* [23], *W. muntiacci* [24], and *W. sagaensis* [25]. A clear, well documented summary of *Weissella* taxonomy and ecology can be found in Fusco et al. [2]. Although *Weissella* remains a relatively new genus compared with other LAB, it has been the focus of research in the past 5 years and has attracted a strong interest from the food industry. Specific *Weissella* strains have been studied for their ability to produce (i) antimicrobial compounds like bacteriocins and hydrogen peroxide and (ii) exopolysaccharides (EPSs) and diacetyl. The latter are of major importance to dairy applications [26–29]. Different strains have been isolated from artisanally produced foods, which demonstrate the contributing role *Weissella* plays in the foods'

characteristic features. Other studies have highlighted *Weissella* strains' ability to survive in the gastrointestinal tract (GIT) and even grow in the gut. Both of these attributes suggest the use of *Weissella* species as beneficial applied probiotics [30–32].

The purpose of this review is to provide an overview of the *Weissella* genus and present the latest research and findings from the past 5 years. The paper presents the most recent techniques and approaches used in *Weissella* identification and characterization. It also covers the bacteria genus' potential health benefits and its main technological features which are of interest to the food industry.

2. *Weissella* Species Identification and Characterization

The past 5 years have seen the development of new methods for identifying *Weissella* species. They now make it possible to distinguish *Weissella* species from other closely related LAB. The polyphasic identification method and sequencing and analysis of the 16S and 23S rRNA genes are two new molecular-based identification methods that allow for rapid differentiation between *Lactiplantibacillus plantarum*, *Pediococcus pentosaceus*, and *Weissella confusa* [33]. Complete genome sequence analysis using bioinformatic analysis can be used to describe metabolic routes and to predict potential uses for the identified strains [34–38]. These methods are the foundation of modern microbial taxonomy, as are matrix-assisted laser desorption ionization time of flight method (MALDI-TOF) and database mass spectrometry [39, 40]. One recent study has shown that the colony morphology of *W. confusa* can change according to the adopted culture media and incubation conditions. *W. confusa*'s morphology can vary from rod-shaped coccobacilli—with characteristic, small, cream-colored colonies gathered in single, short or long chains—to a large, irregularly shaped, transparent morphology that lacks characteristic chain cell organization [41]. *Weissella* strains can also thrive in high-sugar niches and produce large amounts of dextran [41]. *Weissella* strains' behaviour in different substrates is key to differentiating them from other LAB.

Genomic and metatranscriptomic methods have been used to assess the metabolic and fermentative traits of *W. koreensis* during the kimchi fermentation process. This has led to the characterization of metabolic pathways for certain carbohydrates fermented by *W. koreensis*. These include D-glucose, D-mannose, D-lactose, L-malate, D-xylose, L-arabinose, D-ribose, N-acetyl-glucosamine, and

gluconate [42]. The determination of carbohydrate metabolic pathways makes it possible to predict how specific *Weissella* substrates will ferment, as well as pinpoint those which promote higher cell multiplication rates. It also allows for an estimation in type and quantity of products generated by the fermentation process. With this information, it is possible to predict the ecosystems where *Weissella* will survive and thus potential food and pharmaceutical applications for specific species.

Rizzello et al. [43] delved into *Weissella* genus metabolism by focusing on *W. cibaria* and *W. confusa* strains. The resulting data showed that *W. cibaria* and *W. confusa* were able to both produce phytase and use phytic acid, thus counteracting an anti-nutritional factor in legumes. The two strains were also able to reduce the raffinose oligosaccharide (RFO) concentrations of α -galactosides raffinose, verbascose, and stachyoseraphinose. RFO concentrations represent another anti-nutritional factor in vegetables, especially fava beans. When *W. cibaria* and *W. confusa* strains undergo the formation process, RFO hydrolysis enables galactose production. [43]. Legumes such as fava beans, peas, and lentils are a good source of vegetable proteins. Their anti-nutritional factors include saponins, condensed tannins, and protease inhibitors. α -Galactosides and phytic acid [44] have been shown to inhibit them. Fermentation and its subsequent reduction of these anti-nutritional factors would allow these foods to be used as new protein sources for both humans and animals. These new protein sources could in turn be used to develop new products in the food industry.

Many *Weissella* strains have been isolated from fermented products such as kimchi, pozol, jeotgal, and ogi [33, 39, 42, 45]. These strains have been characterized in terms of the carbon source they use and the compounds they can produce. While studying carbon source metabolism, Lopez-Hernandez et al. [45] observed that strains *W. cibaria*, *W. confusa*, and *W. paramesenteroides* were able to metabolize D-xylose, glucose, D-fructose, D-mannose, sucrose, and D-maltose. Some strains were also able to use ribose and esculin ferric citrate. Others were able to use galactose, metabolize cellobiose, and demonstrate β -glucosidase and β -galactosidase activity. The latter is a desirable characteristic in probiotic microorganisms, as β -glucosidase breaks down certain compounds and makes them easier for the body to absorb. Anthocyanins, which have antioxidant and anti-inflammatory properties, are among these compounds [46].

New identification methods have allowed for rapid and accurate identification of *Weissella* strains and reduced the risk of misidentification. Characterizing *Weissella* strains based on their metabolic pathways opens up the possibilities for their use in the medical and food production sectors.

3. Beneficial Properties of *Weissella*

Once the metabolic pathways of *Weissella* have been characterized, isolated strains can be studied for use in the pharmaceutical and food industries. *Weissella* isolates have beneficial, probiotic features and are able to produce EPS. These make them an important bacteria in both pharmaceutical and food production sectors. Abriouel et al. [47] described the technological, functional, and pathogenic potential of *Weissella* genus applications in the food industry, but their review only covered research carried out before 2015. Below, our review presents and discusses the primary benefits of *Weissella* based on these areas of study.

4. Health Benefits of *Weissella*

Few studies in the past 5 years have examined the effects of *Weissella* strains on human health [48–50]. Some research has evaluated their effects on mice [51–55], gerbils [56], and beagles [57]. Most studies have demonstrated how *Weissella*'s probiotic characteristics and antimicrobial properties may offer health benefits. Only two strains of *W. cibaria* have been studied for their probiotic potential for humans. *W. cibaria* JW15 was studied because of how it affects cytokine and immunoglobulin natural killer cell activity, and *W. cibaria* CMU has been the subject of research because of its oral epithelial cell adhesion and oral colonization; *W. cibaria* CMU's antibacterial activity against *Fusobacterium nucleatum* and *Streptococcus mutans* [48–50] represents another important species feature. A recent study found that immune functions were enhanced by an increase in natural killer cell activity for subjects who consumed probiotic *W. cibaria* capsules [50]. Moreover, oral ingestion of *W. cibaria* CMU can help to reduce halitosis and microbiota numbers in the gingival sulcus, thus improving oral health [48, 49].

Fonseca et al. [56] tested strains of *Bifidobacterium longum* and *W. paramesenteroides* to assess their probiotic potential and their impact on parasite load in gerbils. They observed the positive effect these two strains had in reducing the parasitic load of animals infected with *Giardia*. Strains of *W. cibaria* have been studied

as a way to reduce cancer treatment drug side effects in mice [55]. *W. cibaria* strains were shown to assist in the recovery of lymphocytes, hemoglobins, and platelet levels that drop when cyclosporamide (an anti-cancer agent) is administered. Cyclosporamide is widely used in the treatment of acute and chronic leukemia, lymphoma, and a number of autoimmune diseases [55]. *W. cibaria* strains also proved to be efficient in treating atopic dermatitis when administered orally to mice. *W. cibaria* improved the clinical symptoms of lesions, such as erythema/hemorrhage, edema/excoriation, erosion, scarring/dryness, and lichenification [54]. In yet another study, *W. cibaria* was shown to reduce the destruction of periodontal tissue in mice and also protected against alveolar bone destruction in mice. *W. cibaria* co-aggregates with periodontal bacteria to produce hydrogen peroxide and bacteriocins that inhibit the production of proinflammatory cytokines [58]. Although clinical studies using mice are an effective research technique, these studies cannot reliably predict the outcome of human studies [59]. Further study is needed to determine the health benefits of *Weissella* strains for humans.

Although there have been few human or animal studies using *Weissella* strains in vitro research indicates a potential for future use as antimicrobial agents (Table 1) and probiotics (Table 2). Further study on animals and/ or humans is necessary to prove their efficiency as a functional species. The *W. confusa* DD_A7 strain has been shown to be a natural alternative prophylactic agent against multidrug-resistant (MDR) ESBL (extended spectrum β -lactamase) positive *E. coli* bacteria [27, 29]. *W. confusa* has also been shown to be effective destroying *Listeria monocytogenes* (ATCC 7644), *Staphylococcus aureus* (ATCC 12600), *Salmonella* Typhimurium (ATCC 43174), *Bacillus cereus* (ATCC 13061), and *E. coli* O157: H7 (ATCC 43889) pathogens. Several studies have shown that the supernatant from of *W. confusa* culture weakened the membrane of the pathogen *E. coli* O157: H7 (ATCC 43889) and damaged the integrity of its DNA [27, 29]. Another study which paired *W. viridescens* with high pressure processing showed strong results when it came to protecting ready-to-eat salads from *L. monocytogenes*. The research showed that high pressure applied to the weakened pathogen membranes which then facilitated the antimicrobial activity produced by the *W. viridescens* strain [60].

Table 1. *Weissella* species with antimicrobial effects and the respective pathogens that each one inhibits.

Species	Pathogen	Reference
<i>W. viridescens</i>	<i>L. monocytogenes</i>	Stratakos et al. [60]
<i>W. paramesenteroides</i>	<i>Giardia lamblia</i>	Fonseca et al. [56]
	<i>E. coli</i>	Pabari et al. [62]
	<i>S. aureus</i>	
<i>W. cibaria</i>	<i>Porphyromonas gingivalis</i>	Do et al. [57]
	<i>Prevotella intermedia</i>	
	<i>Fusobacterium nucleatum</i>	
	<i>S. enterica</i> subsp. <i>enterica</i>	Tenea et al. [108]
<i>W. confusa</i>	<i>Escherichia coli</i>	
	<i>S. enterica</i> Typhi	Pelyuntha et al. [109]
	<i>E. coli</i>	Dey et al. [29]
	<i>Candida albicans</i>	Rosca et al. [80]
	<i>S. enterica</i>	Lakra et al. [61]
	<i>S. enterica</i> Typhi	
	<i>L. monocytogenes</i>	
	<i>S. aureus</i>	
	<i>L. monocytogenes</i>	Dey et al., [27]
	<i>S. aureus</i>	
<i>S. enterica</i> Typhimurium		
<i>B. cereus</i>		
<i>E. coli</i>		

Table 2. *In vitro* and *in vivo* evaluation of *Weissella* species as a probiotic candidate.

Species	Probiotic property	References
<i>Weissella</i> spp	Cholesterol-reducing potential	Anandharaj et al. [30]
<i>W. cibaria</i>	Enhance in immune functions	Lee et al. [50]
	Reduce the side effects of drugs	Park and Lee [55]
	Improve atopic dermatitis	Lim et al. [54]
	Protective effects against alveolar bone destruction	Kim et al. [58]
	Short-chain fatty acids production and adhesion to HT-29 cells.	Silva et al. [64]
<i>W. confusa</i>	inhibition of $-\alpha$ -amylase activity	Xia et al. [32]

	Thermostability, cholesterol removal, β -galactosidase production and proteolytic activity	Sharma et al. [63]
<i>W. confusa</i> and <i>W. cibaria</i>	reduction in cholesterol, DPPH free radical and inhibition of linoleic acid peroxidation.	Xia et al. [32] Lakra et al. [61]
<i>W. paramesenteroides</i>	Formation of short-chain fatty acids and antimicrobial compounds.	Pabari et al. [62]

Recent in vitro studies have shown that *W. cibaria*, *W. koreensis*, *W. confusa*, and *W. paramesenteroides* strains are strong candidates for probiotic applications [30, 32,61–64]. The survival of these strains in the GIT and their tolerance levels when in contact with bile and acids are important factors for intestinal colonization and pathogen protection. *W. paramesenteroides* has also been shown to form a biofilm which facilitates mucin adherence. It uses fructooligosaccharides (FOS) and galactooligosaccharides (GOS) as prebiotic fibers to the form of compounds that can survive the gut microbiome. *W. paramesenteroides* has also demonstrated antimicrobial activity against *E. coli* and *S. aureus* [62]. In another study, Anandharaj et al. [30] demonstrated that *Weissella* species may help reduce cholesterol due to its ability to use cholesterol from de Man, Rogosa and Sharpe (MRS) agar with added Ovgall and water-soluble cholesterol. Cholesterol reduction is common in probiotic microorganisms isolated from the intestine [65–67]. Interestingly, Kim et al. [58] found that using the *W. cibaria* D30 probiotic strain in fermented *Inula britannica* created a product with improved anti-inflammatory capacity. *W. confusa* and *W. cibaria*—isolated from Plaa Som Fug—have also been shown to produce folate. Folate, or Vitamin B9, plays an important role in DNA replication, repair, and methylation. It can also act as an antioxidant [68]. Folate is important for immune and nervous system functions because it plays an integral part in the synthesis of neurotransmitters in the central nervous system [69]. This is especially the case during the early stages of uterine growth and development.

Extensive research has been carried out on *Weissella* genus antimicrobial activity and its potential use of strains as probiotics. However, other beneficial compounds produced by *Weissella*, including vitamins, have only been touched on in research thus far.

5. EPS Production by *Weissella*

EPSs are extracellular macromolecules that have potential applications in the food, medical, and pharmaceutical industries. It has been shown that *Weissella* strains can produce these molecules, and EPS production has been the focus of several *Weissella* studies (Table 3). EPS production has been primarily documented for *W. confusa* and *W. cibaria* strains, with dextran as the dominant EPS produced by the strains. Structurally, dextran molecules are highly linear. They are formed by α -1,6-linked glucose residues with branches linked to α -1,2, α -1,3, or α -1,4. The α -1,2 bonds are representative of prebiotic properties because they are resistant to adverse GIT conditions and can help to modulate the intestinal microbiota [41, 70–72]. EPS produced by LAB are generally recognized as safe (GRAS), which means they can be used in foods, prebiotics, and medical applications.

Table 3. *Weissella* species and its EPS products.

Species	EPS produced	References
<i>W. confusa</i>	EPS	Benhoua et al. [28]
	Dextran	Heperkan et al. [41]
		Rosca et al. [80]
		Tang et al. [86]
		Devi et al. [73]
	Galactan	Kavitake et al. [81]
		Lakra et al. [110]
Lactose- and cellobiose-derived branched trisaccharides Isomelezitose	Shi et al. [78]	
Homopolysaccharide with glucose monomers	Adesulu-Dahunsi et al. [33]	
Wild and mutant <i>W. confusa</i>	EPS with eight sugar moieties	Adebayo-Tayo et al. [51]
<i>W. cibaria</i>	Dextran	Xu et al. [87]
		Zannini et al. [84]
		Baruah et al. [77,82]
		Kanimozhi et al. [72]
		Yu et al. [75]
	Linear dextran	Ye et al. [74]
Dextran	Hu, Gänzle [71]	

Oligosaccharides	
Heteropolysaccharides	Zhu et al. [76]
Isomalto-oligosaccharide	Baruah et al. [77,82]
	Rolim et al. [85]
EPS mainly composed of glucose and rhamnose sugar units	Adesulu-Dahunsi et al. [26]

Of the LAB *Weissella* species, *W. cibaria* and *W. confusa* are the most important EPS producers, which is why research has focused on them and their production, purification, and characterization [41, 71–76]. Some studies have proposed ways to optimize EPS production and to test different parameters, such as the effects of different temperatures and substrates. Hu and Ganzel [71] showed that *W. cibaria* strains grew faster at 30 °C, but oligosaccharide production was higher at temperatures at or below 20 °C. The greatest dextran production occurred at 15 °C.

The physicochemical properties of bacteria-produced EPS are an important factor for functionality and use. Devi et al. [73] carried out physicochemical characterizations of galactan EPS produced by a *W. confusa* strain. They showed that the EPS displayed a high oil absorption capacity, strong emulsifying activity, and an emulsion kinetic stability of up to 15 days. These results supported the theory that galactan EPS may be a good candidate as an emulsifier in forming long-term colloidal systems in food, pharmaceutical, and cosmetic products. Other studies have shown that several *Weissella* strains use a dextransucrase enzyme to produce dextran. This enzyme has been the focus of isolation and purification studies for the production of EPS [77, 78] and caffeic acids [79]. Dextransucrase produced by a *W. confusa* strain has been shown to synthesize a rare oligosaccharide called isomelezitose. This oligosaccharide has a potent nutraceutical effect due to its ability to promote bifidobacteria growth in the colon [78].

The EPS produced by LAB have antifungal [80], antiinflammatory, and immunomodulatory properties [51] that may be of great interest to the medical sector. EPS can also act as antioxidants [26, 28, 51, 76] and help probiotics survive passage through the GIT [81]. The linear EPS galactan produced by the *W. confusa* strain shows strong potential for encapsulation technology that is used to deliver bioactive compounds, probiotics, and drugs. The EPS produced by a *W. confusa* strain was shown to inhibit the multiplication of *Candida albicans* and was able to destroy up to

70% of the biofilm formed by this pathogen [80]. This suggests that it could be a promising candidate for use in antifungal treatments. Another recent study showed that *W. confusa* can produce mannan, an EPS made up of only monomeric units of mannose. This EPS displayed antibiofilm activity against pathogenic bacteria such as *S. aureus*, *L. monocytogenes*, *S. enterica*, and *S. typhi*. It also demonstrated potential as an antioxidant compound [61].

EPS may also have prebiotic properties. This is the case for the EPS produced by the *W. cibaria* strain isolated from pomelos (*Citrus maximum*) grown in India [82]. To be categorized as a prebiotic compound, the EPS carbohydrates must demonstrate higher metabolization levels from probiotic bacteria than from enteric bacteria present in the intestine. Dextran produced by *W. cibaria* was shown to stimulate the growth of *L. plantarum*, *L. acidophilus*, *B. animalis*, *B. bifidum*, and *B. infantis* more than *E. coli* and *E. aerogenes* [82]. This activity confirmed its classification as a prebiotic compound.

In the food industry, EPS can be used to improve the rheological characteristics and texture of fermented products. EPS can also play a role as an emulsifier and stabilizer [28, 74, 81]. EPS-producing LABs can be used as starter or secondary cultures in the production of fermented products such as yogurt [83]. The use of these cultures leads to the production of EPS in situ, where they act as natural thickeners and can eliminate the need for artificial additives [76]. *Weissella* strains have been used to ferment quinoa-based yogurt [84] and fruit juices [85, 86]. The EPS produced can improve the sensory characteristics of such products. One example: Orange juice concentrate was used to produce oligosaccharides. Then, after a fermentation process involving the sucrose present in the juice itself, a drink was obtained with reduced a sugar content and specific acidity and sweetness levels [85, 86].

EPS are also of interest to the frozen dough industry, as they can modify the texture and improve the end product characteristics of frozen and thawed doughs. Studies have shown that dextrans produced by different microorganisms have different functionalities. For instance, dextran produced by the *W. cibaria* strain demonstrated a greater ability to reinforce gelling properties compared with dextran produced by *L. pseudomesenteroides* when the strains were mixed with fava bean (*Vicia faba* L.) protein isolate [87, 88]. *Weissella* EPS has also been shown to improve the characteristics of wheat gluten during dough freezing and thawing. In general, the freezing-thawing process leads to an increase in water loss, a decrease in gluten

viscoelasticity, and changes in water mobility and distribution. During freezing and thawing, the continuous structure of gluten is destroyed by mechanical damage caused when ice recrystallizes. An addition of *W. confuse*-produced EPS to doughs helped the gluten to maintain its qualities and structural integrity during the freezing-thawing process. This makes *Weissella* EPS promising cryoprotectant candidates for the frozen dough industry [86].

6. *Weissella* as an Emerging Pathogen

Weissella isolates with potential health benefits must also be assessed for any adverse effects. Testing is standard procedure for isolates to be used as food supplements and starter cultures to determine if they harbor virulent or antibiotic-resistant genes, even when the negative effects are not active or apparent. [89]. If or when any noxious genetic properties are present in the isolate genome, they represent a health risk because they may be transferred to other bacteria present in the food matrix or gut [90–92].

Although many LAB cultures found in foods and are considered safe, the biosafety of new cultures must be established before they can be used in food production [93]. Most of the studies carried out to determine *Weissella* strains' probiotic potential also evaluate the strains' health risks. These assessments include tests for hemolytic capacity and resistance to antibiotics, PCR detection of virulence genes, and biogenic amine production [94–96]. Kang et al. [97] assessed the safety of *W. cibaria* CMU and CMS1 when used for oral care probiotics. Their results confirmed the strain is resistant to vancomycin and kanamycin (antibiotics) and that the resistance is intrinsic to the genus. They also demonstrated that the antibiotic resistance of *W. cibaria* CMU was not transferred to recipient strains. This was confirmed by the fact that no homology for a conjugative transposon integron-specific gene was present, i.e., there was no trace of antibiotic resistance genes in either chromosomal or plasmid DNA samples. They found that the strain had negative results for β -hemolysis, mucin degradation, and platelet aggregation tests. The strain also tested negative for various toxic secondary metabolites [97].

Until now, *Weissella* strains have only been associated with “weissellosis,” a disease that affects rainbow trout (*Oncorhynchus mykiss*). The disease leads to the septicemia the fish and can cause important economic losses on fish farms. “Weissellosis” is caused by a specific species, the *W. ceti* [98–100]. Because certain

Weissella strains are resistant to antibiotics such as vancomycin, teicoplanin, ceftazidime, and sulfamethoxazole, human infections remain a possibility. These would primarily affect immunocompromised patients [3, 101]. It should be noted, however, that even generally recognized probiotic strains may cause health complications. *Lacticaseibacillus rhamnosus* GG and *L. helveticus* are two examples of this [102, 103].

Even though there are few cases of this genus causing disease in humans or animals, recent studies have evaluated the safety of *Weissella* use in foods. Any strain considered for use in fermented foods must be approved by government regulatory commissions. Although Brazil, the USA, and European Union countries do not have specific legislation for the production of new types of yeast, they do have recommendations. In the USA, for example, some cultures are classified as safe and suitable for human consumption while others are classified as GRAS and must apply for approval from the FDA (Food and Drug Administration) [104].

In 2011, some *Weissella* species, including *W. confusa*, were added to the International Dairy Federation (IDF) inventory, a list of microorganisms allowed in food fermentation processes. The German Committee for Biological Agents has also placed *W. confusa* in Risk Group 1 for microorganism that are unlikely to cause human diseases [105]. *W. confusa*'s important technological potential as a source of probiotics and other health benefits have led to two studies that address the microorganism's safety [105, 106]. Cupi, Elvig-Jorgensen [106] carried out a series of toxicity tests which included genotoxicity, skin irritation, eye irritation, and sub-chronic oral toxicity in rats. They found that the *W. confusa* strain did not show any toxic effects and therefore can be used as a safe, direct-feed microbial product for animals. Kang et al. [97] showed that *W. cibaria* strains could be granted GRAS status in the future because: (i) they do not have antibiotic-resistant genes, (ii) they lack antibiotic resistance transferability, (iii) their genomic sequences do not include virulent genes related to pathogenic bacteria, and (iv) they tested negative for most virulent factors (β -hemolysis, mucin degradation, etc.) and for toxic metabolic production (ammonia production, β -glucuronidase activity, etc.).

Although many studies have demonstrated the safety of *Weissella*, the genus has not yet been included in the Qualified Presumption of Safety (QPS) list published by the European Food Safety Authority (EFSA) [107]. Research on *W. cibaria* and *W. confusa* has been carried out to this end [96]. To be granted the QPS status, a

microorganism must meet the following criteria: (i) its taxonomic identity must be well defined, (ii) the available body of knowledge must be enough to establish its safety, (iii) the lack of pathogenic properties must be established and substantiated, and (iv) its intended use must be clearly described [107].

7. Conclusion

Over the past 5 years, the *Weissella* genus has been the focus of many studies due to its strong potential as a probiotic that could be used in both the food and pharmaceutical industries. *Weissella* strains' probiotic potential is attributed to its remarkable ability to survive passage through the GIT, produce antimicrobial substances for a variety of pathogens, and promote the formation of compounds that stimulate the gut microbiome. The ability of some *Weissella* strains to produce large amounts of EPS represents another major attribute, as EPS have prebiotic properties. *Weissella* EPS production also demonstrates the strains' potential as a natural thickener for foods, among other possibilities.

Moreover, recent studies have demonstrated that *Weissella* strains do not have antibiotic-resistant genes and they generally test negative for virulent factors. Thus, there is a low probability that the genus could cause foodborne diseases or carry on virulent genes that may be transferred to other bacteria pathogens.

8. References

1. Collins MD, Samelis J, Metaxopoulos J, Wallbanks S (1993) Taxonomic studies on some leuconostoc-like organisms from fermented sausages: description of a new genus *Weissella* for the *Leuconostoc paramesenteroides* group of species. *J Appl Bacteriol* 75:595–603. <https://doi.org/10.1111/j.1365-2672.1993.tb01600.x>
2. Fusco V, Quero GM, Cho GS et al (2015) The genus *Weissella*: taxonomy, ecology and biotechnological potential. *Front Microbiol* 6:22. <https://doi.org/10.3389/fmicb.2015.00155>
3. Huys G, Leisner J, Bjorkroth J (2012) The lesser LAB gods: *Pediococcus*, *Leuconostoc*, *Weissella*, *Carnobacterium*, and affiliated genera. In: Lahtinen S, Ouwehand AC, Salminen S, Von WA (eds) *Lactic Acid Bacteria, Microbiological and Functional Aspects*, fourth. Taylor & Francis Group, New York, pp 94–112
4. Niven CF, Evans JB (1957) *Lactobacillus viridescens* nov. spec., a heterofermentative species that produces a green discoloration of cured meat pigments. *J Bacteriol* 73:758–759
5. Garvie EI (1967) The growth factor and amino acid requirements of species of the genus *Leuconostoc*, including *Leuconostoc paramesenteroides* (sp. nov.) and *Leuconostoc oenos*. *J Gen Microbiol* 48:439–447. <https://doi.org/10.1099/00221287-48-3-439>
6. Holzapfel WH, Van WEP (1982) *Lactobacillus kandleri* sp. nov., a new species of the subgenus *Betabacterium*, with glycine in the peptidoglycan. *Zentralblatt für Bakteriologie Mikrobiologie*

und Hyg I Abt Orig C Allg Angew und ökologische Mikrobiol 3:495–502. [https://doi.org/10.1016/S0721-9571\(82\)80007-0](https://doi.org/10.1016/S0721-9571(82)80007-0)

7. Kandler O, Schillinger U, Weiss N (1983) *Lactobacillus halotolerans* sp. nov., nom. rev. and *Lactobacillus minor* sp. nov., nom. rev. Syst Appl Microbiol 4:280–285. [https://doi.org/10.1016/S0723-2020\(83\)80056-3](https://doi.org/10.1016/S0723-2020(83)80056-3)

8. Tanasupawat S, Shida O, Okada S (2016) *Weissella thailandensis* sp. nov., isolated from fermented fish in Thailand. 023838:1479–1485

9. Magnusson J, Jonsson H, Schnurer J, Roos S (2002) *Weissella soli* sp. nov., a lactic acid bacterium isolated from soil. Int J Syst Evol Microbiol 52:831–834. <https://doi.org/10.1099/ijs.0.02015-0>

10. Bjorkroth KJ, Schillinger U, Geisen R et al (2002) Taxonomic study of *Weissella confusa* and description of *Weissella cibaria* sp. nov., detected in food and clinical samples. Int J Syst Evol Microbiol 52:141–148. <https://doi.org/10.1099/00207713-52-1-141>

11. Lee J-S, Lee KC, Ahn J-S et al (2002) *Weissella koreensis* sp. nov., isolated from kimchi. Int J Syst Evol Microbiol 52:1257–1261

12. De Bruyne K, Camu N, Lefebvre K et al (2008) *Weissella ghanensis* sp. nov., isolated from a Ghanaian cocoa fermentation. Int J Syst Evol Microbiol 58:2721–2725. <https://doi.org/10.1099/ijs.0.65853-0>

13. Padonou SW, Schillinger U, Nielsen DS et al (2010) *Weissella beninensis* sp. nov., a motile lactic acid bacterium from submerged cassava fermentations, and emended description of the genus *Weissella*. Int J Syst Evol Microbiol 60:2193–2198. <https://doi.org/10.1099/ijs.0.014332-0>

14. De Bruyne K, Camu N, De Vuyst L, Vandamme P (2010) *Weissella fabaria* sp. nov., from a Ghanaian cocoa fermentation. Int J Syst Evol Microbiol 60:1999–2005. <https://doi.org/10.1099/ijs.0.019323-0>

15. Vela AI, Fernandez A, de Quiros YB et al (2011) *Weissella ceti* sp. nov., isolated from beaked whales (*Mesoplodon bidens*). Int J Syst Evol Microbiol 61:2758–2762. <https://doi.org/10.1099/ijs.0.028522-0>

16. Snauwaert I, Papalexandratou Z, De Vuyst L, Vandamme P (2013) Characterization of strains of *Weissella fabalis* sp. nov. and *Fructobacillus tropaeoli* from spontaneous cocoa bean fermentations. Int J Syst Evol Microbiol 63:1709–1716. <https://doi.org/10.1099/ijs.0.040311-0>

17. Tohno M, Kitahara M, Inoue H et al (2013) *Weissella oryzae* sp. nov., isolated from fermented rice grains. Int J Syst Evol Microbiol 63:1417–1420. <https://doi.org/10.1099/ijs.0.043612-0>

18. Oh SJ, Shin N-R, Hyun D-W et al (2013) *Weissella diestrammenae* sp. nov., isolated from the gut of a camel cricket (*Diestrammena coreana*). Int J Syst Evol Microbiol 63:2951–2956. <https://doi.org/10.1099/ijs.0.047548-0>

19. Nisiotou A, Dourou D, Filippousi M-E et al (2014) *Weissella uvarum* sp. nov., isolated from wine grapes. Int J Syst Evol Microbiol 64:3885–3890. <https://doi.org/10.1099/ijs.0.066209-0>

20. Heo J, Hamada M, Cho H et al (2019) *Weissella cryptocerci* sp. nov., isolated from gut of the insect *Cryptocercus kyebangensis*. Int J Syst Evol Microbiol 69:2801–2806. <https://doi.org/10.1099/ijs.0.003564>

21. Praet J, Meeus I, Cnockaert M et al (2015) Novel lactic acid bacteria isolated from the bumble bee gut: *Convivina* intestine gen. nov., sp. nov., *Lactobacillus bombicola* sp. nov., and

- Weissella bombi* sp. nov. Antonie van Leeuwenhoek, J Microbiol 107:1337–1349. <https://doi.org/10.1007/s10482-015-0429-z>
22. Lee S, Ku H, Ahn M et al (2015) *Weissella jogaejeotgali* sp. nov., isolated from jogae jeotgal, a traditional Korean fermented seafood. Int J Syst Evol Microbiol 65:4674–4681. <https://doi.org/10.1099/ijsem.0.000631>
23. Choi H, Cheigh C, Kim S et al (2002) *Weissella kimchii* sp. nov., a novel lactic acid bacterium from kimchi. Int J Syst Evol Microbiol 52:507–511
24. Lin S-T, Wang L-T, Wu Y-C et al (2020) *Weissella muntiaci* sp. nov., isolated from faeces of Formosan barking deer (*Muntiacus reevesi*). Int J Syst Evol Microbiol 70:1578–1584
25. Li YQ, Tian WL, Gu CT (2020) *Weissella sagaensis* sp. nov., isolated from traditional Chinese yogurt. Int J Syst Evol Microbiol 70:2485–2492
26. Adesulu-Dahunsi AT, Sanni AI, Jeyaram K (2018) Production, characterization and in vitro antioxidant activities of exopolysaccharide from *Weissella cibaria* GA44. LWT - Food Sci Technol 87:432–442. <https://doi.org/10.1016/j.lwt.2017.09.013>
27. Dey DK, Koo BG, Sharma C, Kang SC (2019) Characterization of *Weissella confusa* DD_A7 isolated from kimchi. LWT - Food Sci Technol 111:663–672. <https://doi.org/10.1016/j.lwt.2019.05.089>
28. Benhouna IS, Heumann A, Rieu A et al (2019) Exopolysaccharide produced by *Weissella confusa*: chemical characterisation, rheology and bioactivity. Int Dairy J 90:88–94. <https://doi.org/10.1016/j.idairj.2018.11.006>
29. Dey DK, Khan I, Kang SC (2019) Anti-bacterial susceptibility profiling of *Weissella confusa* DD_A7 against the multidrug-resistant ESBL-positive *E. coli*. Microb Pathogenesis 128:119–130. <https://doi.org/10.1016/j.micpa.2018.12.048>
30. Anandharaj M, Sivasankari B, Santhanakaruppu R et al (2015) Determining the probiotic potential of cholesterol-reducing *Lactobacillus* and *Weissella* strains isolated from gherkins (fermented cucumber) and south Indian fermented koozh. Res Microbiol 166:428–439. <https://doi.org/10.1016/j.resmi.2015.03.002>
31. Yu H-S, Lee N-K, Choi A-J et al (2019) Antagonistic and antioxidant effect of probiotic *Weissella cibaria* JW15. Food Sci Biotechnol 28:851–855. <https://doi.org/10.1007/s10068-018-0519-6>
32. Xia Y, Qin S, Shen Y (2019) Probiotic potential of *Weissella* strains isolated from horse feces. Microb Pathog 132:117–123. <https://doi.org/10.1016/j.micpa.2019.04.032>
33. Adesulu-Dahunsi AT, Sanni AI, Jeyaram K (2017) Rapid differentiation among *Lactobacillus*, *Pediococcus* and *Weissella* species from some Nigerian indigenous fermented foods. LWT - Food Sci Technol 77:39–44. <https://doi.org/10.1016/j.lwt.2016.11.007>
34. Shin J, Oh G, Bang M et al (2020) Complete genome sequence of *Weissella cibaria* strain BM2, isolated from Korean kimchi. Microbiol Resour Announc 9:10–11
35. Manberger A, Verbrugghe P, Gudmundsdottir EE et al (2020) Taxogenomic assessment and genomic characterisation of *Weissella cibaria* strain 92 able to metabolise oligosaccharides derived from dietary fibres. Sci Rep 10:1–14. <https://doi.org/10.1038/s41598-020-62610-x>
36. Falasconi I, Fontana A, Patrone V et al (2020) Genome-assisted characterization of *Lactobacillus fermentum*, *Weissella cibaria*, and *Weissella confusa* strains isolated from sorghum as starters for sourdough fermentation. Microorganisms 8:1–16. <https://doi.org/10.3390/microorganisms8091388>

37. Patrone V, Al-Surrayai T, Romaniello F et al (2020) Integrated phenotypic-genotypic analysis of candidate probiotic *Weissella cibaria* strains isolated from dairy cows in Kuwait. *Probiotics Antimicrob Proteins*. <https://doi.org/10.1007/s12602-020-09715-x>
38. Mun SY, Chang HC (2020) Characterization of *Weissella koreensis* sk isolated from kimchi fermented at low temperature (Around 0° C) based on complete genome sequence and corresponding phenotype. *Microorganisms* 8:1–17. <https://doi.org/10.3390/microorganisms8081147>
39. Kim E, Cho Y, Lee Y et al (2017) A proteomic approach for rapid identification of *Weissella* species isolated from Korean fermented foods on MALDI-TOF MS supplemented with an in-house database. *Int J Food Microbiol* 243:9–15. <https://doi.org/10.1016/j.ijfoodmicro.2016.11.027>
40. Nacef M, Chevalier M, Chollet S et al (2017) MALDI-TOF mass spectrometry for the identification of lactic acid bacteria isolated from a French cheese : the Maroilles. *Int J Food Microbiol* 247:2–8. <https://doi.org/10.1016/j.ijfoodmicro.2016.07.005>
41. Heperkan ZD, Bolluk M, Bulbul S (2020) Structural analysis and properties of dextran produced by *Weissella confusa* and the effect of different cereals on its rheological characteristics. *Int J Biol Macromol* 143:305–313. <https://doi.org/10.1016/j.ijbiomac.2019.12.036>
42. Jeong SE, Chun BH, Kim KH et al (2018) Genomic and metatranscriptomic analyses of *Weissella koreensis* reveal its metabolic and fermentative features during kimchi fermentation. *Food Microbiol* 76:1–10. <https://doi.org/10.1016/j.fm.2018.04.003>
43. Rizzello CG, Coda R, Wang Y et al (2019) Characterization of indigenous *Pediococcus pentosaceus*, *Leuconostoc kimchii*, *Weissella cibaria* and *Weissella confusa* for faba bean bioprocessing. *Int J Food Microbiol* 302:24–34. <https://doi.org/10.1016/j.ijfoodmicro.2018.08.014>
44. Alcazar-Valle M, Lugo-Cervantes E, Mojica L et al (2020) Bioactive compounds, antioxidant activity, and antinutritional content of legumes: a comparison between four *Phaseolus* Species. *Molecules* 25:3528
45. Lopez-Hernandez M, Rodriguez-Alegria ME, Lopez-Munguia A, Wachter C (2018) Evaluation of xylan as carbon source for *Weissella* spp., a predominant strain in pozol fermentation. *LWT - Food Sci Technol* 89:192–197. <https://doi.org/10.1016/j.lwt.2017.10.030>
46. Cheng JR, Liu XM, Chen ZY et al (2016) Mulberry anthocyanin biotransformation by intestinal probiotics. *Food Chem* 213:721–727. <https://doi.org/10.1016/j.foodchem.2016.07.032>
47. Abriouel H, Lerma LL, del Casado Munoz MC et al (2015) The controversial nature of the *Weissella* genus: Technological and functional aspects versus whole genome analysis-based pathogenic potential for their application in food and health. *Front Microbiol* 6:1–14. <https://doi.org/10.3389/fmicb.2015.01197>
48. Kang MS, Lee DS, Lee SA et al (2020) Effects of probiotic bacterium *Weissella cibaria* CMU on periodontal health and microbiota: a randomised, double-blind, placebo-controlled trial. *BMC Oral Health* 20:1–12. <https://doi.org/10.1186/s12903-020-01231-2>
49. Lee DS, Lee SA, Kim M et al (2020) Reduction of halitosis by a tablet containing *Weissella cibaria* CMU: a randomized, double blind, placebo-controlled study. *J Med Food* 23:649–657. <https://doi.org/10.1089/jmf.2019.4603>
50. Lee YJ, Lee A, Yoo HJ et al (2018) Supplementation with the probiotic strain *Weissella cibaria* JW15 enhances natural killer cell activity in nondiabetic subjects. *J Funct Foods* 48:153–158. <https://doi.org/10.1016/j.jff.2018.07.009>

51. Adebayo-Tayo B, Ishola R, Oyewunmi T (2018) Characterization, antioxidant and immunomodulatory potential on exopolysaccharide produced by wild type and mutant *Weissella confusa* strains. *Biotechnol Reports* 19:e00271. <https://doi.org/10.1016/j.btre.2018.e00271>
52. Elshaghabee FMF, Ghadimi D, Habermann D et al (2020) Effect of oral administration of *Weissella confusa* on fecal and plasma ethanol concentrations, lipids and glucose metabolism in Wistar rats fed high fructose and fat diet. *Hepatic Med Evid Res* 12:93–106
53. Kim HY, Bae WY, Yu HS et al (2020) *Inula britannica* fermented with probiotic *Weissella cibaria* D30 exhibited anti-inflammatory effect and increased viability in RAW 264.7 cells. *Food Sci Biotechnol* 29:569–578. <https://doi.org/10.1007/s10068-019-00690-w>
54. Lim SK, Kwon M-S, Lee J et al (2017) *Weissella cibaria* WIKIM28 ameliorates atopic dermatitis-like skin lesions by inducing tolerogenic dendritic cells and regulatory T cells in BALB/c mice. *Sci Rep* 7:1–9. <https://doi.org/10.1038/srep40040>
55. Park HE, Lee WK (2018) Immune enhancing effects of *Weissella cibaria* JW15 on BALB/c mice immunosuppressed by cyclophosphamide. *J Funct Foods* 49:518–525. <https://doi.org/10.1016/j.jff.2018.09.003>
56. Fonseca JF, Alvim LB, Nunes C et al (2019) Probiotic effect of *Bifidobacterium longum* 51A and *Weissella paramesenteroides* WpK4 on gerbils infected with *Giardia lamblia*. *J Appl Microbiol* 127:1184–1191. <https://doi.org/10.1111/jam.14338>
57. Do K, Park H, Kang M et al (2019) Effects of *Weissella cibaria* CMU on halitosis and calculus, plaque, and gingivitis indices in beagles. *J Vet Dent* 36:135–142. <https://doi.org/10.1177/0898756419872562>
58. Kim J, Jung BH, Lee JH et al (2020) Effect of *Weissella cibaria* on the reduction of periodontal tissue destruction in mice. *J Periodontol* 1–8. <https://doi.org/10.1002/jper.19-0288>
59. Perlman RL (2016) Mouse models of human disease: an evolutionary perspective. *Evol Med Public Heal* 2016:170–176. <https://doi.org/10.1093/emph/eow014>
60. Stratakos AC, Linton M, Tessema GT et al (2016) Effect of high pressure processing in combination with *Weissella viridescens* as a protective culture against *Listeria monocytogenes* in ready to-eat salads of different pH. *Food Control* 61:6–12. <https://doi.org/10.1016/j.foodcont.2015.09.020>
61. Lakra AK, Domdi L, Hanjon G et al (2020) Some probiotic potential of *Weissella confusa* MD1 and *Weissella cibaria* MD2 isolated from fermented batter. *LWT - Food Sci Technol* 125:109261. <https://doi.org/10.1016/j.lwt.2020.109261>
62. Pabari K, Pithva S, Kothari C et al (2020) Evaluation of probiotic properties and prebiotic utilization potential of *Weissella paramesenteroides* isolated from fruits. *Probiotics Antimicrob Proteins* 12:1126–1138. <https://doi.org/10.1007/s12662-019-09630-w>
63. Sharma S, Kandasamy S, Kavitha D, Shetty PH (2018) Probiotic characterization and antioxidant properties of *Weissella confusa* KR780676, isolated from an Indian fermented food. *LWT - Food Sci Technol* 97:53–60. <https://doi.org/10.1016/j.lwt.2018.06.033>
64. Silva MS, Ramos CL, Gonzalez-Avila M et al (2017) Probiotic properties of *Weissella cibaria* and *Leuconostoc citreum* isolated from tejuino – a typical Mexican beverage. *LWT - Food Sci Technol* 86:227–232. <https://doi.org/10.1016/j.lwt.2017.08.009>
65. Ait Seddik H, Bendali F, Cudennec B, Drider D (2017) Antipathogenic and probiotic attributes of *Lactobacillus salivarius* and *Lactobacillus plantarum* strains isolated from feces of Algerian infants and adults. *Res Microbiol* 168:244–254. <https://doi.org/10.1016/j.resmic.2016.12.003>

66. Cavalcante RGS, de Albuquerque TMR, de Luna Freire MO et al (2019) The probiotic *Lactobacillus fermentum* 296 attenuates cardiometabolic disorders in high fat diet-treated rats. *Nutr Metab Cardiovasc Dis* 29:1408–1417. <https://doi.org/10.1016/j.numec d.2019.08.003>
67. Liang X, Lv Y, Zhang Z et al (2020) Study on intestinal survival and cholesterol metabolism of probiotics. *LWT - Food Sci Technol* 124:109–132. <https://doi.org/10.1016/j.lwt.2020.109132>
68. Deatraksa J, Sunthornthummas S, Rangsiruji A et al (2018) Isolation of folate-producing *Weissella* spp. from Thai fermented fish (Plaa Som Fug). *LWT - Food Sci Technol* 89:388–391. <https://doi.org/10.1016/j.lwt.2017.11.016>
69. Hsieh YC, Chou LS, Lin CH et al (2019) Serum folate levels in bipolar disorder: a systematic review and meta-analysis. *BMC Psychiatry* 19:1–9. <https://doi.org/10.1186/s12888-019-2269-2>
70. Bounaix MS, Gabriel V, Morel S et al (2009) Biodiversity of exopolysaccharides produced from sucrose by sourdough lactic acid bacteria. *J Agric Food Chem* 57:10889–10897. <https://doi.org/10.1021/jf902068t>
71. Hu Y, Ganzle MG (2018) Effect of temperature on production of oligosaccharides and dextran by *Weissella cibaria* 10 M. *Int J Food Microbiol* 280:27–34. <https://doi.org/10.1016/j.ijfoodmicro.2018.05.003>
72. Kanimozhi J, Ganesh Moorthy I, Sivashankar R, Sivasubramanian V (2017) Optimization of dextran production by *Weissella cibaria* NITCSK4 using response surface methodology-genetic algorithm based technology. *Carbohydr Polym* 174:103–110. <https://doi.org/10.1016/j.carbpol.2017.06.021>
73. Devi PB, Kavitate D, Shetty PH (2016) Physico-chemical characterization of galactan exopolysaccharide produced by *Weissella confusa* KR780676. *Int J Biol Macromol* 93:822–828. <https://doi.org/10.1016/j.ijbio mac.2016.09.054>
74. Ye G, Chen Y, Wang C et al (2018) Purification and characterization of exopolysaccharide produced by *Weissella cibaria* YB-1 from pickle Chinese cabbage. *Int J Biol Macromol* 120:1315–1321. <https://doi.org/10.1016>
75. Yu Y-J, Chen Z, Chen PT, Ng I-S (2018) Production, characterization and antibacterial activity of exopolysaccharide from a newly isolated *Weissella cibaria* under sucrose effect. *J Biosci Bioeng* 126:769–777. <https://doi.org/10.1016/j.jbiosc.2018.05.028>
76. Zhu Y, Wang C, Jia S et al (2018) Purification, characterization and antioxidant activity of the exopolysaccharide from *Weissella cibaria* SJ14 isolated from Sichuan paocai. *Int J Biol Macromol* 115:820–828. <https://doi.org/10.1016/j.ijbio mac.2018.04.067>
77. Baruah R, Deka B, Goyal A (2017) Purification and characterization of dextransucrase from *Weissella cibaria* RBA12 and its application in in vitro synthesis of prebiotic oligosaccharides in mango and pineapple juices. *LWT - Food Sci Technol* 84:449–456. <https://doi.org/10.1016/j.lwt.2017.06.012>
78. Shi Q, Juvonen M, Hou Y et al (2016) Lactose- and cellobiose derived branched trisaccharides and a sucrose-containing trisaccharide produced by acceptor reactions of *Weissella confusa* dextransucrase. *Food Chem* 190:226–236. <https://doi.org/10.1016/j.foodchem.2015.05.090>
79. Nolte J, Kempa A, Schlockermann A et al (2019) Glycosylation of caffeic acid and structural analogues catalyzed by novel glucansucrases from *Leuconostoc* and *Weissella* species. *Biocatal Agric Biotechnol* 19. <https://doi.org/10.1016/j.bcab.2019.101114>

80. Rosca I, Petrovici AR, Peptanariu D et al (2018) Biosynthesis of dextran by *Weissella confusa* and its in vitro functional characteristics. *Int J Biol Macromol* 107:1765–1772. <https://doi.org/10.1016/j.ijbio mac.2017.10.048>
81. Kavitate D, Devi PB, Shetty PH (2016) Characterization of a novel galactan produced by *Weissella confusa* KR780676 from an acidic fermented food. *Int J Biol Macromol* 86:681–689. <https://doi.org/10.1016/j.ijbio mac.2016.01.099>
82. Baruah R, Maina NH, Katina K et al (2016) Functional food applications of dextran from *Weissella cibaria* RBA12 from pummelo (*Citrus maxima*). *Int J Food Microbiol* 242:124–131. <https://doi.org/10.1016/j.ijfoo dmicr o.2016.11.012>
83. Zannini E, Waters DM, Coffey A, Arendt EK (2016) Production, properties, and industrial food application of lactic acid bacteria-derived exopolysaccharides. *Appl Microbiol Biotechnol* 100:1121–1135. <https://doi.org/10.1007/s0025 3-015-7172-2>
84. Zannini E, Jeske S, Lynch KM, Arendt EK (2018) Development of novel quinoa-based yoghurt fermented with dextran producer *Weissella cibaria* MG1. *Int J Food Microbiol* 268:19–26. <https://doi.org/10.1016/j.ijfoo dmicr o.2018.01.001>
85. Rolim PM, Hu Y, Ganzle MG (2019) Sensory analysis of juice blend containing isomalto-oligosaccharides produced by fermentation with *Weissella cibaria*. *Food Res Int* 124:86–92. <https://doi.org/10.1016/j.foodr es.2018.08.089>
86. Tang X, Zhang B, Huang W et al (2019) Hydration, water distribution and microstructure of gluten during freeze thaw process: Role of a high molecular weight dextran produced by *Weissella confusa* QS813. *Food Hydrocoll* 90:377–384. <https://doi.org/10.1016/j.foodh yd.2018.10.025>
87. Xu Y, Pitkanen L, Maina NH et al (2018) Interactions between fava bean protein and dextrans produced by *Leuconostoc pseudomesenteroides* DSM 20193 and *Weissella cibaria* Sj 1b. *Carbohydr Polym* 190:315–323. <https://doi.org/10.1016/j.carbp ol.2018.02.082>
88. Xu Y, Wang Y, Coda R et al (2017) In situ synthesis of exopolysaccharides by *Leuconostoc* spp. and *Weissella* spp. and their rheological impacts in fava bean flour. *Int J Food Microbiol* 248:63–71. <https://doi.org/10.1016/j.ijfoo dmicr o.2017.02.012>
89. Mathur S, Singh R (2005) Antibiotic resistance in food lactic acid bacteria - a review. *Int J Food Microbiol* 105:281–295. <https://doi.org/10.1016/j.ijfoo dmicr o.2005.03.008>
90. Toomey N, Bolton D, Fanning S (2010) Characterisation and transferability of antibiotic resistance genes from lactic acid bacteria isolated from Irish pork and beef abattoirs. *Res Microbiol* 161:127–135. <https://doi.org/10.1016/j.resmi c.2009.12.010>
91. Džidić S, Šušković J, Kos B (2008) Antibiotic resistance mechanisms in bacteria: biochemical and genetic aspects. *Food Technol Biotechnol* 46:11–21
92. Ammor MS, Belen Florez A, Mayo B (2007) Antibiotic resistance in non-enterococcal lactic acid bacteria and bifidobacteria. *Food Microbiol* 24:559–570. <https://doi.org/10.1016/j.fm.2006.11.001>
93. Perin LM, Miranda RO, Todoroc SD et al (2014) Virulence, antibiotic resistance and biogenic amines of bacteriocinogenic lactococci and enterococci isolated from goat milk. *Int J Food Microbiol* 185:121–126. <https://doi.org/10.1016/j.ijfoo dmicr o.2014.06.001>
94. Fhoula I, Rehaïem A, Najjari A et al (2018) Functional probiotic assessment and in vivo cholesterol-lowering efficacy of *Weissella* sp. associated with arid lands living-hosts. *Biomed Res Int* 2018:11 pages. <https://doi.org/10.1155/2018/16541 51>

95. Mortezaei F, Royan M, Noveirian HA et al (2020) In vitro assessment of potential probiotic characteristics of indigenous *Lactococcus lactis* and *Weissella oryzae* isolates from rainbow trout (*Oncorhynchus mykiss* Walbaum). *J Appl Microbiol* 129:1004–1019. <https://doi.org/10.1111/jam.14652>
96. Quattrini M, Korcari D, Ricci G, Fortina MG (2019) A polyphasic approach to characterize *Weissella cibaria* and *Weissella confusa* strains. *J Appl Microbiol* 128:500–512. <https://doi.org/10.1111/jam.14483>
97. Kang MS, Yeu JE, Hong SP (2019) Safety evaluation of oral care probiotics *Weissella cibaria* CMU and CMS1 by phenotypic and genotypic analysis. *Int J Mol Sci* 20:22 pages. <https://doi.org/10.3390/ijms20112693>
98. Figueiredo SP, Boari CA, de Souza Costa Sobrinho P et al (2015) Características do leite cru e do queijo Minas artesanal do Serro em diferentes meses. *Arch Vet Sci* 20:68–82. <https://doi.org/10.5380/avs.v20i1.37243>
99. Ortega JCC, Fajardo R, Irgang R et al (2018) Isolation characterization, virulence potential of *Weissella ceti* responsible for weissellosis outbreak in rainbow trout (*Oncorhynchus mykiss*) cultured in Mexico. *Transbound Emerg Dis* 65:1401–1407. <https://doi.org/10.1111/tbed.12978>
100. Welch TJ, Good CM (2013) Mortality associated with weissellosis (*Weissella* sp.) in USA farmed rainbow trout: potential for control by vaccination. *Aquaculture* 388–391:122–127. <https://doi.org/10.1016/j.aquaculture.2013.01.021>
101. Aberkane S, Didelot MN, Carriere C et al (2017) Bacteriemie a *Weissella confusa*: un pathogene opportuniste sous-estime. *Med Mal Infect* 47:297–299. <https://doi.org/10.1016/j.medmal.2017.02.001>
102. Freedman SB, Williamson-Urquhart S, Farion KJ et al (2018) Multicenter trial of a combination probiotic for children with gastroenteritis. *N Engl J Med* 379:2015–2026. <https://doi.org/10.1056/NEJMoa1802597>
103. Schnadower D, Tarr PI, Casper T et al (2018) *Lactobacillus rhamnosus* GG versus placebo for acute gastroenteritis in children. *N Engl J Med* 379:2002–2014. <https://doi.org/10.1056/NEJMoa1802598>
104. Laulund S, Wind A, Derkx PMF, Zuliani V (2017) Regulatory and safety requirements for food cultures. *Microorganisms* 5:28. <https://doi.org/10.3390/microorganisms5020028>
105. Sturino JM (2018) Literature-based safety assessment of an agriculture- and animal-associated microorganism: *Weissella confusa*. *Regul Toxicol Pharmacol* 95:142–152. <https://doi.org/10.1016/j.yrtph.2018.03.013>
106. Cupi D, Elvig-Jorgensen SG (2019) Safety assessment of *Weissella confusa* – a direct-fed microbial candidate. *Regul Toxicol Pharmacol* 107:104414. <https://doi.org/10.1016/j.yrtph.2019.104414>
107. Panel EB, Koutsoumanis K, Allende A et al (2019) The list of QPS status recommended biological agents for safety risk assessments carried out by EFSA. <https://doi.org/10.5281/ZENODO.3336268>
108. Tenea GN, Hurtado P, Ortega C (2020) A novel *Weissella cibaria* strain UTNGt210 isolated from wild solanum quitoense fruit: genome sequence and characterization of a peptide with highly inhibitory potential toward Gram-negative bacteria. *Foods* 9(9):1242. <https://doi.org/10.3390/foods9091242>
109. Pelyuntha W, Chaiyasut C, Kantachote D, Sirilun S (2020) Organic acids and 2,4-Di-tert-butylphenol: major compounds of *Weissella confusa* WM36 cell-free supernatant against

growth, survival and virulence of *Salmonella* Typhi. PeerJ 8:e8410. <https://doi.org/10.7717/peerj.8410>

110. Lakra AK, Domdi L, Tilwani YM, Arul V (2020) Physicochemical and functional characterization of mannan exopolysaccharide from *Weissella confusa* MD1 with bioactivities. Int J Biol Macromol 143:797–805. <https://doi.org/10.1016/j.ijbio mac.2019.09.139>

CHAPTER II: Biodiversity and technological features of *Weissella* isolates obtained from Brazilian artisanal cheese-producing regions

Biodiversity and technological features of *Weissella* isolates obtained from Brazilian artisanal cheese-producing regions

The content of this part has been published in:

LWT - Food Science and Technology

Camila Gonçalves Teixeira, Andressa Fusieger, Evandro Martins, Rosângela de Freitas, Martina Vakarelova, Luís Augusto Nero, Antônio Fernandes de Carvalho

License to reuse this content in thesis is in Annex 3

Abstract

Weissella spp. strains (n = 57) were isolated from different Brazilian food-related environments and characterized based on their genetic profiles and technological potential. PFGE and Rep-PCR revealed the high level of biodiversity among isolates: PFGE grouped the isolates in three major profiles (similarity from 60 to 80%), while rep-PCR characterized the isolates as belonging to a single profile. Based on PFGE, isolates with identical pulsotypes were found within a same geographical region (Campo das Vertentes, Minas Gerais state). Based on these profiles, 26 isolates were selected and characterized based on their inhibitory and bacteriocinogenic activity against *Listeria monocytogenes* ATCC 15313, *Staphylococcus aureus* subsp. *aureus* ATCC 6538, *Salmonella enterica* subsp. *enterica* serovar Typhimurium ATCC 14028 and *Escherichia coli* ATCC 11229. Most isolates (n = 20) were able to inhibit the targets through organic acids. Most of the isolates (n = 12) were able to produce diacetyl, proteases, and/or coagulating milk, but none were able to produce exopolysaccharides. Isolate 16, in particular, was characterized as possessing high acidification ability, diacetyl and protease production, with promising technological potential in the dairy industry. The obtained results allowed the understanding of the *Weissella* spp. strains role as starter cultures in the dairy industry.

Keywords: *Weissella*; dairy microbiota; diacetyl profile

1. Introduction

Lactic acid bacteria (LAB) is a microbial group naturally present in a wide variety of foods. LAB play an important role in the production of fermented foods due to their ability to produce aroma, flavor and texture. *Weissella* spp. are a LAB known for lactic acid production acid through carbohydrate fermentation and for its phylogenetic position characterized as Gram-positive, catalase negative, non-spore forming, with coccoid morphology or short bacilli (Björkroth, Dicks, & Holzapfel, 2009; Collins, Samelis, Metaxopoulos, & Wallbanks, 1993).

The genus *Weissella* is made up of 25 validated species, most of which have already been identified in fermented foods such as vegetables, sausages and dairy products, and particularly raw milk cheeses (Jans, Bugnard, Njage, Lacroix, & Meile, 2012; Masoud et al., 2012; Pereira, San Romão, Lolkema, & Barreto Crespo, 2009; Wouters, Ayad, Hugenholtz, & Smit, 2002). In Brazil, *Weissella* spp. were previously been isolated from artisanal cheeses produced in Pará, Minas Gerais and Paraíba states (Martins et al., 2018; Medeiros et al., 2016; Perin, Savo Sardaro, Nero, Neviani, & Gatti, 2017).

Weissella species present many positive attributes for the food industry. Indeed, its wide array of characteristics have been attributed to the strains of this genus. Studies have shown that *Weissella* spp. Strains can be used as starter cultures and to produce antimicrobial substances and exopolysaccharides (Ayeni et al., 2011; Choi, Kim, Hwang, Kim, & Yoon, 2012; Srionnual, Yanagida, Lin, Hsiao, & Chen, 2007; Zannini et al., 2013). Specifically, *W. paramesenteroides* and *W. hellenica*, usually found in dairy products, have been widely studied for their antimicrobial potential, since certain strains have already been characterized as bacteriocinogenic (Chen et al., 2014; Leong et al., 2013; Masuda et al., 2012; Papagianni & Papamichael, 2011). *W. hellenica* was detected in dairy products such as raw milk Croatian cheese, while *W. paramesenteroides* was previously isolated from raw milk artisanal cheeses from different countries, thus demonstrating the relevance of this genus for dairy production (Coppola et al., 2006; Fuka et al., 2013; Gerasi, Litopoulou-Tzanetaki, & Tzanetakis, 2003; Masuda et al., 2012).

Although the phenotypic characteristics of *Weissella* spp. have been studied and *Weissella* spp. strains have been isolated and identified in artisanal cheeses, there is little information on this genus found in Brazilian artisanal cheeses. *Weissella* spp.

function in such products and this genus biodiversity from region to region in Brazil remain unknown. Thus, the present study aimed to identify the microbial biodiversity and evaluate the technological potential of *Weissella* spp. isolates obtained from different Brazilian regions that produce artisanal cheeses.

2. Material and methods

Microorganisms and culture conditions

Fifty-seven (n = 57) *Weissella* spp. isolates were obtained from the InovaLeite (Laboratory of Milk and Dairy Products, Universidade Federal de Viçosa) bacterial culture collection and included in this study. The isolates were taken from samples from four different Brazilian regions that produce artisanal cheeses (Supplementary Table 1). The isolates have been previously identified by 16S rRNA sequencing (Felske, Rheims, Wolterink, Stackebrandt, & Akkermans, 1997) as: *W. cibaria* (n = 38), *W. paramesenteroides* (n = 11) and *W. confusa* (n = 8) (Fig. 1). The isolates obtained were stored at -60 °C in de Man Rogosa and Sharpe (MRS) broth (Oxoid Ltd., Basingstoke, England) and supplemented with 30% (v/v) glycerol (Synth, São Paulo, SP, Brazil). Prior to analysis, stock cultures (1% v/v) were transferred to MRS broth (Oxoid) and incubated at 30 °C for 14 h.




Fig 1. Strains, biotope and geographical origins and identification of *Weissella* spp. isolates obtained from Brazilian artisanal cheese-producing regions.

Biodiversity determination using molecular cluster analysis

2.2.1. PFGE characterization of *Weissella* spp.

Pulsed-Field Gel Electrophoresis (PFGE) was performed as described by Chuat and Dalmaso (2015), with modifications. *Weissella* spp. isolates were grown to an optical density ($\lambda = 650$ nm) of 0.3 in MRS broth (Oxoid). Cells were obtained by centrifuging (3500×g for 10 min at 20 °C) 10 mL of culture. The supernatant was discarded and 5 mL of a TES buffer (10 mmol/L Tris, 1 mmol/L EDTA, pH 8.0, 0.5 mol/L sucrose) was added to the pellet, then homogenized. The pellet was mixed with a lysis solution to obtain a final concentration of 10 g/L of lysozyme (Sigma, St. Louis, MO, USA) and the mixture was centrifuged once more. After an incubation period at 37 °C for 1 h, the suspension was heated at 55 °C for 1 min and 700 μ L of agarose solution (ultrapure agarose, Gibco-BRL, Paisley, United Kingdom) 1% (w/v) in 125 mmol/L EDTA (pH 7.0) at the same temperature were added to the suspension. The mixture was placed in the molds and solidified at 4 °C for 10 min. The blocks were placed in a deproteinization buffer solution (10 mmol/L Tris, 100 mmol/L EDTA, pH 8.0, 10% SDS, 20 g/L proteinase K, Sigma) for 2 h at 55 °C. The agarose blocks were rinsed in sterile water at 55 °C for 10 min and four times in TE buffer (pH 8.0), for 10 min per wash. The agarose blocks were stored at 4 °C in TE Buffer (pH 8.0) until digestion.

For digestion, the DNA in the agarose blocks was cleaved with 10 U of restriction enzyme SmaI (Promega Corp., Madison, WI, USA) for 4h at 24 °C, and transferred to 1% (w/v) agarose gel (PFGE certified agarose, Bio-Rad, Hercules, CA, USA). DNA fragments were separated by PFGE in 0.5 x TBE buffer (45 mmol/L Tris, 45 mmol/L borate, 1.0 mmol/L EDTA, pH 8.3) at 14 °C in a CHEF DR II apparatus (Bio-Rad) with a pulse of 2–25 s, voltage 6 V/cm, for 18 h at 14 °C. The gel was stained with GelRed (3x in 0.1 M NaCl solution) (Biotium Inc., Hayward, CA, USA) for 15 min and the images were taken under ultraviolet light. PFGE gel images were obtained and the band profiles were analyzed using BioNumerics software, version 6.6 (Applied Maths, Kortrijk, Belgium). The normalized band profiles were compared using the Dice similarity coefficient (1%). Cluster analysis for the compiled matrix was established using the unweighted pair group method and an arithmetic average (UPGMA) clustering algorithm.

2.2.2. Rep-PCR characterization of *Weissella* spp.

Aliquots of 1 mL of *Weissella* spp. cultures were centrifuged at 10,000×g for 10 min at 4 °C, and the obtained pellets were subjected to DNA extraction using a DNA Purification Wizard® Genomic kit (Promega). DNA concentration was standardized at 100 ng/μL (NanoDrop™ Lite Spectrophotometer, Thermo Fisher Scientific, Waltham, MA, USA).

Rep-PCR analysis was performed according to the protocol described by Dal Bello et al. (2010) using a single universal primer (GTG)₅ (5'-GTGGTGGTGGTGGTG-3'). PCR reactions contained 12.5 μL of Go Taq Green Master Mix 2x (Promega), 0.5 μL of the primer (100 mol/L), 100 ng DNA and ultra-pure water (Promega) for a final volume of 25 μL. PCR amplification was carried out in a thermal cycler, with the following conditions: 95 °C for 5 min as an initial step, 95 °C for 30 s, annealing at 40 °C for 30 s, then 65 °C for 8 min for the next 30 cycles, and 65 °C for 16 min to conclude the amplification.

The PCR products were electrophoresed on 2% (w/v) agarose gel for 2 h at a constant voltage of 75 V in 0.5x TBE buffer. The gels were stained using GelRed (Biotium Inc.) and developed using an LPIX transilluminator (Loccus Biotechnology, São Paulo, SP, Brazil). The band profile was analyzed using BioNumerics software 6.6 (Applied Maths). The similarities between the profiles were calculated using the Dice similarity coefficient (1%). Dendrograms were constructed using the UPGMA method.

Antagonism analysis and bacteriocin production

Based on the fingerprinting profiles, a representative number of isolates (n = 26) were selected for antimicrobial activity evaluation. *Weissella* spp. isolates were cultured in MRS broth (Oxoid) at 30 °C for 24 h and a cell-free supernatant (CFS) was obtained by centrifugation (10,000×g for 10 min at 4 °C) and filtration (Ø = 0.45 μm). The strains' CFS levels were tested using microdilution assays (Arena et al., 2016; Vijayakumar & Muriana, 2015) to determine antagonism against *Listeria monocytogenes* ATCC 15313, *Staphylococcus aureus* subsp. *aureus* ATCC 6538, *Salmonella enterica* subsp. *enterica* serovar Typhimurium ATCC 14028 and *Escherichia coli* ATCC 11229. Target strains were cultured in a brain heart infusion (BHI, Oxoid) at 37 °C for 24 h and aliquots (100 μL, at a concentration of about 10⁴ CFU/mL) were transferred to 96-well microtiter plates. Then, 50 μL aliquots of the

selected *Weissella* spp. isolates' CFS were added to each well. The microplates were incubated at 37 °C for 24 h in a Multiskan™ GO Microplate Spectrophotometer (Thermo Fisher). The cultures' optical densities (OD) were measured every 30 min (λ = 600 nm) (Arena et al., 2016; Vijayakumar & Muriana, 2015). The CFS pH was measured both before (pH0h) and after incubation (pH24h), and results were presented as the difference of both records (Δ pH = pH0h – pH24h). Growth controls were prepared in wells added with the target culture (100 μ L) and MRS broth (50 μ L, Oxoid). Negative controls were prepared in wells added with BHI (150 μ L, Oxoid) alone. Inhibition assays were conducted in triplicate and over two independent repetition. Recorded ODs were corrected based on negative controls, and OD mean values calculated over the reading time; OD from positive controls (ODc) were compared with OD from CFS treated cultures (ODt) and inhibition values were calculated based on the following equation:

$$\text{Inhibition} = 100 - (\text{ODt} \times 100/\text{ODc})$$

Based on the antagonism test results, isolates that presented antimicrobial activity higher than 60% were selected to verify their bacteriocinogenic activity. CFS of the selected isolates was obtained as previously described. Furthermore, it was treated as following: pH was adjusted to 6.5 with 1 mol/L NaOH to neutralize the medium and a heat treatment was applied for 10 min, at 80 °C, to eliminate peroxides (Cavicchioli, Camargo, Todorov, & Nero, 2017). The treated CFS were tested using the same target strains as described above. The same positive and negative controls were used, and the same result interpretation methods were used. This assay was conducted in triplicate and over two independent repetitions.

Technological potential

Selected isolates (n = 13) were also subjected to phenotypical analysis in order to characterize the following technological features: diacetyl production, coagulation, proteolysis, acidifying activity and exopolysaccharide (EPS) production. Before each assay, the isolates were cultured in MRS (Oxoid) at 30 °C for 14 h and centrifuged at 14,000 \times g for 2 min at 4 °C. After the supernatant was discarded, the cell pellets were suspended in NaCl 0.85% (w/v) until a turbidity similar to McFarland tube 1 (approximately 3×10^8 CFU/mL). The obtained cell suspension of the cultures were used in the assays described below, which were conducted in triplicate and over two

independent repetitions. MilliQ water was used as a negative control. To assess diacetyl production, 0.1 mL aliquots of the cell suspension of the cultures were transferred to 10 mL of sterile reconstituted skim milk (10% w/v, Nestlé, São Paulo, SP, Brazil), and incubated at 30 °C for 24 h. Then, 1 mL aliquots of the obtained cultures were added to 0.5 mL of α -naphthol (1% w/v) and KOH (16% w/v) and incubated at 30 °C for 10 min: the formation of a red ring at the top of the tubes indicated diacetyl production (Franciosi, Settanni, Cavazza, & Poznanski, 2009). *Lactococcus lactis* subsp. *lactis* bv. *diacetylactis* ATCC 13675 was used as positive control. Milk coagulation patterns were assessed by inoculating 0.1 mL aliquots of the cell suspension of the cultures into 10 mL of sterile reconstructed skim milk (10% w/v, Nestlé), followed by incubation at 30 °C for 24 h. Based on the formed clot characteristics, the coagulation patterns were described using an empirical analysis and classified as: uniform, uniform with presence of serum, uniform and fragile (appearance), broken with presence of serum, and absence of clot (Fusieger, Martins, de Freitas, Nero, & de Carvalho, 2020). *L. lactis* subsp. *lactis* bv. *diacetylactis* Q1C5 was used as positive control. A screening assay was conducted to identify presumed extracellular proteolytic activity of the isolates as described by Franciosi et al. (2009). Two μ L aliquots of bacterial cultures were spotted onto the surface of a Plate Count Agar (PCA, HiMedia, Mumbai, MH, India) added to skim milk (10% w/v, Nestlé), and the cultures were incubated at 30 °C for 4 days. Presumed proteolytic activity was indicated by a clear zone around the colonies and *Pseudomonas fluorescens* 07A (Alves, Salgado, Eller, Vidigal, & Carvalho, 2016) was used as positive control. EPS production from lactose present in milk was evaluated according to protocol described by Dal Bello et al. (2012). 0.1 mL aliquots of selected cell suspension of the cultures were inoculated into 10 mL of sterile reconstituted skim milk (10% v/v, Nestlé) and incubated at 30 °C for 48 h. Then, the EPS production was determined by the quality and degree of stringiness: isolates were characterized as EPS producers when the coagulated cultures could be tied to a string with an inoculating loop. Acidifying activity was assessed by adding 100 μ L of the selected cell suspension of the cultures to skim milk (10% w/v, Nestlé), followed by an incubation at 30 °C for 24 h pH values were measured at the time of inoculation (T0), then after 6 (T6), 12 (T12), and 24 h (T24) using a digital pHmeter (Hanna Instruments, São Paulo, SP, Brazil). Also, titratable acidity was expressed as lactic acid (Dal Bello et al., 2012). Results have been shown as means of the triplicates and repetitions.

2. Results

Biodiversity by molecular analysis

Following *Sma*I macro-restriction, the *Weissella* spp. isolates presented 51 pulsotypes. These were grouped in three major clusters with similarity indexes among the grouped isolates varying from 60 to 80% (Fig. 2).




Fig 2. Dendrograms based on UPGMA clustering of *Sma*I PFGE profile of *Weissella* spp. isolates obtained from Brazilian artisanal cheese-producing regions.

Clusters I and II included isolates from different species and from all four geographical regions, while cluster III was composed only of isolates from Campos das Vertentes and Southern Pará. Also, cluster III presented the highest index similarity among isolates (around 80%), and included isolates from different origins that shared

identical pulsotypes: isolates 28, 29, 46 and 53, obtained from silage, pasture and cow milk (*W. cibaria*), respectively, and isolates 26, 27 and 44, obtained from soil and pasture samples (*W. cibaria*).

Based on the rep-PCR tests, the isolates presented genetic profiles with bands ranging from 500 to 5000 bp. This allowed the characterization of 45 different profiles (Fig. 3).




Fig 3. Dendrograms generated after cluster analysis of rep-PCR fingerprints of *Weissella* spp. obtained from Brazilian artisanal cheese-producing regions.

Unlike PFGE, the rep-PCR profiles were visually more diverse, which allowed the isolates to be clustered into two major groups, with similarity indexes among the included isolates ranging from 35 to 45%. Isolates that presented identical profiles according to rep-PCR analysis did not show the same SmaI restriction profiles (Figs.

2 and 3). However, as we had observed for the PFGE results, rep-PCR allowed the isolates with identical genetic profiles from different samples to be identified. As examples, see: samples 25, 29, 35, 39 and 54, from soil, pasture and cow milk (*W. cibaria*); 20, 22 and 37, from soil and pasture (*W. cibaria*); 24 and 38, from pasture and soil (*W. cibaria*); 26, 43 and 45, from soil and pasture (*W. cibaria*), and 2, 5 and 7, from fermented curd, cow milk and utensils (*W. confusa*).

Inhibitory activity

Most of the tested isolates demonstrated antagonism against the target strains, with a minimum level of 60% of inhibitory activity (Table 1). The pH range of the inoculated MRS after the isolates were added and incubated for 24 h is shown in Table 1. *L. monocytogenes* ATCC 15313 was susceptible to all isolates from Marajó Island and Guanambi. Only isolate 16, from Southern Pará, did not demonstrate any antagonistic activity above 60%. *S. aureus* ATCC 6538, *S. Typhimurium* ATCC 14028 and *E. coli* ATCC 11229 were greatly inhibited by almost all of the isolates from Marajó Island, Southern Pará and by all isolates from Guanambi. Isolate 21 (*W. cibaria*) presented the highest inhibition level against *L. monocytogenes* ATCC 15313. Isolate 11 (*W. paramesenteroides*), from Southern Pará, presented the inhibitory activity against *S. aureus* ATCC 6538 and *S. Typhimurium* ATCC 14028. Despite the described inhibitory activity in the antagonism test, for the test with treated CFS (neutralized and heat treated) none isolate presented bacteriocinogenic activity against the target strains (data not shown).

Technological features

Among the 13 tested *Weissella* isolates, six exhibited diacetyl production (Table 2): isolate 25 (*W. cibaria*) results exhibited the most intense red color (data not shown). Four of these six isolates were obtained from cheese (Southern Pará) and two from pasture samples (Campo das Vertentes). *W. confusa* isolates demonstrated no diacetyl production.

In our research, four of the *Weissella* spp. isolates tested showed uniform and consistent clotting after milk coagulation (Table 2). Two demonstrated homogeneous coagulation (16 and 25, *W. paramesenteroides* and *W. cibaria*, respectively) while the other two isolates (17 and 22, *W. paramesenteroides* and *W. cibaria*, respectively)

showed homogeneous clots, but only after 48 h of incubation. The other isolates did not coagulate milk.

Presumed extracellular proteolytic activity was observed in eleven *Weissella* isolates (Table 2); five of these isolates were obtained from artisanal cheeses (Southern Pará). EPS production was determined qualitatively and all strains proved to be negative (Table 2). Most samples showed some ability to acidify skim milk, producing a weak reduction of pH after 24 h of incubation, except for isolate 16 (*W. paramesenteroides*) which lowered milk pH by over 2.0 pH units after 12 h (Table 2). The titratable acidity results expressed as lactic acid content confirm the pH measurement results. (Table 2).

Table 1. Inhibitory potential of the cell-free supernatant of *Weissella* spp. isolates against target strains.

Codes	Identification	<i>L. monocytogenes</i> ATCC 15313	<i>S. aureus</i> ATCC 6538	<i>S. Typhimurium</i> ATCC 14028	<i>E. coli</i> ATCC 11229	pH
1	<i>W. confusa</i>	89.2	53.0	80.6	67.5	4.12
3	<i>W. confusa</i>	88.1	37.0	77.3	71.1	4.35
5	<i>W. confusa</i>	88.7	86.4	78.1	86.1	4.03
8	<i>W. confusa</i>	85.3	85.2	76.6	87.9	3.95
9	<i>W. paramesenteroides</i>	62.4	25.0	50.9	29.2	4.03
10	<i>W. paramesenteroides</i>	89.4	85.1	81.9	84.1	4.04
11	<i>W. paramesenteroides</i>	90.3	89.2	83.0	85.6	3.94
12	<i>W. paramesenteroides</i>	87.1	79.8	77.5	83.6	4.04
14	<i>W. paramesenteroides</i>	88.0	86.6	79.5	87.3	4.01
15	<i>W. paramesenteroides</i>	60.1	45.0	26.7	49.5	4.74
16	<i>W. paramesenteroides</i>	31.8	36.8	43.4	46.3	4.36
17	<i>W. paramesenteroides</i>	88.0	84.0	79.5	85.4	4.00
18	<i>W. paramesenteroides</i>	88.1	68.8	78.1	65.6	4.17
19	<i>W. paramesenteroides</i>	88.6	88.1	82.0	86.1	3.94
21	<i>W. cibaria</i>	98.5	87.1	76.8	82.8	4.79
22	<i>W. cibaria</i>	39.4	39.9	15.2	63.4	4.49
25	<i>W. cibaria</i>	44.0	22.2	49.9	18.6	4.41
30	<i>W. cibaria</i>	48.7	28.9	26.3	40.2	4.67
32	<i>W. cibaria</i>	26.6	37.0	4.2	21.8	4.71
35	<i>W. cibaria</i>	86.8	65.9	74.7	70.0	4.20
42	<i>W. cibaria</i>	89.1	88.5	80.3	86.2	3.92
49	<i>W. cibaria</i>	85.2	27.4	0.7	20.6	4.72
51	<i>W. cibaria</i>	86.7	63.8	78.4	74.0	4.01
52	<i>W. cibaria</i>	11.4	4.0	8.5	2.2	4.46
54	<i>W. cibaria</i>	78.5	30.5	70.7	28.0	4.75
56	<i>W. cibaria</i>	87.9	48.4	77.2	56.6	4.39

Percentage of inhibition is expressed as (%) = $100 - (\text{mean OD}_{600 \text{ nm}} / \text{mean OD}_{600 \text{ nm}} \text{ target strain}) * 100$

Table 2. Technological features of *Weissella* spp. isolates obtained from Brazilian artisanal cheese-producing regions

Codes	Identification	Diacetyl	Coagulation	Proteolysis	EPS	Acidifying activity							
						pH				% lactic acid			
						T0	T6	T12	T24	T0	T6	T12	T24
1	<i>W. confusa</i>	-	absence of clot	+	-	6.85	6.50	6.44	6.15	0.14	0.15	0.13	0.19
5	<i>W. confusa</i>	-	absence of clot	+	-	6.85	6.59	6.26	6.15	0.14	0.11	0.16	0.19
10	<i>W. paramesenteroides</i>	-	absence of clot	+	-	6.85	6.62	6.40	6.33	0.14	0.13	0.11	0.20
12	<i>W. paramesenteroides</i>	+	absence of clot	+	-	6.85	6.60	6.42	5.46	0.14	0.13	0.13	0.23
14	<i>W. paramesenteroides</i>	+	absence of clot	+	-	6.85	6.63	6.45	6.27	0.14	0.12	0.14	0.18
15	<i>W. paramesenteroides</i>	+	absence of clot	+	-	6.85	6.62	6.43	6.36	0.14	0.13	0.17	0.16
16	<i>W. paramesenteroides</i>	+	uniform	+	-	6.85	5.41	4.65	4.60	0.14	0.37	0.57	0.77
17	<i>W. paramesenteroides</i>	-	uniform	+	-	6.85	6.53	6.14	5.84	0.14	0.12	0.20	0.27
18	<i>W. paramesenteroides</i>	-	absence of clot	+	-	6.85	6.69	6.53	6.37	0.14	0.12	0.13	0.17
19	<i>W. paramesenteroides</i>	-	absence of clot	+	-	6.85	6.57	6.40	6.24	0.14	0.13	0.15	0.18
22	<i>W. cibaria</i>	+	uniform	+	-	6.85	6.30	5.78	5.35	0.14	0.17	0.29	0.38
25	<i>W. cibaria</i>	+	uniform	-	-	6.85	6.47	5.58	5.00	0.14	0.15	0.35	0.60
54	<i>W. cibaria</i>	-	absence of clot	-	-	6.85	6.77	6.73	6.66	0.14	0.12	0.13	0.12

Results: +, positive; -, negative

3. Discussion

Based on the profiles obtained by PFGE and rep-PCR (Figs. 2 and 3), the studied isolates presented high variability; the two techniques grouped the isolates differently, which is to be expected because the approaches used are totally different. Nisiotou, Dourou, Filippousi, Banilas, and Tassou (2014) have described differences found in *W. uvarum* groupings obtained using PFGE and rep-PCR in a study on grape isolates (Nemea region, Greece). rep-PCR testing presented a greater number of genetic profiles compared to PFGE (23 and 29, respectively), and the profiles had fewer similarities compared to the results obtained using PFGE. These results have also been observed by other authors who used both rep-PCR and PFGE techniques with different lactic bacteria (Cavicchioli et al., 2015; Perin & Nero, 2014). In general, clustering obtained by PFGE confirmed the rep-PCR results, where clustering tendencies between isolates species and regions have been observed (Figs. 2 and 3). However, similarity and distribution levels within the groups were not the same for the two techniques: PFGE was able to establish clearer associations between similar isolates, thus demonstrating better discriminatory properties and genetic characterizations of these isolates. As PFGE and rep-PCR require very different approaches, the first based on the digestion of entire DNA and the second based on the amplification of specific DNA regions, the profiles recorded were expected to be different (Tenover et al., 1995). Freitas et al. (2015) compared PFGE to random amplified polymorphic DNA (RAPD), another molecular technique, and concluded that PFGE provides better heterogeneity resolution between strains of the same species. Thus, in this study, the PFGE technique was chosen to screen select samples from each cluster for antimicrobial activity assays.

The three *Weissella* species studied were present in almost all types of original samples including milk, cheese, pasture, and utensils. This demonstrates the high diversity of strains found in the regions studied. Fuka et al. (2013) and others have also identified *Weissella* strains in dairy environments and isolated *W. paramesenteroides* and *W. hellenica* in Croatian raw sheep's milk cheese. Masoud et al. (2012) have also isolated the same *Weissella* species in raw milk and raw milk cheeses. The regions studied represent specific ecological niches as isolated strains from different regions did not present similar PFGE restriction profiles (100% similarity). Freitas et al. (2015) in the research of *Propionibacterium* biodiversity obtained the same results with each

of the farms they studied in Campo das Vertentes, representing a different bacterial ecological niche.

Using PFGE, identical pulsotypes were found within the same geographical regions (Fig. 2). In addition to belonging to the same regional classification (Campo das Vertentes), these isolates were taken from very closely related and/or similar samples, such as isolates 20 and 21 from pasture and isolates 44, 26 and 27, from soil and pasture from the same region (Fig. 1). Martins et al. (2020) used PFGE to compare 23 isolates of *L. lactis* subsp. *lactis* isolated from dairy environment in four different regions in Brazil and showed that isolates presented identical profiles by PFGE. As here, the isolates belonged to the same region and similar samples.

The high susceptibility levels of *L. monocytogenes* to *Weissella* supernatants (Table 1) were justified because antimicrobial activity among related microorganisms is common in Gram-positive bacteria. *E. coli* was used as a Gram-negative target microorganism and it was also shown to be highly inhibited by supernatants. Ortolani, Yamazi, Moraes, Viçosa, and Nero (2010) also reported LAB antagonist activity for foodborne pathogens *L. monocytogenes* and *S. aureus* when they studied isolates taken from raw milk and soft cheese. When Espeche, Otero, Sesma, and Nader-Macias (2009) isolated 6 strains of *W. paramesenteroides* from bovine milk samples, one sample also showed antimicrobial activity for *Streptococcus dysgalactiae* subsp. *dysgalactiae* ATCC 27957 and pathogenic *E. coli*. It should be noted that only cells were removed from *Weissella* sp. supernatants in this test, and the cells were obtained under optimum growth conditions, what does not prevent the production of more than one antimicrobial compound. LAB can produce a variety of antimicrobial compounds including hydrogen peroxide, organic acids (lactic acid, acetic acid, formic acid, for example), carbon dioxide, bacteriocins, ethanol and diacetyl (Leroy & De Vuyst, 2004). The bacteriocin production test revealed that the isolates did not demonstrate any production capacity under the growth conditions (data not shown). However, *W. paramesenteroides* DX is able to produce the bacteriocin weissellin A when grown in M17 + 2% glucose broth at 30 °C for 60 h, according to Papagianni and Papamichael (2011). Sriannual et al. (2007) also incubated *W. cibaria* 110 in MRS broth at 30 °C and obtained bacteriocin weissellicin 110.

Decreases in pH indicate that acids were produced during the incubation of isolates in MRS. Because MRS includes dextrose and ammonium citrate in its formulation, it can be assumed that the acids produced by the strains are lactic acid

and acetic acid, via glycolytic and phosphoketolase pathways, as well as acetic acid via a diacetyl/acetoin pathway (Lynch et al., 2015). These organic acids have antimicrobial capacities when the pathogenic bacteria's cytoplasmic medium experiences a decrease in pH. According to Russell (1992), the non-dissociated form of organic acids is liposoluble and able to passively pass through cell cytoplasmic membranes. It changes the pH within cells by modifying the proton gradient and the electric charge when it interferes with the amino acid transport system and inactivates enzymes.

Weissella sp. are autochthonous to certain Brazilian raw milk artisanal cheeses and to their production environments. The fact that they present antimicrobial activity against important foodborne pathogens indicates the microbiota may interfere with pathogens present in raw milk and inhibit them. According to Jay (1996), the effects of autochthonous microorganisms on pathogens is well established. This effect is greater in animal-derived products containing a rich autochthonous microbiota.

The six isolates capable of producing diacetyl included two *W. cibaria* and four *W. paramesenteroides* (Table 2). Studies have shown that the *W. cibaria* MG1 strain isolated from pozol can produce diacetyl via a diacetyl/acetoin pathway (L'opez-Hernández, Rodríguez-Alegría, L'opez-Munguía, & Wachter, 2018). Additionally, Pakdeeto, Naranong, and Tanasupawat (2003) observed diacetyl production by *W. confusa* strains under incubation conditions in modified MRS broth. Diacetyl is a volatile compound generated as a final product when citrate is converted to pyruvate. It confers a buttery aroma to fermented dairy products and it is a common LAB feature that is usually associated with *L. lactis* subsp. *lactis* bv. *diacetylactis* (Fusieger, Martins, Freitas, Nero, & Carvalho, 2020; Leroy & De Vuyst, 2004; Smit, Smit, & Engels, 2005).

Of the four isolates that were able to coagulate milk, two were *W. cibaria* (Table 2). Some studies have shown the ability of *W. cibaria* strains to ferment lactose and galactose (Bounaix et al., 2010; Lynch et al., 2014). According to Lynch et al. (2015), *W. cibaria* MG1 possesses all the necessary genes for the phosphoketolase pathway and metabolizes galactose via the Leloir pathway. Milk coagulation is important in the production of fermented products such as yogurt and cheeses. Coagulation occurs when milk pH is lowered by LAB acids production. When the caseins' isoelectric point is reached, this adds to the effects of the expelled serum to form yogurt gels and cheese curds (Robinson, 2002; Tamime & Robinson, 2007).

Presumed proteolytic activity was observed for almost all the isolates selected in our study (Table 2). Lynch et al. (2014) have also evaluated the proteolysis activity of strains of *W. cibaria* MG1 using tests in milk agar and report that this enzymatic activity is strain-specific. LAB proteolytic and lipolytic activities in cheese lead to aroma and flavor generation and contribute to the release of bioactive peptides in milk that are beneficial to consumer health (Meisel & Bockelmann, 1999; Wouters et al., 2002).

According to Cogan et al. (1997), isolates with acidification capacities must reduce milk pH in 1.3 units after 6 h of incubation. The isolates examined in this study can be characterized as poor acid producers, except for isolate 16 (*W. paramesenteroides*), which was the only one able to reduce the pH to 5.41 after 6 h of incubation at 30 °C. It also produced a large amount of acid after 24 h of incubation (Table 2). Isolate 25 (*W. cibaria*) also produced a large amount of acid at the end of 24 h, but it required 12 h for it to reach a pH below 5.75 (Table 2).

Isolate 16 (*W. paramesenteroides*) was isolated from cheese samples (Southern Pará) which could justify its strong acidification properties given the optimal adaptation it has undergone. In addition, we can conclude that this isolate may play an important role in cheese. Not only did it show positive results for acidification and pH reduction in milk, but it also showed promising results for diacetyl production (which generates aroma), proteolytic activity (texture) and homogeneous clot formation.

4. Conclusion

This study shows the genetic diversity of *Weissella* strains isolated from different sources and different Brazilian regions that produce artisanal cheeses. It demonstrates the relevance of characterizing molecular diversity of autochthonous *Weissella* in the various environments. The technological characterizations helped understand the role *Weissella* strains play in artisanal cheese microbiota and demonstrated their potential features. Further studies will be conducted to elucidate the contributing of *Weissella* species on cheese quality and sensory aspects, allowing propose their use as starter cultures in the dairy industry.

5. Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.lwt.2021.111474>.

6. References

- Alves, M. P., Salgado, R. L., Eller, M. R., Vidigal, P. M. P., & Carvalho, A. F. (2016). Characterization of a heat-resistant extracellular protease from *Pseudomonas fluorescens* 07A shows that low temperature treatments are more effective in deactivating its proteolytic activity. *Journal of Dairy Science*, *99*(10), 7842–7851. <https://doi.org/10.3168/jds.2016-11236>
- Arena, M. P., Silvain, A., Normanno, G., Grieco, F., Drider, D., Spano, G., et al. (2016). Use of *Lactobacillus plantarum* strains as a bio-control strategy against food-borne pathogenic microorganisms. [Original Research]. *Frontiers in Microbiology*, *7*(464). <https://doi.org/10.3389/fmicb.2016.00464>
- Ayeni, F. A., S´anchez, B., Adeniyi, B. A., de los Reyes-Gavil´an, C. G., Margolles, A., & Ruas-Madiedo, P. (2011). Evaluation of the functional potential of *Weissella* and *Lactobacillus* isolates obtained from Nigerian traditional fermented foods and cow’s intestine. *International Journal of Food Microbiology*, *147*(2), 97–104. <https://doi.org/10.1016/j.ijfoodmicro.2011.03.014>
- Björkroth, J., Dicks, L. M. T., & Holzapfel, W. H. (2009). Genus III. *Weissella*. In P. De Vos, G. M. Garrity, D. Jones, N. R. Krieg, W. Ludwig, F. A. Rainey, et al. (Eds.), *Vol. 3. Bergey’s manual of systematic bacteriology, the firmicutes* (pp. 643–654). New York, NY: Springer.
- Bounaix, M.-S., Robert, H., Gabriel, V., Morel, S., Remaud-Siméon, M., Gabriel, B., et al. (2010). Characterization of dextran-producing *Weissella* strains isolated from sourdoughs and evidence of constitutive dextransucrase expression. *FEMS Microbiology Letters*, *311*(1), 18–26. <https://doi.org/10.1111/j.1574-6968.2010.02067.x>
- Cavicchioli, V. Q., Camargo, A. C., Todorov, S. D., & Nero, L. A. (2017). Novel bacteriocinogenic *Enterococcus hirae* and *Pediococcus pentosaceus* strains with antilisterial activity isolated from Brazilian artisanal cheese. *Journal of Dairy Science*, *100*(4), 2526–2535. <https://doi.org/10.3168/jds.2016-12049>
- Cavicchioli, V. Q., Dornellas, W. S., Perin, L. M., Pieri, F. A., Franco, B. D. G. M., Todorov, S. D., et al. (2015). Genetic diversity and some aspects of antimicrobial

activity of lactic acid bacteria isolated from goat milk. *Applied Biochemistry and Biotechnology*, 175(6), 2806–2822. <https://doi.org/10.1007/s12010-015-1511-8>

Chen, C., Chen, X., Jiang, M., Rui, X., Li, W., & Dong, M. (2014). A newly discovered bacteriocin from *Weissella hellenica* D1501 associated with Chinese Dong fermented meat (Nanx Wudl). *Food Control*, 42, 116–124. <https://doi.org/10.1016/j.foodcont.2014.01.031>

Choi, H., Kim, Y.-W., Hwang, I., Kim, J., & Yoon, S. (2012). Evaluation of *Leuconostoc citreum* HO12 and *Weissella koreensis* HO20 isolated from kimchi as a starter culture for whole wheat sourdough. *Food Chemistry*, 134(4), 2208–2216. <https://doi.org/10.1016/j.foodchem.2012.04.047>

Chuat, V., & Dalmasso, M. (2015). Pulsed-Field gel electrophoresis for *Leuconostoc mesenteroides* and *L. Pseudomesenteroides*. In K. Jordan, & M. Dalmasso (Eds.), *Pulse field gel Electrophoresis: Methods and protocols* (pp. 241–251). New York, NY: Springer (New York).

Cogan, T. M., Barbosa, M., Beuvier, E., Bianchi-Salvadori, B., Cocconcelli, P. S., Fernandes, I., ... Rodriguez, E. V. A. (1997). Characterization of the lactic acid bacteria in artisanal dairy products. *Journal of Dairy Research*, 64(3), 409–421. <https://doi.org/10.1017/S0022029997002185>

Collins, M. D., Samelis, J., Metaxopoulos, J., & Wallbanks, S. (1993). Taxonomic studies on some *Leuconostoc*-like organisms from fermented sausages: Description of a new genus *Weissella* for the *Leuconostoc paramesenteroides* group of species. *Journal of Applied Bacteriology*, 75(6), 595–603. <https://doi.org/10.1111/j.1365-2672.1993.tb01600.x>

Coppola, S., Fusco, V., Andolfi, R., Aponte, M., Blaiotta, G., Ercolini, D., et al. (2006). Evaluation of microbial diversity during the manufacture of Fior di Latte di Agerola, a traditional raw milk pasta-filata cheese of the Naples area. *Journal of Dairy Research*, 73(3), 264–272. <https://doi.org/10.1017/S0022029906001804>

Dal Bello, B., Cocolin, L., Zeppa, G., Field, D., Cotter, P. D., & Hill, C. (2012). Technological characterization of bacteriocin producing *Lactococcus lactis* strains employed to control *Listeria monocytogenes* in Cottage cheese. *International Journal*

of *Food Microbiology*, 153(1), 58–65. <https://doi.org/10.1016/j.ijfoodmicro.2011.10.016>

Dal Bello, B., Rantsiou, K., Bellio, A., Zeppa, G., Ambrosoli, R., Civera, T., et al. (2010). Microbial ecology of artisanal products from North West of Italy and antimicrobial activity of the autochthonous populations. *Lebensmittel-Wissenschaft und -Technologie- Food Science and Technology*, 43(7), 1151–1159. <https://doi.org/10.1016/j.lwt.2010.03.008>

Espeche, M. C., Otero, M. C., Sesma, F., & Nader-Macias, M. E. F. (2009). Screening of surface properties and antagonistic substances production by lactic acid bacteria isolated from the mammary gland of healthy and mastitic cows. *Veterinary Microbiology*, 135(3), 346–357. <https://doi.org/10.1016/j.vetmic.2008.09.078>

Felske, A., Rheims, H., Wolterink, A., Stackebrandt, E., & Akkermans, A. D. L. (1997). Ribosome analysis reveals prominent activity of an uncultured member of the class Actinobacteria in grassland soils. *Microbiology*, 143(9), 2983–2989. <https://doi.org/10.1099/00221287-143-9-2983>

Franciosi, E., Settanni, L., Cavazza, A., & Poznanski, E. (2009). Biodiversity and technological potential of wild lactic acid bacteria from raw cows' milk. *International Dairy Journal*, 19(1), 3–11. <https://doi.org/10.1016/j.idairyj.2008.07.008>

Freitas, R., Chuat, V., Madec, M.-N., Nero, L. A., Thierry, A., Valence, F., et al. (2015). Biodiversity of dairy *Propionibacterium* isolated from dairy farms in Minas Gerais, Brazil. *International Journal of Food Microbiology*, 203, 70–77. <https://doi.org/10.1016/j.ijfoodmicro.2015.03.006>

Fuka, M. M., Wallisch, S., Engel, M., Welzl, G., Havranek, J., & Schloter, M. (2013). Dynamics of bacterial communities during the ripening process of different Croatian cheese types derived from raw Ewe's milk cheeses. *PloS One*, 8(11), Article e80734. <https://doi.org/10.1371/journal.pone.0080734>

Fusieger, A., Martins, M. C. F., Freitas, R., Nero, L. A., & Carvalho, A. F. (2020). Technological properties of *Lactococcus lactis* subsp. *lactis* bv. *diacetylactis* obtained from dairy and non-dairy niches. *Brazilian Journal of Microbiology*, 51(1), 313–321. <https://doi.org/10.1007/s42770-019-00182-3>

- Gerasi, E., Litopoulou-Tzanetaki, E., & Tzanetakis, N. (2003). Microbiological study of Manura, a hard cheese made from raw ovine milk in the Greek island Sifnos. *International Journal of Dairy Technology*, 56(2), 117–122. <https://doi.org/10.1046/j.1471-0307.2003.00085.x>
- Jans, C., Bugnard, J., Njage, P. M. K., Lacroix, C., & Meile, L. (2012). Lactic acid bacteria diversity of African raw and fermented camel milk products reveals a highly competitive, potentially health-threatening predominant microflora. *Lebensmittel-Wissenschaft und -Technologie- Food Science and Technology*, 47(2), 371–379. <https://doi.org/10.1016/j.lwt.2012.01.034>
- Jay, J. M. (1996). Microorganisms in fresh ground meats: The relative safety of products with low versus high numbers. *Meat Science*, 43, 59–66. [https://doi.org/10.1016/0309-1740\(96\)00055-1](https://doi.org/10.1016/0309-1740(96)00055-1)
- Leong, K.-H., Chen, Y.-S., Lin, Y.-H., Pan, S.-F., Yu, B., Wu, H.-C., et al. (2013). Weissellicin L, a novel bacteriocin from sian-sianzih-isolated *Weissella hellenica* 4-7. *Journal of Applied Microbiology*, 115(1), 70–76. <https://doi.org/10.1111/jam.12218>
- Leroy, F., & De Vuyst, L. (2004). Lactic acid bacteria as functional starter cultures for the food fermentation industry. *Trends in Food Science & Technology*, 15(2), 67–78. <https://doi.org/10.1016/j.tifs.2003.09.004>
- López-Hernández, M., Rodríguez-Alegría, M. E., López-Munguía, A., & Wachter, C. (2018). Evaluation of xylan as carbon source for *Weissella* spp., a predominant strain in pozol fermentation. *Lebensmittel-Wissenschaft und -Technologie- Food Science and Technology*, 89, 192–197. <https://doi.org/10.1016/j.lwt.2017.10.030>
- Lynch, K. M., Lucid, A., Arendt, E. K., Sleator, R. D., Lucey, B., & Coffey, A. (2015). Genomics of *Weissella cibaria* with an examination of its metabolic traits. *Microbiology*, 161(4), 914–930. <https://doi.org/10.1099/mic.0.000053>
- Lynch, K. M., McSweeney, P. L. H., Arendt, E. K., Uniacke-Lowe, T., Galle, S., & Coffey, A. (2014). Isolation and characterisation of exopolysaccharide-producing *Weissella* and *Lactobacillus* and their application as adjunct cultures in Cheddar cheese. *International Dairy Journal*, 34(1), 125–134. <https://doi.org/10.1016/j.idairyj.2013.07.013>

Martins, M. C. F., Freitas, R., Deuvaux, J. C., Eller, M. R., Nero, L. A., & Carvalho, A. F. (2018). Bacterial diversity of artisanal cheese from the Amazonian region of Brazil during the dry and rainy seasons. *Food Research International*, *108*, 295–300. <https://doi.org/10.1016/j.foodres.2018.03.060>

Martins, M. C. F., Fusieger, A., Freitas, R., Valence, F., Nero, L. A., & Carvalho, A. F. (2020). Novel sequence types of *Lactococcus lactis* subsp. *lactis* obtained from Brazilian dairy production environments. *Lebensmittel-Wissenschaft und -Technologie- Food Science and Technology*, *124*, 109146. <https://doi.org/10.1016/j.lwt.2020.109146>

Masoud, W., Vogensen, F. K., Lillevang, S., Abu Al-Soud, W., Sørensen, S. J., & Jakobsen, M. (2012). The fate of indigenous microbiota, starter cultures, *Escherichia coli*, *Listeria innocua* and *Staphylococcus aureus* in Danish raw milk and cheeses determined by pyrosequencing and quantitative real time (qRT)-PCR. *International Journal of Food Microbiology*, *153*(1), 192–202. <https://doi.org/10.1016/j.ijfoodmicro.2011.11.014>

Masuda, Y., Zendo, T., Sawa, N., Perez, R. H., Nakayama, J., & Sonomoto, K. (2012). Characterization and identification of weissellicin Y and weissellicin M, novel bacteriocins produced by *Weissella hellenica* QU 13. *Journal of Applied Microbiology*, *112*(1), 99–108. <https://doi.org/10.1111/j.1365-2672.2011.05180.x>

Medeiros, R. S., Araújo, L. M., Queiroga Neto, V., Andrade, P. P., Melo, M. A., & Gonçalves, M. M. B. P. (2016). Identification of lactic acid bacteria isolated from artisanal Coalho cheese produced in the Brazilian Northeast. *CyTA - Journal of Food*, *14*(4), 613–620. <https://doi.org/10.1080/19476337.2016.1185468>

Meisel, H., & Bockelmann, W. (1999). Bioactive peptides encrypted in milk proteins: Proteolytic activation and thropho-functional properties. *Antonie Van Leeuwenhoek*, *76*(1), 207–215. <https://doi.org/10.1023/A:1002063805780>

Nisiotou, A., Dourou, D., Filippousi, M.-E., Banilas, G., & Tassou, C. (2014). *Weissella uvarum* sp. nov., isolated from wine grapes. *Microbiology*, *64*(Pt_11), 3885–3890. <https://doi.org/10.1099/ijfs.0.066209-0>

Ortolani, M. B. T., Yamazi, A. K., Moraes, P. M., Viçosa, G. N., & Nero, L. A. (2010). Microbiological quality and safety of raw milk and soft cheese and detection of

autochthonous lactic acid bacteria with antagonistic activity against *Listeria monocytogenes*, *Salmonella* spp., and *Staphylococcus aureus*. *Foodborne Pathogens and Disease*, 7(2), 175–180. <https://doi.org/10.1089/fpd.2009.0390>

Pakdeeto, A., Naranong, N., & Tanasupawat, S. (2003). Diacetyl of lactic acid bacteria from milk and fermented foods in Thailand. *Journal of General and Applied Microbiology*, 49(5), 301–307. <https://doi.org/10.2323/jgam.49.301>

Papagianni, M., & Papamichael, E. M. (2011). Purification, amino acid sequence and characterization of the class IIa bacteriocin weissellin A, produced by *Weissella paramesenteroides* DX. *Bioresource Technology*, 102(12), 6730–6734. <https://doi.org/10.1016/j.biortech.2011.03.106>

Pereira, C. I., San Romão, M. V., Lolkema, J. S., & Barreto Crespo, M. T. (2009). *Weissella halotolerans* W22 combines arginine deiminase and ornithine decarboxylation pathways and converts arginine to putrescine. *Journal of Applied Microbiology*, 107 (6), 1894–1902. <https://doi.org/10.1111/j.1365-2672.2009.04371.x>

Perin, L. M., & Nero, L. A. (2014). Antagonistic lactic acid bacteria isolated from goat milk and identification of a novel nisin variant *Lactococcus lactis*. *BMC Microbiology*, 14(1), 36. <https://doi.org/10.1186/1471-2180-14-36>

Perin, L. M., Savo Sardaro, M. L., Nero, L. A., Neviani, E., & Gatti, M. (2017). Bacterial ecology of artisanal Minas cheeses assessed by culture-dependent and -independent methods. *Food Microbiology*, 65, 160–169. <https://doi.org/10.1016/j.fm.2017.02.005>

Robinson, R. K. (2002). *Dairy microbiology handbook* (3rd ed.). New York, NY: Wiley-Interscience. Russell, J. B. (1992). Another explanation for the toxicity of fermentation acids at low pH: Anion accumulation versus uncoupling. *Journal of Applied Bacteriology*, 73(5), 363–370. <https://doi.org/10.1111/j.1365-2672.1992.tb04990.x>

Smit, G., Smit, B. A., & Engels, W. J. M. (2005). Flavour formation by lactic acid bacteria and biochemical flavour profiling of cheese products. *FEMS Microbiology Reviews*, 29 (3), 591–610. <https://doi.org/10.1016/j.fmre.2005.04.002> (JFEMSMicrobiology Reviews).

Srionnual, S., Yanagida, F., Lin, L.-H., Hsiao, K.-N., & Chen, Y.-s. (2007). Weissellicin 110, a newly discovered bacteriocin from *Weissella cibaria* 110, isolated from plaasom, a fermented fish product from Thailand. *Applied and Environmental Microbiology*,

73(7), 2247–2250. <https://doi.org/10.1128/AEM.02484-06> (J Applied and Environmental Microbiology).

Tamime, A. Y., & Robinson, R. K. (2007). *Tamime and Robinson's yoghurt - science and technology* (3rd ed.).

Boca Raton, FL: Woodhead Publishing. Tenover, F. C., Arbeit, R. D., Goering, R. V., Mickelsen, P. A., Murray, B. E., Persing, D. H., et al. (1995). Interpreting chromosomal DNA restriction patterns produced by pulsed-field gel electrophoresis: Criteria for bacterial strain typing. *Journal of Clinical Microbiology*, 33(9), 2233–2239.

Vijayakumar, P. P., & Muriana, P. M. (2015). A microplate growth inhibition assay for screening bacteriocins against *Listeria monocytogenes* to differentiate their mode-of-action. *Biomolecules*, 5(2), 1178–1194.

Wouters, J. T. M., Ayad, E. H. E., Hugenholtz, J., & Smit, G. (2002). Microbes from raw milk for fermented dairy products. *International Dairy Journal*, 12(2), 91–109. [https://doi.org/10.1016/S0958-6946\(01\)00151-0](https://doi.org/10.1016/S0958-6946(01)00151-0)

Zannini, E., Mauch, A., Galle, S., Ganzle, M., Coffey, A., Arendt, E. K., ... Waters, D. M. (2013). Barley malt wort fermentation by exopolysaccharide-forming *Weissella cibaria* MG1 for the production of a novel beverage. *Journal of Applied Microbiology*, 115(6), 1379–1387. <https://doi.org/10.1111/jam.12329>

CHAPTER III: *Weissella cibaria* W25:
Comparative genomic analyses

Genomic analyses of *Weissella cibaria* W25, a potential bacteriocin-producing strain isolated from pasture in Campos das Vertentes, Minas Gerais, Brazil

The content of this part has been published in:

Microorganisms

Camila Gonçalves Teixeira, Rafaela da Silva Rodrigues, Ricardo Seiti Yamatogi, Anca Lucau-Danila, Djamel Drider, Luís Augusto Nero, Antônio Fernandes de Carvalho

License to reuse this content in thesis is in Annex 4

Abstract

Weissella is a genus containing Gram-positive, heterofermentative bacteria belonging to lactic acid bacteria (LAB) group. These bacteria are endowed with promising technological and antimicrobial attributes. *Weissella cibaria* W25 was isolated from a dairy environment where raw milk cheeses are produced. Therefore, we sequenced and assembled the W25 draft genome sequence, which consists of 41 contigs totaling ~2.4Mbp, with a G+C content of 45.04%. Then we carried out comprehensive a comparative genomic analysis with *W. cibaria* 110, known to produce the weissellicin 110 bacteriocin, and four other non-bacteriocin-producing *W. cibaria* strains.

Keywords: *Weissella*; genome sequencing; bacteriocins genes

1. Introduction

The study of microbial diversity in dairy and non-dairy environments plays a pivotal role in understanding the presence of these microorganisms in such ecosystems and their impact on the final product, especially when we refer to traditional and artisanal products. Each environment has unique and specific characteristics that favor and allow the development of different bacterial species [1].

Handmade cheeses and raw milk are considered potential sources of new strains of LAB [2]. The way these cheeses are made can determine the fermentation to be conducted by bacteria present in the grazing, animal skin, utensils, surfaces and other places that may come into contact with the cheese during production [3]. The study of the bacterial community present in artisanal cheeses revealed the presence of species that had not yet been related to cheeses and a high diversity of lactic bacteria with differentiated technological characteristics [4]. In addition, non-dairy environments such as grass, different types of silage and even animal skin have also been important sources of novel strains that have adapted and therefore, can provide interesting features to be explored [5].

The diversity of *Weissella* isolated from dairy and non-dairy environments is of great interest for the enrichment in knowledge of this microorganism in the final products. The genus *Weissella* is composed of bacteria classified as Gram-positive, catalase negative, non-spore forming, coccoid morphology or short bacilli. They belong to the group of LAB, mainly due its production of lactic acid from the fermentation of carbohydrates [6]. The main purpose of this study is to announce and analyze the sequencing and annotation of *Weissella cibaria* W25 genome and carry out a comprehensive comparative genomic analysis with *W. cibaria* 110, known to produce the weissellicin 110 bacteriocin, and four other non-bacteriocin-producing *W. cibaria* strains.

2. Materials and Methods

Bacterial strain

The strain W25 was previously isolated from pasture sampled from a dairy farm located in the Campos da Vertentes region, in the Southeast of the Minas Gerais state, Brazil. This strain was named as “isolate id 25” by Teixeira, et al. [7], identified as

Weissella cibaria after sequencing of the gene 16S rRNA and characterized as possessing a technological potential due to the ability of coagulating milk, producing diacetyl and non-proteolytic. Of note, the strain presented antimicrobial activity against a panel of Gram-positive and Gram-negative foodborne pathogens [7].

Genome sequencing and assembly

The whole genome of *W. cibaria* W25 was sequenced with Nextera technology by a whole-genome shotgun strategy using the MiSeq v3 machine and Solexa platform (Illumina, USA) by Neoprosperta (Florianópolis, SC, Brazil). The trimming was performed using the Trimmomatic v.0.36 [8] and the Phred value > 20. The raw read files were trimmed of adapter sequences, and low-quality reads. After trimming, sequence reads were checked for quality using the fastQC v.0.11.5 [9] and then used for de novo genome assembly. Genome assembly was conducted by using MIRA Assembler v.4.9.6 [10], mode “genome, accurate”. The assembling quality was determined with QUILT v.5.0.2 [11], and ContEst16S was used to check contamination [12].

Genome annotation and analysis

Gene prediction and annotation were performed using the Rapid Prokaryotic Genome Annotation (PROKKA) v.1.14.5 [13], executed with default parameters and also performed by RAST automated web server [14]. To identify secondary metabolite bio-synthetic gene clusters and bacteriocins, we used the antiSMASH v.6.0 [15] and the BAGEL4 webserver [16]. Moreover, the research for plasmid was evaluated by plasmidFinder [17] and the web tool PathogenFinder [18] was used to check the presence of potential virulence factors.

Phylogenetic analyses

The identification of the genus and species was carried out using KmerFinder [19,20] and the Type (Strains) Genome Server (TYGS) [21]. The phylogenetic trees were visualized and edited using the online tool iTol v.6 (<https://doi.org/10.1093/nar/gkw290>).

Comparative genomic

For establishing the relationship between *W. cibaria* W25 and other members of this species in the bacteriocin production context we selected four published non-bacteriocins-producing *W. cibaria* strains and *W. cibaria* 110 known to produce the weissellicin 110 bacteriocin (Table 1). The genomes obtained from the GenBank were annotated using Prokka before subjecting to analyses in order to standardize the annotations.

All the strains were submitted to the Type (Strains) Genome Server (TYGS) to confirm genus and species. Moreover, for establish the genetic similarity between all strains, analysis was done with Digital DDH (DNA–DNA hybridization) similarities based on the GGDC (Genome-to-Genome Distance Calculator) web server, version 3.0 [22]. The core genome of each group was determined with OrthoVenn2 (e-value of 10⁻⁵) [23] and the CGView Server [24] was used to comparative genome analysis using BLAST with default parameters.

Availability of nucleotide sequence data

This Whole Genome Shotgun project was deposited at DDBJ/ENA/GenBank under the accession JAFNKE000000000. The version described in this paper is version JAFNKE010000000. The raw sequencing data were submitted to Sequence Read Archive (SRA) database under accession number SRR16076638.

3. Results and Discussion

Genome sequencing, annotation and analysis

The genome features of *W. cibaria* W25, *W. cibaria* 110, *W. cibaria* B3b, *W. cibaria* ffPR, *W. cibaria* JCM 12495 and *W. cibaria* MG1 are presented in table 1. The whole-genome sequencing of *W. cibaria* W25 resulted in maximum size of reads for the forward sequence of 305 and for the reverse sequence of 205 and with a total number of sequences of 2,906,916 bp. After genome assembly using the MIRA software, we obtained a draft genome with 41 contigs, N50 202,649 bp and maximum length of 331,445 bp (contigs over 500 bases).

The genome of *W. cibaria* W25 contains 2,412,435 bp, which is very similar to the genome of *W. cibaria* MG1, slightly bigger than *W. cibaria* 110, *W. cibaria* ff3PR

and *W. cibaria* JCM 12495. However, the GC content are very similar between all of them varying from 44.7 to 45.1%. Genome annotation using Prokka identified a total of 2,190 of coding DNA sequences (CDS) which were more abundant than in the genome of *W. cibaria* JCM 12495 and less abundant than other genomes. *W. cibaria* W25 presented the highest quantity of tRNA and rRNA with 11 copies of 5S ribosomal RNA (rRNA) genes, 3 copies of 16S and one single copy of 23S rRNA genes.

Table 1. Comparison of the genomic feature of *W. cibaria* W25 with a bacteriocin-producing *W. cibaria*, strains 110, and four other non-bacteriocin-producing strains.

Genome feature	W25	110	AB3b	ff3PR	JCM 12495 ^T	MG1
Accession ^a	<u>JAFNKE000</u>	<u>LRRC00</u>	<u>JWHT000</u>	<u>JWHV0000</u>	<u>BJEF00000</u>	<u>JWHU000</u>
	<u>000000</u>	<u>000000</u>	<u>00000</u>	<u>0000</u>	<u>000</u>	<u>00000</u>
Reference	This study	[25]	[26]	[26]	[27]	[26]
Contigs	41	18	88	60	25	44
Size (pb)	2,412,435	2,347,049	2,465,158	2,357,128	2,323,953	2,436,232
GC content (%) ^b	45.04	44.9	44.7	44.9	45.1	44.7
CDS	2,190	2,209	2,348	2,228	2,124	2,284
tRNA	84	76	67	70	77	62
rRNA	15	5	7	5	9	4
tmRNA	1	1	1	1	1	1
Repeat region	1	1	0	0	0	0

^a GenBank accession number; ^b using RAST program.

No plasmid gene was detected by plasmidFinder in *W. cibaria* W25 and this input organism was predicted as non-human pathogen by PathogenFinder. These results are an indicative for the safety use of this strain for future human consumption as a probiotic or as a bioprotective culture in food, for example.

According to AntiSMASH *W. cibaria* W25 possess two putative bacteriocins genes clusters, one lassopeptide (MicJ25) and one RiPP-like bacteriocin_Ilc. Of, note, the Bagel4 software did not allow identification of any bacteriocin gene. Previously, we showed that *W. cibaria* W25 has a narrow spectrum of inhibition against the most common foodborne pathogens [7] reinforcing that this strain is a suitable probiotic candidate. Li et al. (2017)[25] showed that *W. cibaria* 110 present similar results showing a large spectrum of inhibition against other LAB but the bacteriocin weissellicin 110, produced by *W. cibaria* 110, unlike most class II bacteriocins, has no inhibitory activity against *Listeria monocytogenes*.

1.1. Phylogenetic analyses and comparative genomic

Weissella phylogeny was constructed using available 16S rRNA sequences of se-sequenced *Weissella* species, including complete and draft genomes (Figure 1) and using the whole genome sequences (Figure 2). Figure 1 and 2 shows the formation of two major clusters, using only the 16S rRNA sequence we noticed that the cluster 1 comprised 4 strains (*W. confusa*, *W. cibaria* and *W. muntiaci*) and the cluster 2 comprises the other strains. On the other hand when we used the whole genome sequence, the cluster 1 comprised *W. ceti* and *W. diestrammenae* and the cluster 2 all the other strains. Also figures 1 and 2 show that *W. cibaria* W25 and *W. cibaria* JCM 12495 are phylogenetically closely related. This results along with the one from KmerFinder software confirmed the genus *Weissella* and the species *cibaria* for the strain W25 as previously announced by Teixeira et. al (2021).




Figure 1. Tree inferred with FastME 2.1.6.1[28] from GBDP distances calculated from 16S rDNA gene sequences. The branch lengths are scaled in terms of GBDP distance formula d_5 . The numbers above branches are GBDP pseudo-bootstrap support values > 60 % from 100 replications, with an average branch support of 67.6 %. The tree was rooted at the midpoint [29].




Figure 2. Tree inferred with FastME 2.1.6.1 [28] from GBDP distances calculated from genome sequences. The branch lengths are scaled in terms of GBDP distance formula d_5 . The numbers above branches are GBDP pseudo-bootstrap support values > 60 % from 100 replications, with an average branch support of 55.9 %. The tree was rooted at the midpoint [29].

The digital DDH genomic similarity revealed that between the strains tested in this study *W. cibaria* W25 possess more similarity with *W. cibaria* JCM 12495 (87.10%), a non-bacteriocin producer, and less similarity with *W. cibaria* 110 (63.80%), a strain known as a bacteriocin producer.

According to the Venn diagram from OrthoVenn2, *W. cibaria* W25 shares the same quantity of protein cluster genes with *W. cibaria* 110 and *W. cibaria* JCM 12495 (Figure 3) when comparing just the three of them. When we compared with all the strains used in this study, *W. cibaria* W25 shares more unique protein cluster genes [18] with *W. cibaria* 110 (Figure 4, B) than with the others. Besides that, *W. cibaria* W25 shares 25 with all the non-bacteriocin-producing strains (Figure 4, C). There were 1,852 protein cluster genes conserved in all *W. cibaria* strains, and five of them are unique to *W. cibaria* W25 (Figure 4, A) with 10 paralogs. Among the five protein clusters coding genes the OrthoVenn2 identified three of them, one related to lipopolysaccharide biosynthetic process, one to O antigen biosynthetic process and one to oxidoreductase activity. Li et al. (2017)[25] also compared *W. cibaria* 110 with four other strains (*W. cibaria* MG1, *W. cibaria* AB3b, *W. cibaria* ff3PR and *W. cibaria*

KACC11862) and the comparative genomic analysis also showed the presence of unique genes that encoded the novel bacteriocin weissellicin 110 and defense system.




Figure 3. Venn diagram showing the protein coding genes or pseudogenes in each sequenced of *W. cibaria* W25, *W. cibaria* 110 and *W. cibaria* JCM 12495. Overlapped regions represent shared proteins and the numbers in the non-overlapped regions indicate the unique protein coding genes or pseudogenes.




Figure 4. Venn diagrams showing the protein coding genes or pseudogenes of six *W. cibaria* strains. Overlapped regions represent shared proteins and the numbers in the non-overlapped regions indicate the unique protein coding genes or pseudogenes. A: normal Venn diagram; B: Venn diagram highlighting the cluster between *W. cibaria* W25 and *W. cibaria* 110; C: Venn diagram highlighting the clusters between *W. cibaria* W25 and the non-bacteriocin-producing strains.

We constructed a comparative genetic map using CGview server to help demonstrate the similarity between the strains (Figure 5). We used the *W. cibaria* 110 as the reference for comparison because it is a well-known bacteriocin-producing strain. Moreover, gene confirmation for weissellicin 110 was performed by the BLASTn

program at the National Center for Biotechnology Information (NCBI) using the deposited sequence under accession number LC010242 and the predicted genes of the genome from *W. cibaria* 110. We obtained results to query coverage and percent identity of 100%. Among the regions that showed to be similar between *W. cibaria* W25 and *W. cibaria* 110, we observed that the strain under study presented a similar and more complete region, when compared to the region containing weissellicin 110, than the non-bacteriocin-producing strains (Figure 5, B). This indicates that perhaps in this region of *W. cibaria* W25 there are genes for a bacteriocin, however a distinct one.




Figure 5. Comparative genome map generated using CGView server, showing a full circular map obtained using BLAST (A) and an expanded view of the weissellicin 110 region (B). The rings in (A) and (B) green, purple, yellow, red, and brown indicate the *W. cibaria* strains W25, JCM 12495, MG1, ff3PR, and AB3b, subsequently, that were compared with the *W. cibaria* 110. The blue ring in (B) indicate the coding sequences (CDS) of the genome from *W. cibaria* 110.

4. Conclusions

According to the bioinformatics results obtained in this study *W. cibaria* W25 has great potential to be used for human consumption once it was predicted as non-human pathogen. Also, despite of *W. cibaria* W25 shows more genomic similarity with *W. cibaria* JCM 12495 (according to DDH similarly) OrthoVenn2 showed that it has its unique protein cluster genes which can be related with the bacteriocins genes

indicated by AntiSMASH confirming the possibility of producing two different bacteriocins.

5. References

1. Finnegan, W.; Yan, M.; Holden, N.M.; Goggins, J. A review of environmental life cycle assessment studies examining cheese production. *Int. J. Cycle Assess.* **2018**, *23*, 1773–1787.
2. Silva, E.O.O. e; Nespolo, C.R.; Sehn, C.P.; Pinehiro, F.C.; Stefani, L.M. Lactic acid bacteria with antimicrobial, proteolytic and lipolytic activities isolated from ovine dairy products. *Food Sci. Technol.* **2020**, *40*, 293–299.
3. Pyz-lukasik, R.; Knysz, P.; Gondek, M. Hygiene Quality and Consumer Safety of Traditional Short- and Long-Ripened Cheeses from Poland. *J. Food Qual.* **2018**, *2018*.
4. Aldrete-tapia, A.; Escobar-ramírez, C.M.; Tamplin, M.L.; Hernández-Iturriaga, M. Characterization of Bacterial Communities in Mexican Artisanal Raw Milk “Bola de Ocosingo” Cheese by High-Throughput Sequencing. *Front. Microbiol.* **2018**, *9*, 2598, doi:10.3389/fmicb.2018.02598.
5. McAuliffe, O. Symposium review: *Lactococcus lactis* from nondairy sources: Their genetic and metabolic diversity and potential applications in cheese. *J. Dairy Sci.* **2018**, *101*, 3597–3610, doi:10.3168/jds.2017-13331.
6. Collins, M.D.; Samelis, J.; Metaxopoulos, J.; Wallbanks, S. Taxonomic studies on some leuconostoc-like organisms from fermented sausages: description of a new genus *Weissella* for the *Leuconostoc paramesenteroides* group of species. *J. Appl. Bacteriol.* **1993**, *75*, 595–603, doi:10.1111/j.1365-2672.1993.tb01600.x.
7. Teixeira, C.G.; Fusieger, A.; Martins, E.; de Freitas, R.; Vakarelova, M.; Nero, L.A.; Carvalho, A. fernandes de Biodiversity and technological features of *Weissella* isolates obtained from Brazilian artisanal cheese-producing regions. *LWT - Food Sci. Technol.* **2021**, *147*, 111474, doi:10.1016/j.lwt.2021.111474.
8. Bolger, A.M.; Lohse, M.; Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **2014**, *30*, 2114–2120, doi:10.1093/bioinformatics/btu170.
9. Andrews, S. FastQC: a quality control tool for high throughput sequence data. *Available online* 2010.
10. Chevreur, B.; Wetter, T.; Suhai, S. Genome Sequence Assembly Using Trace Signals and Additional Sequence Information. In Proceedings of the Computer Science and Biology: Proceedings of the German Conference on Bioinformatics; 1999; Vol. 99, pp. 45–46.
11. Gurevich, A.; Saveliev, V.; Vyahhi, N.; Tesler, G. QUAST : quality assessment tool for genome assemblies. *Bioinformatics* **2013**, *29*, 1072–1075, doi:10.1093/bioinformatics/btt086.
12. Lee, I.; Chalita, M.; Ha, S.; Na, S.; Yoon, S.; Chun, J. ContEst16S : an algorithm that identifies contaminated prokaryotic genomes using 16S RNA gene sequences. *Int J Syst Evol Microbiol* **2017**, *67*, 2053–2057, doi:10.1099/ijsem.0.001872.

13. Seemann, T. Prokka: rapid prokaryotic genome annotation. *Bioin* **2014**, *30*, 2068–2069, doi:10.1093/bioinformatics/btu153.
14. Aziz, R.K.; Bartels, D.; Best, A.A.; Dejongh, M.; Disz, T.; Edwards, R.A.; Formsma, K.; Gerdes, S.; Glass, E.M.; Kubal, M.; et al. The RAST Server : Rapid Annotations using Subsystems Technology. *BCM Genomics* **2008**, *9*, 1–15, doi:10.1186/1471-2164-9-75.
15. Blin, K.; Shaw, S.; Kloosterman, A.M.; Charlop-Powers, Z.; van Wezel, G.P.; Medema, M.H.; Weber, T. antiSMASH 6.0: improving cluster detection and comparison capabilities. *Nucleic Acids Res.* **2021**, *49*, 0–7, doi:10.1093/nar/gkab335.
16. Heel, A.J. Van; Jong, A. De; Song, C.; Viel, J.H.; Kok, J.; Kuipers, O.P. BAGEL4 : a user-friendly web server to thoroughly mine RiPPs and bacteriocins. *Nucl. Acids Res.* **2018**, *46*, 278–281, doi:10.1093/nar/gky383.
17. Carattoli, A.; Zankari, E.; García-fernández, A.; Larsen, M.V.; Lund, O.; Villa, L.; Aarestrup, F.M.; Hasman, H. In Silico Detection and Typing of Plasmids using PlasmidFinder and Plasmid Multilocus Sequence Typing. *Antimicrob. Agents Chemother.* **2014**, *58*, 3895–3903, doi:10.1128/AAC.02412-14.
18. Cosentino, S.; Larsen, M.V.; Aarestrup, F.M.; Lund, O. PathogenFinder - Distinguishing Friend from Foe Using Bacterial Whole Genome Sequence Data. *Plus One* **2013**, *8*, e77302, doi:10.1371/journal.pone.0077302.
19. Hasman, H.; Saputra, D.; Sicheritz-ponten, T.; Lund, O.; Svendsen, C.A.; Frimodt-møller, N. Rapid Whole-Genome Sequencing for Detection and Characterization of Microorganisms Directly from Clinical Samples. *J. Clin. Microbio* **2014**, *52*, 139–146, doi:10.1128/JCM.02452-13.
20. Larsen, M. V; Cosentino, S.; Lukjancenko, O.; Saputra, D.; Rasmussen, S.; Hasman, H.; Sicheritz-pontén, T.; Aarestrup, F.M.; Ussery, D.W.; Lund, O. Benchmarking of Methods for Genomic Taxonomy. *J. Clin. Microbiol.* **2014**, *52*, 1529–1539, doi:10.1128/JCM.02981-13.
21. Meier-kolthoff, J.P.; Göker, M. TYGS is an automated high-throughput platform for state-of-the-art genome-based taxonomy. *Nat. Commun.* **2019**, *10*, 2182, doi:10.1038/s41467-019-10210-3.
22. Meier-kolthoff, J.P.; Carbasse, J.S.; Peinado-Olarte, R.L.; Göker, M. TYGS and LPSN : a database tandem for fast and reliable genome-based classification and nomenclature of prokaryotes. *Nucleic Acids Res.* **2021**, 1–7.
23. Xu, L.; Dong, Z.; Fang, L.; Luo, Y.; Wei, Z.; Guo, H.; Zhang, G.; Gu, Y.Q.; Coleman-Derr, D.; Xia, Q.; et al. OrthoVenn2 : a web server for whole-genome comparison and annotation of orthologous clusters across multiple species. *Nucl. Acids Res.* **2019**, *47*, 52–58, doi:10.1093/nar/gkz333.
24. Stothard, P.; Wishart, D.S. Circular genome visualization and exploration using CGView. *Bioinformatics* **2005**, *21*, 537–539, doi:10.1093/bioinformatics/bti054.
25. Li, S.; Chen, Y.; Lee, Y.; Yang, C.; Srionnual, S.; Wu, H.; Chang, C. Comparative genomic analysis of bacteriocin-producing *Weissella cibaria* 110. *Appl. Microbiol. Biotechnol.* **2017**, *101*, 1227–1237, doi:10.1007/s00253-016-8073-8.

26. Lynch, K.M.; Lucid, A.; Arendt, E.K.; Sleator, R.D.; Lucey, B.; Coffey, A. Genomics of *Weissella cibaria* with an examination of its metabolic traits. *Microbiology* **2015**, *161*, 914–930, doi:10.1099/mic.0.000053.
27. Björkroth, K.J.; Schillinger, U.; Geisen, R.; Weiss, N.; Hoste, B.; Holzapfel, W.H.; Korkeala, H.J.; Vandamme, P. Taxonomic study of *Weissella confusa* and description of *Weissella cibaria* sp. nov., detected in food and clinical samples. *Int. J. Syst. Evol. Microbiol.* **2002**, *52*, 141–148, doi:10.1099/00207713-52-1-141.
28. Lefort, V.; Desper, R.; Gascuel, O. FastME 2.0: A Comprehensive, Accurate, and Fast Distance-Based Phylogeny Inference Program. *Mol. Biol. Evol.* **2015**, *32*, 2798–2800, doi:10.1093/molbev/msv150.
29. Farris, J.S. Estimating Phylogenetic Trees from Distancec Matrices. *Am. Nat.* **1972**, *106*, 645–668.

CHAPTER IV: *Weissella cibaria* strains isolated from Campo das Vertentes (MG), a potential bacteriocin producers

Genome characterization of *Weissella* strains isolated from Campos das Vertentes, Minas Gerais, Brazil focused revealed new bacteriocins with a large spectrum of activity

Camila Gonçalves Teixeira, Rafaela da Silva Rodrigues, Anca Lucau-Danila, Luís Augusto Nero, Antônio Fernandes de Carvalho, Djamel Drider

Abstract

The ability of lactic acid bacteria (LAB) to produce different antimicrobial compounds, like bacteriocins, that could be used in the food and pharmaceutical industries. LAB are commonly found in the gut microbiota and in natural fermented foods, and some LAB possess the “Generally Recognized as Safe” status. Bacteriocins can be used as alternative to aging antibiotics, and numerous studies are steadily reporting the use of bacteriocins as a novel therapeutic options to fight antibiotic-resistant pathogens. On the other hand, bacteriocins, mainly those produced by LAB referred as LAB-bacteriocins can largely replce chemical preservatives in the food industry.

This work aimed at characterizing the *Weissella cibaria* strains isolated from Campos das Vertentes, Minas Gerais, Brazil focusing on their antimicrobial compounds. To this end, three strains of *W. cibaria* (W21, W25, and W42) were previously selected, and the genomes of W21 and W42 were sequenced, while that of strain W25 was already sequenced (Teixeira et al., 2022). Afterwhich, we investigated the presence of genes coding for potential secondary metabolites in the genomes of the strains using the AntiSmash webtool, and discovered that they have a common region in the genome with a putative bacteriocin identified as Bacteriocin_IIc. Subsequently, we assessed experimentally *in vitro* this antimicrobial activity using the supernatant of these strains. According to the data, the strain W21, which has one transport-related gene missing from the bacteriocin operon comparing to the others two, likewise showed no differences in the *in vitro* antimicrobial test when comparing the outcomes of the three treatments (supernatant without treatment, neutralized and treated with proteinase K). Also, we compared the nucleotides and amino acids of this putative bacteriocin in the NCBI platform and there was no similarity with other so far described bacteriocin from the genus *Weissella*. In conclusion, two of the three *Weissella* strains tested in this

work have the potential to produce a new bacteriocin not yet described which needs to be deeply characterized and proposed for food protection and/or therapeutic options as alternatives to chemicals in foods or aging antibiotics, respectively.

Key words: *Weissella cibaria*, bacteriocins, antimicrobial activity

1. Introduction

For many years antibiotics have been used as the basis of treatment against bacterial infections and their misuse and exacerbated the development of antibiotic-resistant species (Blair et al., 2015). The World Health Organization (WHO) encourages new strategies and interventions to avoid the development of new multi-drug-resistant pathogens (Neill, 2016). LAB and their diverse metabolites such as organic acids, hydrogen peroxide (H₂O₂), and bacteriocins offer interesting prospects (Hernández-González et al., 2021).

Bacteriocins are ribosomally synthesized antimicrobial peptides/proteins produced by Gram-positive, Gram-negative, and Archaea (Hammami et al., 2010; Nes, 2011) with the aim to defeat their cospecific strains living in the ecological niches (Soltani et al., 2021). Bacteriocins can be active against phylogenetically related or phylogenetically distant target strains, but will not against producing strains themselves because of their developed specific immunity proteins (Drider et al., 2006). As bacteriocins are commonly produced by strains from the gut microbiota and natural food sources, in which some of them are “Generally Recognized as Safe” (GRAS status), interest for LAB-bacteriocins are ever-growing for the food and pharmaceutical sectors (Field et al., 2018). In addition, bacteriocins are non-toxic to cells from Eukarya domain and can be used against many food spoilage microbes and pathogenic bacteria (Belguesmia et al., 2011; Field et al., 2018).

Despite some LAB genera being known for their antimicrobial activity against food spoilage and pathogenic microorganisms, not all of them are effective. Some of them produce a low amount of antimicrobial compounds or have a limited spectrum of activity (Tenea et al., 2020). Therefore, the discovery of novel antimicrobial compounds endowed with broad inhibitory spectra is of interest for their potential applications as alternatives to overage antibiotics or as bio-preservative agents for the food industry. Of note, applications of LAB-bacteriocins in the food industry as bio-preservative

agents is feasible and constitute an eco-friendly approach that could replace the use of chemical preservatives. The biopreservative approach is timely as it meets the consumers expectations for natural and bioprotected foods (Venegas-Ortega et al., 2019).

Within the LAB group, strains of the *Weissella* genus are known to produce bacteriocins such as weissellicin 110, MBF, L, D, M, Y, and weissellicin A from strains of *W. cibaria*, *W. paramesenteroides* and *W. hellenica* (Chen et al., 2014; Di Cagno et al., 2011; Kariyawasam et al., 2019; Leong et al., 2013; Masuda et al., 2012; Srionnual et al., 2007). Related to that, we isolated strains of *W. cibaria* from different artisanal cheese-producing regions in Brazil and established their technological and bacteriocinogenic traits (Teixeira et al., 2021). Most of the studied isolates demonstrated broad inhibition against several pathogens, both gram-positive and gram-negative (Teixeira et al., 2021).

The aim of this study is to identify the antimicrobial compounds produced by *Weissella* strains isolated from Campos das Vertentes, Minas Gerais, Brazil, with the aim to schedule their future application as pathogens inhibitors in the food industry and preservation.

2. Material and Methods

2.1. Microorganisms and Culture Conditions

Three different strains of *Weissella cibaria*, previously isolated from dairy farms located in the Campos da Vertentes region, in the southeast of the Minas Gerais state, Brazil were used in this work. The strains W21 (isolate id 21) and W25 (isolate id 25) were isolated from pasture samples, whereas strain W42 (isolate id 42) was isolated from soil (Teixeira et al., 2021). These 3 strains were identified as *Weissella cibaria* after sequencing of the gene 16S rRNA and presented antimicrobial activity against a panel of Gram-positive and Gram-negative foodborne pathogens (Teixeira et al., 2021). The isolates obtained were stored at -60°C in de Man Rogosa and Sharpe (MRS) broth (Oxoid Ltd., Basingstoke, England) and supplemented with 30% (v/v) glycerol (Synth, São Paulo, SP, Brazil). Before analysis, stock cultures (1% v/v) were transferred to MRS broth (Oxoid) and incubated at 30 °C overnight.

2.2. Genome Sequencing, Assembly, and Annotation

The genome sequence of the W25 strain was previously deposited in GenBank under accession number JAFNKE000000000 (Teixeira et al., 2022). The whole genomes of *W. cibaria* W21 and W42 were sequenced, trimmed, and assembled as recently performed for the W25 strain (Teixeira et al., 2022). Gene prediction and annotation were performed using the Rapid Prokaryotic Genome Annotation (PROKKA) v.1.14.5 (Seemann, 2014), executed with default parameters, and also performed by RAST automated web server (Aziz et al., 2008). The assembled genome was conducted by using MIRA Assembler v.4.9.6 (Chevreux et al., 1999), mode “genome, accurate”. The assembling quality was determined with QCAST v.5.0.2 (Gurevich et al., 2013), and ContEst16S was used to check contamination (Lee et al., 2017).

2.3. Bacterial Species Confirmation

The identification of the genus and species was carried out using KmerFinder (Hasman et al., 2014; Larsen et al., 2014) and the Type (Strains) Genome Server (TYGS) (Meier-kolthoff & Göker, 2019). The phylogenetic tree was visualized and edited using the online tool iTol v.6 (Letunic & Bork, 2016).

Genus and species were also confirmed by Matrix-assisted laser desorption/ionization-time of flight (MALDI-TOF) mass spectrometry on an Autoflex Speed™ (Bruker Daltonics, Bremen, Germany) with flex control software (Bruker Daltonics).

Protein extraction for bacterial identification was made according to Bruker's recommendation after 24 h of culture in MRS media. Briefly, bacterial pellets were washed with 70% ethanol solution, centrifuged to remove all added ethanol, and incubated for 5 min at room temperature with 50 µL of 70% formic acid, and another 5 min with an equal volume of 100% acetonitrile. After a quick step of centrifugation at maximum speed for 2 minutes, 1 µL of supernatant was spotted 3 times on a polished steel MALDI target plate (Bruker Daltonics) and dried at room temperature. Each spot was overlaid with 1 µL of matrix solution [10 mg/ml of HCCA dissolved in ACN/water/TFA (50/47.5/2.5; v/v/v)] and dried again (Heuson et al., 2019; Nacef et al., 2017; Zidour et al., 2017). MALDI-TOF mass spectrometer calibration was performed using the bacterial standard test recommended by Bruker. MALDI-MS profiles were

acquired in positive linear mode across the m/z range of 2–20 kDa using the manufacturer's automatic method MBT_FC.par (Bruker Daltonics).

2.4. Antimicrobial Potential

To identify secondary metabolite biosynthetic gene clusters and bacteriocins, we used the antiSMASH v.6.0 (Blin et al., 2021) and the BAGEL4 web server (Heel et al., 2018). The bacteriocin genes detected were blasted on the NCBI platform and the *in silico* prediction of promoters (-10 and -35) and terminator region for the bacteriocin coding gene was performed manually and using Arnold Finding Terminator (Gautheret et al., 2001), respectively. The putative promoter region of each gene was suggested based on the consensus sequences for the -10 box (TATAAT) and -35 box (TTGACA) and considering their position upstream of the start codon.

We tested the antimicrobial activity against *Escherichia coli* ATCC 8739, *Kocuria rhizophila* CIP 5345, *Listeria innocua* CIP 80.11, and *Salmonella enterica* using the microdilution assays according to Teixeira et al. (Teixeira et al., 2021) and the spot on the lawn assay according to Moraes et al. (Moraes et al., 2010). *Weissella* spp. strains were cultured in MRS broth (Oxoid) at 30 °C for 24 h and a cell-free supernatant (CFS) was obtained by centrifugation ($10,000 \times g$ for 10 min at 4 °C) and filtration ($\emptyset = 0.45 \mu\text{m}$). Target strains were cultured in a brain heart infusion (BHI, Oxoid) at 37 °C for 24 h. for the microdilution assay aliquots of the supernatant (100 μL , at a concentration of about 10^4 CFU/mL) were transferred to 96-well microtiter plates. Then, 50 μL aliquots of the selected *Weissella* spp. strains' CFS were added to each well. The microplates were incubated at 37 °C for 24 h in a Multiskan™ GO Microplate Spectrophotometer (Thermo Fisher Scientific, Waltham, MA, USA). The cultures' optical densities (OD) were measured every 30 min ($\lambda = 600 \text{ nm}$). For the spot-on-the-lawn assay the supernatant was ten-fold concentrated and spotted on BHI agar plate. After absorption at room temperature (10-15 min), the agar was covered with an overlayer of semi-solid BHI (8 ml) inoculated with 10^5 CFU/mL of the target and incubated at 37 °C for 24 h. The absence of an inhibition zone in the proximity of the well indicates no antimicrobial activity of the supernatant.

After confirming the antimicrobial activity against *E. coli*, *K. rhizophila*, *L. innocua*, *S. enterica*, and *S. aureus*, the supernatant of the *Weissella* strains were treated with proteinase K in order to confirm the proteinaceous nature of the inhibitory substances. On other hand, the pH of the culture supernatant was adjusted/neutralized to discard

the effect of any inhibition linked to acid production. Of note, after these treatments, the antibacterial activity was confirmed against the aforementioned target organisms with both assays as described above.

3. Results and Discussion

3.1. Genome Sequencing, Assembly, and Annotation

The information obtained after genome sequencing, assembly, and annotation is presented in table 1.

Table 1. Genome assembly and annotation statistics for *W. cibaria* W21, W25, and W42 strains.

Parameters	Strains		
	W25 ¹	W21	W42
Contigs	41	50	42
Total length (bp)	2,412,435	2,461,013	2,455,505
Largest contig	331,445	290,433	290,388
N50	202,649	141,851	159,842
N^a	40	434	424
G+C content (%)	45.04	44.96	44.95
CDS	2,190	2,244	2,237
rRNA	15	19	18
tRNA	84	87	86
tmRNA	1	1	1
Repeat region	1	1	1

¹ Results according to Teixeira et al. [19]; ^a The total number of uncalled bases in the assembly.

A slight difference between the three genomes regarding the number of contigs and the size of the genomes (Table 1). According to Björkroth et al. (Björkroth et al., 2002), the G+C content in *W. cibaria* species should be 44 to 45 mol% DNA base ratio, which was what we observed. We can also see that the quantities of rRNA and tRNA genes are comparable between the same species but slightly different due to they are different strains from the same species.

3.2. Bacterial Species Confirmation

The identification of the genus and species using KmerFinder (Hasman et al., 2014; Larsen et al., 2014) and the MALDI-TOF are in agreement with the molecular

identification based on the 16S rRNA/DNA sequencing realized by Teixeira et al. (Teixeira et al., 2021). The MALDI-TOF analysis is utilized for the identification and confirmation of the LAB bacteria genus and species based on its recognized robustness and appropriateness for LAB (Abdelkader et al., 2021; Nacef et al., 2017) including those belonging to the *Weissella* genus (Abdelkader et al., 2021; Kim et al., 2021; Ogunremi et al., 2022). Besides that, it has been proven to be a fast method of identification of bacteria at the species and genus level and can be utilized for routine analysis due to its effectiveness (Nacef et al., 2017; Zidour et al., 2017). This technique has been already used to identify of *Weissella* genus and species (Abdelkader et al., 2021; Kim et al., 2021; Ogunremi et al., 2022).

The results from the genetic relatedness of *W. cibaria* W21, W25, and W42, constructed using their whole-genomes and those from other *Weissella*, are expressed in the phylogenetic tree in figure 1.




Figure 1. Tree inferred with FastME 2.1.6.1 (Lefort et al., 2015) from GBDP distances calculated from genome sequences. The branch lengths are scaled in terms of GBDP distance formula $d5$. The numbers above branches are GBDP pseudo-bootstrap support values > 60 % from 100 replications, with average branch support of 56.6 %. The tree was rooted at the midpoint (Farris, 1972).

Figure 1 besides confirming the genus and species of the three sequenced strains also confirmed the similarity of the results from table 1 grouping the three strains in the same cluster with a high percentage of similarity.

3.3. Antimicrobial potential

In the previous report [15], we mentioned the presence of two putative bacteriocins genes cluster (Teixeira et al., 2022), one lasso peptide (MicJ25), and one RiPP-like bacteriocin_IIc. According to the analysis for *W. cibaria* W25, AntiSMASH (Blin et al., 2021) detected three regions in its genome, C6, C18, and C75.

The C6 region contains 2 core biosynthetic genes, one additional biosynthetic gene, one transport-related gene, and nine putative genes. The two core biosynthetic genes are a possible bacteriocin classified as Bacteriocin_IIc (Figure 2) and a possible protein related to the ABC transport. The additional biosynthetic gene was identified as a phenylalanine-specific permease and the transport-related gene was identified as a possible ABC transporter ATP-binding protein.

In the C18 region, AntiSMASH (Blin et al., 2021) detected one core biosynthetic gene named biosynthetic (rule-based-clusters) hydroxymethylglutaryl-CoA synthase, and one additional biosynthetic gene identified as biosynthetic-additional (rule-based-clusters) Flavoprotein, one transport-related gene identified as transport SMC0G1184: major facilitator transporter and 29 putative genes.

Finally, in the C75 there was one core biosynthetic gene named biosynthetic (rule-based clusters) lasso peptide: micJ25 and three putative genes.

In the genome of W21, we detected two regions C1 and C9 with two and one core biosynthetic genes, respectively. In the C1 region, we found, as in the W25, a possible bacteriocin gene named Bateriocin_IIc. The difference between this region compared to the C6 in the genome of the W25 strain is that there are all the same genes except for the transport-related gene, as we found in W25. Region C9 of W21 has the same genes as region C18 in the W25 genome strain.

In the genome of strain W42, AntiSMASH (Blin et al., 2021) detected two different regions and one possible bacteriocin. Region C3 has the same genes as the C6 region in the W25 genome strain. Also, the C2 region has the same genes as the region C18 in W25.

The bacteriocin_IIc detected by AntiSMASH in the region C6 of *W. cibaria* W25 is represented in figure 2.




Figure 2. In silico prediction of promoters (-10 and -35) and terminator region for the bacteriocin coding gene. Figure adapted from AntiSMASH results.

The gene for the bacteriocin_Ilc was identified in all three *Weissella* strains, and the region containing the gene has also other genes such as the ABC transport protein-related gene and other Orfs non-identified by AntiSMASH. However, the strain W21 is missing one transport-related gene. When the sequence of amino acids of this putative bacteriocin was blasted in the NCBI platform it shows a 98.48% of identity with Blp family class II bacteriocin from *W. cibaria*.

The putative sequences of the promoter region for the bacteriocin_Ilc gene are depicted in Figure 2. The results for the terminator region indicate that it is a Rho-independent terminator which is based on a palindrome sequence downstream from the target gene. The distance between the Orf05 and Orf06 is very short, (3 nucleotides) and therefore we could not identify any putative promoter sequences for gene Orf06 and terminator sequences for gene Orf05 which means that these Orfs could be co-transcribed.

There are already some well-known bacteriocins from the *Weissella* genus such as Weissellicin 110, MBF, L, D, M, Y, and Weissellicin A from strains of *W. cibaria*, *W. paramesenteroides*, and *W. hellenica* (Chen et al., 2014; Di Cagno et al., 2011; Kariyawasam et al., 2019; Leong et al., 2013; Masuda et al., 2012; Srionnual et al., 2007). The Weissellicin 110 is the most characterized bacteriocin from the specie *W. cibaria*, despite of that no other bacteriocin was well-study from the strain *W. cibaria* and the identified sequence by AntiSMASH didn not reveal any similarity with the studied bacteriocins from *Weissella* genus.

The *in vitro* evaluation of antimicrobial activity against *E. coli*, *K. rhizophila*, *L. innocua*, and *S. enterica* is presented in Figure 3 for the microdilution assay. The results of the spot-on-the-lanw assay shows that all the supernatant previous the

treatment presented a clear zone around the spot, but in the test with the treated supernatant no clear zone was detected.




Figure 3. Antimicrobial activity of *Weissella* strains supernatants (Non-treated, Neutralized, and Neutralized, and treated with proteinase K) against the targets *E. coli*, *K. rhizophila*, *L. innocua*, and *S. enterica*. The analysis was in duplicate.

The results showed that when we used the supernatant without treatment there was no growth of any target, but when the supernatant is neutralized or neutralized and treated with proteinase K there was a huge decrease in the antimicrobial activity. However, as we can see in some growth curves from figure 3 (A, C, D, F, and I) after neutralization of the supernatant there was a remaining antimicrobial activity and this one was lost after the treatment with proteinase K. This is an indication that the major compound responsible for antimicrobial activity can be some organic acids produced by the tested *Weissella* strains or a combination between produced compounds. Several compounds produced by *Weissella* strains has antimicrobial activity such as organic acids, hydrogen peroxide, bacteriocins, and other (Cheng et al., 2016; Fhoula

et al., 2022; Yeu et al., 2021; Yu et al., 2019). Also, the synergism between two or more compounds, such as lactic acid and acid citric with bacteriocin, can increase the antimicrobial activity of the supernatant (Soltani et al., 2022). Teixeira et al. (Teixeira et al., 2021) also tested the W25 and W21 strains against different targets such as *L. monocytogenes* ATCC 6538, *S. Typhimurium* ATCC 14028, *S. aureus* ATCC 6538, and *E. coli* ATCC 1122. The authors observed a higher level of inhibition in the supernatant of W21 compared to W25 and also W21 presented the highest inhibition level against *L. monocytogenes* ATCC 15313. However, in the present work comparing the same strains we can see that after the treatment with proteinase K there was no difference in the antimicrobial activity which means that the inhibitory activity from W21 is not related to the production of bacteriocins. This can be confirmed by the AntiSMASH results, once in this genome, the region that has the lasso peptide has no transport-related gene as in the genome of W25 and W42. According to AntiSMASH results, *W. cibaria* W25 and W42 possess one putative lasso peptide and both of them have the same genes in the region containing this bacteriocin. Also, these two strains are the ones in the antimicrobial test that maintained some antimicrobial activity after neutralizing and lost it after being treated with proteinase K.

4. Conclusions

The three *W. cibaria* strains studied in this work present high antimicrobial inhibitory activity of the non-treated supernatant against all the target strains tested. Besides that, the genome of all of them presents the putative gene for bacteriocin identified as bacteriocin_IIc by the AntiSMASH webserver. However, the strain *W. cibaria* W21 did not present the transport-related gene presented in the others two. This fact was confirmed by the *in vitro* test with the supernatant neutralized and with the supernatant neutralized treated with proteinase K. Once the supernatant was neutralized it lost antimicrobial activity, however when the supernatant of W25 and W42 was treated with proteinase K, the remaining antimicrobial activity was lost and for W21 there was no difference between the tested supernatants against all targets. This is an indication of the production of bacteriocin. Moreover, the bacteriocin identified has no similarity with the well-known bacteriocin produced by *W. cibaria* nor with the other strains from this genus being a potential new bacteriocin for the food and pharmaceutical industries.

5. Data Availability

Draft genome sequence of the *W. cibaria* strains W21 and W42 was submitted to NCBI under GenBank accession numbers JALRNM000000000 and JALRNN000000000, respectively.

6. Conflict of Interest

The authors confirm that they have no conflicts of interest with respect to the work described in this manuscript.

7. Acknowledgements

We are thankful for the financial support provided by the Brazilian agencies: Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq, Brasília, DF, Brazil), Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG, Belo Horizonte, MG, Brazil), and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES, Brasília, DF, Brazil, Financial code 001).

8. References

- Abdelkader, A., Cnockaert, M., Aerts, M., & Daube, G. (2021). Comparison of MALDI TOF MS profiling and 16S rRNA gene identification of presumptive lactic acid bacteria isolated from the traditional Algerian date product "Btana". *Applied Biology in Saharan Areas*, 3(3), 27–43.
- Aziz, R. K., Bartels, D., Best, A. A., Dejongh, M., Disz, T., Edwards, R. A., Formsma, K., Gerdes, S., Glass, E. M., Kubal, M., Meyer, F., Olsen, G. J., Olson, R., Osterman, A. L., Overbeek, R. A., McNeil, L. K., Paarmann, D., Paczian, T., Parrello, B., ... Zagnitko, O. (2008). The RAST Server: Rapid Annotations using Subsystems Technology. *BCM Genomics*, 9(75), 1–15. <https://doi.org/10.1186/1471-2164-9-75>
- Belguesmia, Y., Madi, A., Sperandio, D., Merieau, A., Feuilloley, M., Prévost, H., Drider, D., & Connil, N. (2011). Growing insights into the safety of bacteriocins: The case of enterocin S37. *Research in Microbiology*, 162(2), 159–163. <https://doi.org/10.1016/j.resmic.2010.09.019>
- Björkroth, K. J., Schillinger, U., Geisen, R., Weiss, N., Hoste, B., Holzappel, W. H., Korkeala, H. J., & Vandamme, P. (2002). Taxonomic study of *Weissella confusa* and description of *Weissella cibaria* sp. nov., detected in food and clinical samples. *International Journal of Systematic and Evolutionary Microbiology*, 52(1), 141–148.

<https://doi.org/10.1099/00207713-52-1-141>

- Blair, J. M. A., Webber, M. A., Baylay, A. J., Ogbolu, D. O., & Piddock, L. J. V. (2015). Molecular mechanisms of antibiotic resistance. *Nature Reviews Microbiology*, 13(January), 42–51. <https://doi.org/10.1038/nrmicro3380>
- Blin, K., Shaw, S., Kloosterman, A. M., Charlop-Powers, Z., van Wezel, G. P., Medema, M. H., & Weber, T. (2021). antiSMASH 6.0: improving cluster detection and comparison capabilities. *Nucleic Acids Research*, 49(W1), 0–7. <https://doi.org/10.1093/nar/gkab335>
- Chen, C., Chen, X., Jiang, M., Rui, X., Li, W., & Dong, M. (2014). A newly discovered bacteriocin from *Weissella hellenica* D1501 associated with Chinese Dong fermented meat (Nanx Wudl). *Food Control*, 42, 116–124. <https://doi.org/10.1016/j.foodcont.2014.01.031>
- Cheng, J. R., Liu, X. M., Chen, Z. Y., Zhang, Y. S., & Zhang, Y. H. (2016). Mulberry anthocyanin biotransformation by intestinal probiotics. *Food Chemistry*, 213(July), 721–727. <https://doi.org/10.1016/j.foodchem.2016.07.032>
- Chevreur, B., Wetter, T., & Suhai, S. (1999). Genome Sequence Assembly Using Trace Signals and Additional Sequence Information. *Computer Science and Biology: Proceedings of the German Conference on Bioinformatics*, 99(1995), 45–46.
- Di Cagno, R., Minervini, G., Rizzello, C. G., De Angelis, M., & Gobbetti, M. (2011). Effect of lactic acid fermentation on antioxidant, texture, color and sensory properties of red and green smoothies. *Food Microbiology*, 28(5), 1062–1071. <https://doi.org/10.1016/j.fm.2011.02.011>
- Drider, D., Fimland, G., Héchard, Y., McMullen, L. M., & Prévost, H. (2006). The Continuing Story of Class IIa Bacteriocins. *Microbiology and Molecular Biology Reviews*, 70(2), 564–582. <https://doi.org/10.1128/mnbr.00016-05>
- Farris, J. S. (1972). Estimating Phylogenetic Trees from Distance Matrices. *The American Naturalist*, 106(951), 645–668.
- Fhoula, I., Boumaiza, M., Tayh, G., Rehaïem, A., Klibi, N., & Ouzari, I.-H. (2022). Antimicrobial activity and safety features assessment of *Weissella* spp. from environmental sources. *Food Science and Nutrition*, 00(April), 1–15. <https://doi.org/10.1002/fsn3.2885>
- Field, D., Ross, R. P., & Hill, C. (2018). Developing bacteriocins of lactic acid bacteria into next generation biopreservatives. *Current Opinion in Food Science*, 20(April), 1–6. <https://doi.org/10.1016/j.cofs.2018.02.004>

- Gautheret, D., Lambert, A., & Case, L. (2001). Direct RNA Motif Definition and Identification from Multiple Sequence Alignments using Secondary Structure Profiles. *Journal of Molecular Biology*, 313(5), 1003–1011. <https://doi.org/10.1006/jmbi.2001.5102>
- Gurevich, A., Saveliev, V., Vyahhi, N., & Tesler, G. (2013). QUASt: quality assessment tool for genome assemblies. *Bioinformatics*, 29(8), 1072–1075. <https://doi.org/10.1093/bioinformatics/btt086>
- Hammami, R., Zouhir, A., Le Lay, C., Ben Hamida, J., & Fliss, I. (2010). BACTIBASE second release: A database and tool platform for bacteriocin characterization. *BMC Microbiology*, 10(22). <https://doi.org/10.1186/1471-2180-10-22>
- Hasman, H., Saputra, D., Sicheritz-ponten, T., Lund, O., Svendsen, C. A., & Frimodt-møller, N. (2014). Rapid Whole-Genome Sequencing for Detection and Characterization of Microorganisms Directly from Clinical Samples. *Journal of Clinical Microbiology*, 52(1), 139–146. <https://doi.org/10.1128/JCM.02452-13>
- Heel, A. J. Van, Jong, A. De, Song, C., Viel, J. H., Kok, J., & Kuipers, O. P. (2018). BAGEL4: a user-friendly web server to thoroughly mine RiPPs and bacteriocins. *Nucleic Acids Research*, 46(May), 278–281. <https://doi.org/10.1093/nar/gky383>
- Hernández-González, J. C., Martínez-Tapia, A., Lazcano-Hernández, G., García-Pérez, B. E., & Castrejón-Jiménez, N. S. (2021). Bacteriocins from Lactic Acid Bacteria. A Powerful Alternative as Antimicrobials, Probiotics, and Immunomodulators in Veterinary Medicine. *Animals*, 11(April), 979.
- Heuson, E., Etchegaray, A., Filipe, S. L., Beretta, D., Chevalier, M., Phalip, V., & Coutte, F. (2019). Screening of Lipopeptide-Producing Strains of *Bacillus* sp. Using a New Automated and Sensitive Fluorescence Detection Method. *Biotechnology Journal*, 14(4), 1–8. <https://doi.org/10.1002/biot.201800314>
- Kariyawasam, K. M. G. M. M., Jeewanthi, R. K. C., Lee, N.-K., & Paik, H.-D. (2019). Characterization of cottage cheese using *Weissella cibaria* D30: Physicochemical, antioxidant, and antilisterial properties. *Journal of Dairy Science*, 102, 1–7. <https://doi.org/10.3168/jds.2018-15360>
- Kim, E., Cho, E., Yang, S., Kim, M., & Kim, H. (2021). Novel approaches for the identification of microbial communities in kimchi: MALDI-TOF MS analysis and high-throughput sequencing. *Food Microbiology*, 94(June 2020), 103641. <https://doi.org/10.1016/j.fm.2020.103641>
- Larsen, M. V, Cosentino, S., Lukjancenko, O., Saputra, D., Rasmussen, S., Hasman, H.,

- Sicheritz-pontén, T., Aarestrup, F. M., Ussery, D. W., & Lund, O. (2014). Benchmarking of Methods for Genomic Taxonomy. *Journal of Clinical Microbiology*, *52*(5), 1529–1539. <https://doi.org/10.1128/JCM.02981-13>
- Lee, I., Chalita, M., Ha, S., Na, S., Yoon, S., & Chun, J. (2017). ContEst16S : an algorithm that identifies contaminated prokaryotic genomes using 16S RNA gene sequences. *Int J Syst Evol Microbiol*, *67*, 2053–2057. <https://doi.org/10.1099/ijsem.0.001872>
- Lefort, V., Desper, R., & Gascuel, O. (2015). FastME 2.0: A Comprehensive, Accurate, and Fast Distance-Based Phylogeny Inference Program. *Molecular Biology and Evolution*, *32*(10), 2798–2800. <https://doi.org/10.1093/molbev/msv150>
- Leong, K. H., Chen, Y. S., Lin, Y. H., Pan, S. F., Yu, B., Wu, H. C., & Yanagida, F. (2013). Weissellicin L, a novel bacteriocin from sian-sianzih-isolated *Weissella hellenica* 4-7. *Journal of Applied Microbiology*, *115*(1), 70–76. <https://doi.org/10.1111/jam.12218>
- Letunic, I., & Bork, P. (2016). Interactive tree of life (iTOL) v3: an online tool for the display and annotation of phylogenetic and other trees. *Nucleic Acids Research*, *44*(4), W242–W245. <https://doi.org/10.1093/nar/gkw290>
- Masuda, Y., Zendo, T., Sawa, N., Perez, R. H., Nakayama, J., & Sonomoto, K. (2012). Characterization and identification of weissellicin Y and weissellicin M, novel bacteriocins produced by *Weissella hellenica* QU 13. *Journal of Applied Microbiology*, *112*(1), 99–108. <https://doi.org/10.1111/j.1365-2672.2011.05180.x>
- Meier-kolthoff, J. P., & Göker, M. (2019). TYGS is an automated high-throughput platform for state-of-the-art genome-based taxonomy. *Nature Communications*, *10*, 2182. <https://doi.org/10.1038/s41467-019-10210-3>
- Moraes, P. M., Perin, L. M., Tassinari Ortolani, M. B., Yamazi, A. K., Viçosa, G. N., & Nero, L. A. (2010). Protocols for the isolation and detection of lactic acid bacteria with bacteriocinogenic potential. *Lwt*, *43*(9), 1320–1324. <https://doi.org/10.1016/j.lwt.2010.05.005>
- Nacef, M., Chevalier, M., Chollet, S., Drider, D., & Flahaut, C. (2017). MALDI-TOF mass spectrometry for the identification of lactic acid bacteria isolated from a French cheese: The Maroilles. *International Journal of Food Microbiology*, *247*, 2–8. <https://doi.org/10.1016/j.ijfoodmicro.2016.07.005>
- Neill, J. O. (2016). An audience with... Jim O'Neill. *Nature Reviews Drug Discovery*, *15*(August), 526. <https://doi.org/10.1038/nrd.2016.160>
- Nes, I. F. (2011). *Prokaryotic Antimicrobial Peptides: From Genes to Applications* (D. Djamel

- & S. Rebuffat (eds.); March, 8th). Springer-Verlag. https://doi.org/10.1007/978-1-4419-7692-5_1
- Ogunremi, O. R., Leischtfeld, S. F., & Schwenninger, S. M. (2022). MALDI-TOF MS profiling and exopolysaccharide production properties of lactic acid bacteria from Kunu-zaki - A cereal-based Nigerian fermented beverage. *International Journal of Food Microbiology*, 366(February), 109563. <https://doi.org/10.1016/j.ijfoodmicro.2022.109563>
- Seemann, T. (2014). Prokka: rapid prokaryotic genome annotation. *Bioin*, 30(14), 2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>
- Soltani, S., Biron, E., Said, L. Ben, Subirade, M., & Fliss, I. (2022). Bacteriocin-Based Synergetic Consortia: a Promising Strategy to Enhance Antimicrobial Activity and Broaden the Spectrum of Inhibition. *Microbiology Spectrum*, 10(1), e00406-21.
- Soltani, S., Hammami, R., Cotter, P. D., Rebuffat, S., Said, L. Ben, Gaudreau, H., Bédard, F., Biron, E., Drider, D., & Fliss, I. (2021). Bacteriocins as a new generation of antimicrobials: toxicity aspects and regulations. *FEMS Microbiology Reviews*, 45(March 2020), 1–24. <https://doi.org/10.1093/femsre/fuaa039>
- Sriornual, S., Yanagida, F., Lin, L. H., Hsiao, K. N., & Chen, Y. S. (2007). Weissellicin 110, a newly discovered bacteriocin from *Weissella cibaria* 110, isolated from plaa-som, a fermented fish product from Thailand. *Applied and Environmental Microbiology*, 73(7), 2247–2250. <https://doi.org/10.1128/AEM.02484-06>
- Teixeira, C. G., Fusieger, A., Martins, E., Freitas, R. de, Vakarelova, M., Nero, L. A., & Carvalho, A. F. de. (2021). Biodiversity and technological features of *Weissella* isolates obtained from Brazilian artisanal cheese-producing regions. *LWT - Food Science and Technology*, 147(April), 111474. <https://doi.org/10.1016/j.lwt.2021.111474>
- Teixeira, C. G., Rodrigues, R. da S., Yamatogi, R. S., Lucau-Danila, A., Drider, D., Nero, L. A., & de Carvalho, A. F. (2022). Genomic Analyses of *Weissella cibaria* W25, a Potential Bacteriocin-Producing Strain Isolated from Pasture in Campos das Vertentes, Minas Gerais, Brazil. *Microorganisms*, 10(2), 314. <https://doi.org/10.3390/microorganisms10020314>
- Tenea, G. N., Hurtado, P., & Ortega, C. (2020). A Novel *Weissella cibaria* Strain UTNGt210 Isolated from Wild Solanum quitoense Fruit: Genome Sequence and Characterization of a Peptide with Highly Inhibitory Potential toward Gram-Negative Bacteria. *Foods*, 9(9), E1242.
- Venegas-Ortega, M. G., Flores-Gallegos, A. C., Martínez-Hernández, J. L., Aguilar, C. N., &

- Nevárez-Moorillón, G. V. (2019). Production of Bioactive Peptides from Lactic Acid Bacteria: A Sustainable Approach for Healthier Foods. *Comprehensive Reviews in Food Science and Food Safety*, 18(June), 1039–1051. <https://doi.org/10.1111/1541-4337.12455>
- Yeu, J., Lee, H., Park, G., Lee, J., & Kang, M.-S. (2021). Antimicrobial and Antibiofilm Activities of *Weissella cibaria* against Pathogens of Upper Respiratory Tract Infections. *Microorganisms*, 9(6), 1181.
- Yu, H. S., Lee, N. K., Choi, A. J., Choe, J. S., Bae, C. H., & Paik, H. D. (2019). Antagonistic and antioxidant effect of probiotic *Weissella cibaria* JW15. *Food Science and Biotechnology*, 28(3), 851–855. <https://doi.org/10.1007/s10068-018-0519-6>
- Zidour, M., Chevalier, M., Belguesmia, Y., Cudennec, B., Grard, T., Drider, D., Souissi, S., & Flahaut, C. (2017). Isolation and characterization of bacteria colonizing *Acartia tonsa* copepod eggs and displaying antagonist effects against *Vibrio anguillarum*, *Vibrio alginolyticus* and Other pathogenic strains. *Frontiers in Microbiology*, 8(OCT), 1919. <https://doi.org/10.3389/fmicb.2017.01919>

CHAPTER V: *Weissella cibaria* strains isolated from Campo das Vertentes (MG), biosecurity and probiotics analyses

Biosecurity of *Weissella* strains isolated from Campos das Vertentes, Minas Gerais, Brazil focused on its potential as probiotic

Camila Gonçalves Teixeira, Yanath Belguesmia, Rafaela da Silva Rodrigues, Anca Lucau-Danila, Luís Augusto Nero, Antônio Fernandes de Carvalho, Djamel Drider

Abstract

Weissella cibaria referenced as W21, W25, and W42 have previously been shown to present antimicrobial potential against Gram-positive and Gram-negative foodborne pathogens. These antimicrobial properties are a good attribute to design these strains as potential probiotics as they can protect the gut microbiota from infections attributed to diverse pathogens. To reach the probiotic status, it is requested to establish additional criteria. This work aimed at evaluating the safety of these three strains of *W. cibaria*. Of note, the complete genome sequences of the aforementioned strains were previously sequenced and assembled. Here, we analyzed the genomes for safety-related genes, studied the antibiotic susceptibility, cytotoxicity in a mix of human colon adenocarcinoma Caco2 cell and HT-29, inflammation capacity in the promonocytic human cell line U937, and hemolytic activity of *W. cibaria* W21, W25, and W42. To strengthen the probiotic grade of these strains, we studied their survivability through the simulated human gastric digestion and adhesion to intestinal cell culture and exclusion of *S. aureus* MRSA SA1 and *E. coli* 184 in the intestinal cell culture. The results obtained showed overall that all strains have no acquired antimicrobial resistance genes and/or mobile genetic elements (MGE) in their genomes. *W. cibaria* strains were resistant to one antibiotic (vancomycin) in the *in vitro* test and this result was confirmed by genomic analysis, the resistance to vancomycin is intrinsic to this genus. Also, no hemolytic activity and no one of the known virulence factors were detected. In addition, the defense system CRISPR-Cas was verified and the genomes of the 3 strains harbor one CRISPR array each with evidence level 4. Besides they didn't present any inflammation or cytotoxicity capacity against intestinal cells and the three *Weissella* strains were predicted as a non-human pathogens. *W. cibaria* W21 presents higher potential, once it demonstrates excellent results of

adhesion to intestinal cells and exclusion of *S. aureus* MRSA SA1. However, none of the three strains survived through the simulated gastrointestinal tract. These results indicate that the tested strains of *W. cibaria* can be safely used as a potential probiotic for human consumption and animal feed and need some strategy to safety delivery them to the gut.

1. Introduction

Brazil is a country with several regions producing artisanal cheeses, leading to large collections and sources of beneficial microorganisms like lactic acid bacteria (LAB) that could be used in the food and pharmaceutical sectors after their characterization. [1]. Handmade cheeses and raw milk are considered potential sources of novel LAB strains [2], and each handmade cheese is supposed to contain its own microbiota because of the way it has been made. The microbiota present in the environment can get in contact with the product and then determine the fermentation process conferring therefore specific characteristics to the final product [3].

Weissella is a genus of the LAB group which is composed of bacteria characterized as Gram-positive, deprived of catalase activity, non-spore-forming, coccoid morphology, or short bacilli. Of note, strains belonging to the *Weissella* genus were isolated from different Brazilian regions producing artisanal cheeses and many strains have displayed antimicrobial activities against different foodborne pathogens [4].

The ability of LAB to produce antimicrobial compounds is of major interest to the food and pharmaceutical industries because they have less chance of being harmful to consumers. This is because they are frequently present in the gut microbiota and natural food fermentation such as raw milk cheese, and some members (*L. lactis* subsp. *lactis*, e.g.), possess the “Generally Recognized as Safe” status [5]. In addition, they can be used as alternatives to aging antibiotics, which are steadily facing harsh resistance and compromising therapeutic outcomes [6]. On the other hand, they can be substitutes to replace chemical preservatives in the food industry [7] and therefore meet the consumers' expectations.

Nonetheless, the use of a novel strain as an ingredient either for human or animal feed requires fulfilling a panel of strict criteria related to safety and discarding any adverse effect like the hemolytic activity, carriage of genes coding for antibiotic resistance, or pro-inflammation reaction, among others. [8,9]. This work aimed to study

the characteristics of three strains of *Weissella cibaria* (W21, W25, and W42), isolated in different farms, in Campos das Vertentes, in Minas Gerais, Brazil, that produce artisanal cheese made with raw milk, and establish their probiotic related grade in the view of their application in human consumption or animal feed.

2. Material and Methods

2.1. Safety of *Weissella* strains

2.1.1. *In silico* analyses of *Weissella*'s Genomes

To locate genes coding for virulence factors in the genomes of these *Weissella* strains, we used web tools recommended by the EFSA (European Food Safety Authority). Therefore, the research of mobile DNA elements like plasmids was evaluated by plasmidFinder [10], whilst the PathogenFinder web tool [11] was used to test the strains as potential human pathogens. We also checked the presence of genes coding the virulence factors using the “*Virulence Factors Database*” (VFDB) (<http://www.mgc.ac.cn/VFs/main.htm>) [12]. The search for genes coding for antibiotics resistance was realized using the RGI (Resistance Gene Identifier) tool and the “*Comprehensive Antibiotic Resistance Database*” (CARD) (<https://card.mcmaster.ca/>) [13]. For this analysis we import the contigs selecting the parameters “perfect and strict hits only” and “high-quality coverage”. Finally, the ResFinder 4.1 [14] server (<https://cge.cbs.dtu.dk/services/ResFinder/>) was employed to identify the acquired antimicrobial resistance genes with a selected % ID threshold of 90 % and the selected minimum length of 60 %.

For the detection and annotation of prophage sequences within bacterial genomes, the PHASTER [15] (PHAge Search Tool Enhanced Release) (<http://phaster.ca>) tool was used. The CRISPRCasFinder [16] (<https://crisprcas.i2bc.paris-saclay.fr/CrisprCasFinder/Index>) was also used to detect the CRISPR and Cas genes by importing the contig FASTA sequences and selecting “Perform cas detection” and leaving the rest of the parameters by default.

2.1.2. *Hemolytic activity and antibiotic susceptibility*

We realized *in vitro* tests for hemolytic activity and resistance to antibiotics according to Colombo et al. [17]. For hemolytic activity, the cultures were streaked on the Columbia Blood Agar Settle plate (Merck KGaA, Darmstadt, Germany). After a

period of incubation of 24 hours at 37 °C, the plates were inspected for any hemolytic activity associated with each strain, taking into consideration a total or β -hemolysis (clear halos around the colonies), partial or α -hemolysis (greenish halos around the colonies), or γ -hemolysis (absence of hemolysis) [17].

Antibiotics resistance was characterized based on the disk diffusion assay (Oxoid) according to Colombo et al. [17]. Each strain was diluted using 0.85 % NaCl (w/v) until the turbidity was equivalent to 0.5 McFarland standard and swabbed onto the surface of an MRS agar plate (where the antibiotic disks were placed). A panel of 7 antibiotics [imipenem (carbapenem), vancomycin (glycopeptide), clindamycin (lincosamide), erythromycin (macrolide), ampicillin (penicillin), ciprofloxacin (fluoroquinolone) and Amoxicillin + Clavulanic Acid]. The plates were incubated at 37 °C for 18 h, when the diameter of the inhibition halos was measured and the isolates were characterized as presenting resistance (R), intermediary resistance (IR), or sensitivity (S), and the size of the halo was also measured, and analyzed according to CLSI breakpoints [18].

2.2. Cytotoxicity and Inflammatory activity of *Weissella* strains in human cell

2.2.1. Cytotoxicity assay

For the cytotoxicity test, we utilized a mix of human cell cultures with 80 % of Caco2 and 20 % of HT-29. The preparation of the mix of human cell culture was according to Pouille et al. [19]. The mix of human colon adenocarcinoma Caco2 cell and HT-29 line were cultured in Dulbecco's Modified Eagle Medium (DMEM, Gibco, Thermo Fisher Scientific) under standard cell culture conditions (37 °C and 5 % CO₂). The culture medium was changed regularly and when the cells reached sub-confluence (80– 90 %), they were sub-passaged. The cells were cultured at 1.5×10^4 cells/mL in 96-well plates for 24 h and then exposed to the 3 different *Weissella* strains.

The preparation of the culture of *Weissella* strains to contact the mix of human cell culture was realized as described below. One percent of frozen *Weissella* strains were propagated in MRS broth (Oxoid) for 24h, centrifugated at 10000 rpm for 5 min and the supernatant was removed. After the obtained pellet was washed with 5 mL of phosphate buffer saline (PBS, pH 7.4), and the pellet was resuspended in 2 mL of DMEM without antibiotics and fetal bovine serum and diluted until a final concentration of 10^7 CFU/mL. The suspension of cells was incubated at 37 °C until contact with the

mix of human cells. For the contact, the mix of human cell culture fixed in the bottle of the 96 wells plate was washed with DMEM and the suspensions of *Weissella* strains were added in a proportion of 1:1 and 1:10 (v/v). After the plate was incubated for 24h at 37 °C.

Cell viability of Caco2 and HT-29 after 24 h of incubation with *Weissella* strains was measured using a cell counting kit-8 assay (CCK-8, Dojindo Molecular Technologies, Inc. Japan). After incubation, the 96-well plate was washed with DMEM with antibiotics (100 U/mL penicillin and 100 µg/mL streptomycin). After the washing procedure and removing all *Weissella* strains cells, 150 µL of CCK-8 solution (5 %) was added to each well of the 96-well plate, followed by incubation for 1.5 h at 37 °C. Then, 10 mL of stop solution provided in the CCK-8 kit was added to each well. The absorbance of the mixture was measured at 450 nm using a microplate reader (Synergy H1, BioTek, USA). Cell viability was calculated based on the relative absorbance compared with the control group.

2.2.2. Inflammatory effects of *Weissella* strains

For the inflammatory and anti-inflammatory capacity of *Weissella* strains, we used the promonocytic human cell line U937 according to the protocol described by Pouille et al. [19]. The cell line U937 was cultured in Roswell Park Memorial Institute 1640 Medium (RPMI, Gibco, Thermo Fisher Scientific) supplemented with 10 % fetal bovine serum (FBS), 100 U/mL penicillin, 100 µg/mL streptomycin, and 2 mM glutamine in a humidified 5 % CO₂ atmosphere at 37 °C. The culture medium was replaced regularly and when the cells reached sub-confluence (80–90 %), they were sub-passaged. For the differentiation of the cell line in macrophages, U937 cells were seeded at approximately 1×10^5 cells/well in 96-well plates with 60 ng/mL of phorbol-12-myristate-13-acetate (PMA, Thermo Fisher Scientific) for 48 h. The adherent cells were washed with PBS and incubated for 2 h with LPS (10 µg·mL⁻¹; LPS from *E. coli* O26:B6, Millipore Corporation) and *Weissella* strains. Cell supernatants were collected on ice and centrifuged (1500 rpm 5 min) to eliminate cell debris. The supernatants were aliquoted and stored at -80 °C until further analysis.

Weissella strains samples were prepared as described above, resuspended in not supplemented RPMI, and added to each well in 1:10 MOI (multiplicity of infection). The inflammatory control is U937 differentiated with LPS (50 µg/mL) and the anti-

inflammatory control is U937 differentiated with LPS (50 µg/mL) and dexamethasone (20 µM).

Multiple cytokines in culture supernatants of the U937 cell line were detected with a customized Milliplex Map kit (Human High Sensitivity T Cell Magnetic Bead Panel, Millipore, Billerica, MA), following the manufacturer's instructions. Briefly, antibody-immobilized beads for detection of interleukin -1 β (IL-1 β), IL-2, IL-4, IL-5, IL-6, IL-7, IL-10, IL-12 (p70), IL-13, tumor necrosis factor-alpha (TNF- α), granulocyte-macrophage colony-stimulating factor (GM-CSF) and interferon- γ (IFN γ) were incubated with 5-times-diluted culture supernatants of U937. The beads were subsequently treated with detection antibodies as well as PE-conjugated streptavidin and the quantification was carried out using the Luminex® 100/200 (Luminex Corporation, Austin, TX, USA) system and the Luminex xPONENT® for LX100/200 software. The cytokine IL-8 was quantified using a human IL-8/CXCL8 Quantikine® ELISA kit (R&D Systems, Inc., Minneapolis, MN, USA). Medium samples were diluted 100-fold according to the kit recommendations.

2.3. Survival of *Weissella cibaria* strains under conditions mimicking a human gastric digestion

The survival of *Weissella* strains in simulated human gastric and intestinal juices was assessed as described by Fonseca et al. [20] with modifications.

The composition of the simulated gastric juice was pepsin (2 g/L, Sigma Aldrich) and PBS pH 3.5 (adjusted with HCl 0.5 M). Each isolate was grown at 37 °C overnight in sterile MRS broth (Oxoid) and inoculated in the solution until the final concentration of 1×10^{10} CFU/mL and followed by incubation at 37°C for 90 min with constant shaking at 160 rpm. After incubation, the samples were taken and serially diluted, then plated on MRS agar plates and incubated at 37 °C for 24 hours. Then, the simulated duodenal juice was prepared by adding pancreatin (Dinâmica, Brazil) and bile salts solutions to obtain final concentrations of 0.2 % and 0.5 % (w/v), respectively, and the pH was then adjusted to 7.0 by adding 1 M sodium hydroxide (NaOH). After mixing, samples were incubated at 37 °C for 180 min with agitation (160 rpm), and then viable cell counts were determined. All sample counts were determined by plating on MRS (Oxoid) agar. The experiments were repeated three times and performed in triplicate. Results were expressed as mean log colony-forming units per mL (CFU/mL). The rate of survival

was calculated as the percentage of *Weissella* colonies grown on MRS agar relative to the initial bacterial concentration:

$$\text{Survival rate (\%)} = \frac{\log CFUN1}{\log CFUN0} \times 100$$

Where N1 is the viable count of isolates after incubation and N0 is the initial viable count.

2.4. Adhesion of *Weissella* strains to the human Caco-2 cell line

The adhesion capacity test for the three selected strains to the human colon adenocarcinoma cell lines (Caco-2) was performed according to Fonseca et al. [20] with slight modifications. The Caco-2 cells were sub-cultured (2×10^5 cells/mL) in 24-well tissue culture plates (Sarstedt, Germany) and grown at 37 °C in a humidified atmosphere of 5 % CO₂ for 21 days to promote differentiation in cell media. The culture medium was changed on alternate days. For the adhesion assay, bacteria were cultured in MRS broth overnight at 32 °C, and after washing twice with PBS, the cultures were resuspended in the DMEM at a concentration of approximately 10⁷ CFU/mL. After the DMEM of the 24-well tissue culture plates was removed and the culture cells were washed with the same media. Then to each well was added 400 µL of DMEM and 100 µL of *Weissella* strains. The plate was incubated at 37 °C overnight in a 5 % CO₂ atmosphere. Subsequently, the cells were washed three times with 1 mL of PBS to remove non-adherent bacteria cells and then lysed with 300 µL of trypsin. After 15 min of incubation at 37 °C, the solution with released bacteria cells was transferred to 1.5 mL Eppendorf and the bacteria cells were centrifugated (9000 × g/ 5 min), the supernatant was removed and the pellet was resuspended in 500 µL of PBS. After recovering the bacteria cells were serially diluted and plated on MRS agar. The plates were incubated at 32 °C for 24 h. After this time, enumerations of *W. cibaria* strains colonies were performed. Experiments were performed in duplicate and repeated three times.

2.5. In situ inhibition of pathogenic bacteria

For this analysis, Caco-2 cell line cultures were maintained and *Weissella* strains were propagated as above described. Before the experiment, the DMEM of the 24-well tissue culture plates was removed and the culture cells were washed with the same media. For the pathogen inhibition test, the exclusion assay was performed

according to the protocol described by Fonseca et al. [20] with modifications. The pathogens used in this experiment were *Escherichia coli* ATCC 8739 and *Staphylococcus aureus* ATCC 43300, which were propagated separately in BHI medium (Oxoid) at 37 °C for 24h.

Caco-2 cells were first preincubated with 100 µL of *Weissella* strains (10^7 CFU/mL) for 2h at 37 °C in a 5 % CO₂ atmosphere and then 100 µL of *E. coli* (10^7 CFU/mL) or *S. aureus* (10^7 CFU/mL) was added to each well. The plates were incubated at 37 °C overnight in a 5 % CO₂ atmosphere.

After the incubation period, the cells were washed three times with 1 mL of PBS to remove non-adherent bacteria cells and then lysed with 300 µL of trypsin. After 15 min of incubation at 37°C, the solution with released bacteria cells was transferred to 1.5 mL Eppendorf and the bacteria cells were centrifugated ($9000 \times g/ 5 \text{ min}$), the supernatant was removed and the pellet was resuspended in 500 µL of PBS. After recovering, the bacteria cells were serially diluted and plated. *E. coli* was plated in Eosin Methylene Blue agar (EMB) and *S. aureus* in BHI supplemented with the antibiotic erythromycin (100 µL/mL) that prevented the growth of *Weissella* strains. The plates were incubated at 37 °C for 24 h. After this time, enumerations of pathogen colonies were performed. Experiments were performed in duplicate and repeated three times.

3. Results and Discussion

3.1. Safety of *Weissella* strains

Based on the ResFinder data, no acquired antimicrobial resistance genes were found in the genomes of the 3 tested *Weissella* strains. Moreover, the research for plasmid evaluated by plasmidFinder [10] concluded the absence of mobile genetics elements, arguing for the stability of the genomes.

[13], [13].

According to the findings of CRISPRCasFinder [16], each of the three *Weissella* strains' genomes harbor a single CRISPR array, with evidence level 4 indicating that these arrays may be legitimate [16].

The CRISPR-associated protein Csn2_TypellA is a mandatory protein of the Type IIA system and the gene is present in the three sequenced genomes. According to Roberts and Barrangou [21], the Type II systems are rare in nature compared to

types I and III, which are also more well-characterized [21]. The CRISPR-Cas system is a defense mechanism used by bacteria to protect themselves against infections caused by foreign genetic elements. Thus, strains endowed with the CRISPR-Cas system could acquire fewer sequences originating from other bacteria preventing horizontal gene transference and therefore leading to low level of drug resistance [22]. Related to that, the CARD analysis only locate genes involved in the resistance to vancomycin, (*vanY* gene in the *vanB* cluster and *vanT* gene in the *vanG* cluster) for all 3 tested strains (Table 2), and this resistance was interestingly confirmed by the *in vitro* test for antibiotic resistance (Table 3), demonstrating the convergence of these two approaches and their complementarity. In a direct line, Muñoz-Atienza et al. reported that 60% of *Weissella* species are resistant to several antibiotics [23]. Recently, Tenea and Hurtado found putative resistance genes in the genome of *W. cibaria* UTNGt210 [24].

Table 1. CRISPR and Cas genes found in genomes of the 3 *Weissella cibaria* strains isolated from Campos das Vertentes, Minas Gerais, Brazil.

Strain	Region	CRISPR region					CAS protein			
		Start	End	DR consensus	DR Length	Spacers	Gene name	Start	End	Orientation
W25	C4	12728	13621	GTTTTAGT GTCATGTT GAATAGAA TGCTTCTC AAAC	36	13	Type: CAS-TypellA Start: 6,032; End: 12,578			
							cas9_Typell	6,032	10,498	+
							cas1_Typell	10,710	11,582	+
							cas2_TypeI-II-III	11,579	11,914	+
							csn2_TypellA	11,907	12,578	+
W42	C22	95084	95977	GTTTGAG AAGCATT TATTCAAC ATGACACT AAAAC	36	13	Type: CAS-TypellA Start: 96,127; End: 102,673			
							csn2_TypellA	96,127	96,798	-
							cas2_TypeI-II-III	96,791	97,126	-
							cas1_Typell	97,123	97,995	-
							cas9_Typell	98,207	102,673	-
W21	C25	12686	13579	GTTTTAGT GTCATGTT GAATAGA ATGCTTCT CAAAC	36	13	Type: CAS-TypellA Start: 11,865; End: 12,536			
							csn2_TypellA	11,865	12,536	+
							Type: CAS-TypellU Start: 5,990; End: 12,536			
							cas9_Typell	5,990	10,456	+
							cas1_Typell	10,668	11,540	+
							csn2_TypellA	11,865	12,536	+

Table 2. Antibiotic resistance genes identified in *Weissella cibaria* strains using CARD (Comprehensive Antibiotic Resistance Database) webserver.

Strain	RGI Criteria	ARO Term	Detection Criteria	AMR Gene Family	Drug Class	Resistance Mechanism	% Identity of Matching Region	% Length of Reference Sequence
W21 W25 W42	Strict	vanY gene in vanB cluster	Protein homolog model	vanY, Glycopeptide resistance gene cluster	Glycopeptide antibiotic	Antibiotic Target alteration	34.95	97.39
	Strict	vanT gene in vanG cluster	protein homolog model	Glycopeptide resistance gene cluster, Van T	Glycopeptide antibiotic	Antibiotic target alteration	33.33	69.38

Table 3. Results from *in vitro* antibiotic resistance test.

Code	Antibiotics spectrum (mm)						
	Ampicillin	Amoxicillin + clavulanic acid	Imipenem	ciprofloxacin	Vancomycin	Erythromycin	Clindamycin
W21	29	29	35	24	0	24	24
W25	31	31	33	20	0	27	26
W42	28	32	32	22	0	27	26

Results are expressed in mm of inhibition halo from antibiotics. Interpretation according to CLSI breakpoints [18].

Overall, the CARD and the *in vitro* antibiotic tests data obtained here corroborate the low level of resistance, once the three tested strains presented resistance to vancomycin, which is intrinsic for this genus [25]. Intrinsic resistance is not redhibitory or a safety concern [8] to proceed with the selection of these strains as probiotics candidates. Moreover, the number of spacers in the CRISPR-Cas system represents the ability to eliminate heterologous genes, such as prophage. Thus, the more spacer sequences are present, the more the strain has a limited capacity to acquire foreign sequences like prophages [22]. Here, we only could locate using PHASTER only two prophages (1 entire and 1 incomplete) in the genomes of the three strains (Table 4). Dong et al. demonstrated that the correlation between spacers and prophages is negative [22].

Table 4. Prophage regions identified in *Weissella cibaria* strains genome using the PHASTER webserver.

Strain	Region	Length (kb)	Completeness*	Score	GC %	Most Common Phage
W21	1	17.9	Incomplete	20	41.52	PHAGE_Lactoc_PLgT_1_NC_031016(3)

	2	32	Intact	110	42.64	PHAGE_EnterophiFL1A_NC_013646(14)
W42	1	17.9	Incomplete	20	41.51	PHAGE_Lactoc_PLgT_1_NC_031016(3)
	2	35.8	Intact	110	42.30	PHAGE_EnterophiFL1A_NC_013646(14)
W25	1	17.9	Incomplete	20	41.54	PHAGE_Lactoc_PLgT_1_NC_031016(3)
	2	35.8	Intact	110	42.30	PHAGE_EnterophiFL1A_NC_013646(14)


*Intact (score > 90); Questionable (score 70-90); Incomplete (score < 70)

Prophages are MGE which is considered the primary factor in genetic diversity enabling niche adaptation in bacteria [26]. As above-mentioned, two prophages were located in the genomes of the three strains (W21, W25, and W42), one of them was intact, whereas the second was incomplete in each of the tested strains. The length of the intact prophage varies from 32 to 35.8 Kb with the most common phage: PHAGE_EnterophiFL1A_NC_013646(14). The presence of intact regions could be a selective advantage and help bacteria against other prophage infections [27,28].

Several genes were predicted as putative virulence factors by VFDB (data not shown). However, no one of the well-known virulence genes *gelE* (gelatinase), *hyl* (hyaluronidase), *asa1* (aggregation substance), *esp* (enterococcal surface protein), *cylA* (cytolysin), *efaA* (endocarditis antigen), and *ace* (adhesion of collagen), were detected in the genomes of our strains. Besides, most of the genes detected by the VFDB webserver are related to other cellular functions such as genes for putative enolase (*eno*) phosphopyruvate hydratase. Enolase is a key glycolytic enzyme in the cytoplasm of prokaryotic essential for the degradation of carbohydrates via glycolysis [29].

Some researchers reported the presence of some virulence-encoded genes such as collagen adhesin proteins and genes associated with toxin production systems (including botulinum neurotoxin homolog) in some *Weissella* species like *W. cebi*, *W. confusa* [30] and *W. oryzae* [31], respectively. None of these genes were found in our *W. cibaria* strains tested. Also, Tenea and Hurtado report two genes (*hlyD*, *hlyIII*) in some strains of *W. cibaria* [24], however in our study no genes involved in hemolysis activity were observed in the genomes of the *W. cibaria* strains W21, W25, and W42; and this was phenotypically confirmed by assessing this activity on the blood-agar

medium (Figure 1). Therefore, slight hemolysis that could be graded as an α -hemolysis was detected.



Control +: *S. aureus* ATCC 29213

Figure 1. Hemolytic activity of *Weissella* strains W21, W25, and W42.

Despite several virulence factors detected by the VFDB web server, the results indicate that no one known virulence factor was present among them and the analysis of PathogenFinder [11] indicates that the three input organisms were predicted as a non-human pathogens. Therefore, these three strains of *W. cibaria* can be considered safe for use as probiotics.

3.2. Cytotoxicity and Inflammatory activity of *W. cibaria* strains in human cell

For the cytotoxicity test, we used a mix of human cell cultures with 80 % of Caco2 and 20 % of HT-29 5 (Figure 2A) and another experiment also with the human U937 cells (Figure 2B). It is noteworthy that no cytotoxicity trait was found against these human lines, matching with other studies that underpinned the safety of *W. cibaria* strains on RAW 264.7 macrophage cell line and human mouth epithelial cell line KCLB 10017 [32,33]. Interestingly, we assessed the inflammatory and anti-inflammatory capacity of *Weissella* strains against the human U937 cells and these strains showed neither inflammatory nor anti-inflammatory responses. Taken all together, these data

strengthen the safety aspects of these strains and open the gate for their potential use for human consumption and animal feed.




Figure 3. Effect of *W. cibaria* strains on cytotoxicity of 80 % of Caco2 + 20 % of HT-29 cells (A); and U937 cells (B).

3.3. Survival of *W. cibaria* strains in the simulated gastro-intestinal tract

We noted during the assessment of the survivability of *Weissella* strains in the gastrointestinal tract, a small loss of the viability of the cells after 1.5 h in conditions mimicking the stomach compartment, and also no survivability was observed upon 3 h of incubation in conditions simulating those of the duodenal compartment (Table 5).

Table 2. Survival of *Weissella cibaria* strains under conditions simulating gastric digestion.

Strains	Time						Survival rate (%) of time 1.5h
	0h		1.5h		4.5h		
	pH	log10 (CFU/mL)	pH	log10 (CFU/mL)	pH	log10 (CFU/mL)	
W25	3.5	10.04	3.5	9.68	7	ND	96.4
W42	3.5	9.80	3.5	9.37	7	ND	95.6
W21	3.5	9.80	3.5	9.60	7	ND	98.0

ND: not detected by technique limit

The ability of bacteria to survive under gastric digestion conditions is considered a key criterion for probiotic selection [34]. Of note, there are strategies like encapsulation that could help the passage through the gastrointestinal tract enabling protection of bacteria cells from stress conditions and making possible the control of their delivery process directly in the intestine where they can act as probiotics and provide their benefits for the host [35,36].

Besides that, the method of assessing gastrointestinal tract survival by agar enumeration does not take into account the viable but nonculturable cells (VBNC) [37]. Previous studies used the cytometry method, which includes both viable bacteria and VBNC and as expected, the survival percentages obtained by this method were significantly higher in comparison to the data obtained after their growth and enumeration by agar plating method [34].

3.4. Adhesion of *W. cibaria* strains and Inhibition of pathogenic bacteria in Caco-2 cell line

The adhesion of *W. cibaria* strains to the Caco-2 cell lines was almost 100 % effective (Table 6). The initial inoculum was 7-8 log CFU/mL and all the 3 tested strains were able to adhere in the Caco-2 cell line. The adhesion to the intestinal cell is important and renders LAB strains a barrier against enteropathogenic bacteria avoiding them to adhere in the intestinal cells and competing with them for nutrients [38,39].

Table 3. Adhesion of *Weissella cibaria* strains to Caco-2 cells and exclusion of pathogenic bacteria by *W. cibaria* strains.

Strains	Adhesion	Exclusion	
	log ₁₀ (CFU/mL)	<i>S. aureus</i> MRSA SA1	<i>E. coli</i> 184
Control		6.53228176	6.46542394
W25	7.335792102	6.6515239	6.49991994
W42	6.33243846	6.85370046	6.42275553
W21	7.33243846	ND	6.44388297

ND: not detected by technique limit

The data from the exclusion assays revealed that only *W. cibaria* W21 was able to impede the adhesion of *S. aureus* MRSA SA1 to the Caco-2 cells (Table 6) being very effective, suggesting a strong candidate as a probiotic strain. Here only two

pathogenic strains were tested: *S. aureus* MRSA SA1 and *E. coli* 184, but of note, we reported that the strains W21, W25, and W42 are also able to inhibit *L. monocytogenes*, and *Salmonella enterica* [4]. The inhibition of adhesion of pathogens in human cells may be related to different mechanisms: antimicrobial substances produced by LAB, and competition for binding sites, among others [40]. Besides, some pathogens have different levels of adhesion and invasion capacities depending on the eukaryotic cell, some pathogens have more or less capacity in mucus-producing intestinal cell models (HT-29-MTX) compared to non-mucus-producing cells (Caco-2) [41]. That ability is directly related to the expression of specific proteins, pili, fimbriae, and flagella [42].

4. Conclusion

All three *W. cibaria* strains appear to be safe for use for human consumption according to the genome analysis, presenting no acquired antimicrobial resistance genes and no mobile elements in the genome. Moreover, it was shown that the genomes have a CRISPR-Cas system which is a defense system against infections caused by heterologous genes. They also didn't present any antibiotic resistance genes except the vancomycin resistance gene (confirmed by *in vitro* tests) which is an intrinsic resistance gene for the *Weissella* genus and other LAB with GRAS status. Only one intact prophage was detected in the genome of these strains, which can also help the bacteria against other prophage infections. Finally, no one of the known virulence factors was detected in these genomes and no *in vitro* hemolytic activity was detected. No cytotoxicity and no inflammatory effect were detected for the *Weissella* strains tested in this study.

Regarding the probiotic traits of the *W. cibaria* W21, this strain shows higher potential as it demonstrates a high exclusion ability of *S. aureus* MRSA SA1 in the intestinal cells. However, none of the three strains survived through the simulated gastrointestinal tract. To overcome this further studies regarding protective strategies to help the passage of these strains to the gastric system are needed and more studies with other enteropathogenic strains should be explored.

5. References

1. Finnegan, W.; Yan, M.; Holden, N.M.; Goggins, J. A review of environmental life cycle assessment studies examining cheese production. *Int. J. Cycle Assess.* **2018**, *23*, 1773–1787.
2. Silva, E.O.O. e; Nespolo, C.R.; Sehn, C.P.; Pinehiro, F.C.; Stefani, L.M. Lactic acid bacteria with antimicrobial, proteolytic and lipolytic activities isolated from ovine dairy products. *Food Sci. Technol.* **2020**, *40*, 293–299.
3. Pyz-Łukasik, R.; Knysz, P.; Gondek, M. Hygiene Quality and Consumer Safety of Traditional Short- and Long-Ripened Cheeses from Poland. *J. Food Qual.* **2018**, *2018*.
4. Teixeira, C.G.; Fusieger, A.; Martins, E.; Freitas, R. de; Vakarelova, M.; Nero, L.A.; Carvalho, A.F. de Biodiversity and technological features of *Weissella* isolates obtained from Brazilian artisanal cheese-producing regions. *LWT - Food Sci. Technol.* **2021**, *147*, 111474, doi:10.1016/j.lwt.2021.111474.
5. Chittora, D.; Meena, B.R.; Jain, T.; Sharma, K. Biopreservation : bacteriocins and lactic acid bacteria. *J. Postharvest Technol.* **2022**, *10*, 1–15.
6. Fernandes, A.; Jobby, R. Bacteriocins from lactic acid bacteria and their potential clinical applications. *Appl. Biochem. Biotechnol.* **2022**, *194*, 4377–4399, doi:10.1007/s12010-022-03870-3.
7. Mokoena, M.P.; Omatola, C.A.; Olaniran, A.O. Applications of Lactic Acid Bacteria and Their Bacteriocins against Food Spoilage Microorganisms and Foodborne Pathogens. *Molecules* **2021**, *26*, 7055.
8. Rychen, G.; Aquilina, G.; Azimonti, G.; Bampidis, V.; Bastos, M. de L.; Bories, G.; Chesson, A.; Cocconcelli, P.S.; Flachowsky, G.; Gropp, J.; et al. Guidance on the characterisation of microorganisms used as feed additives or as production organisms. *EFSA J.* **2018**, *16*, 1–24, doi:10.2903/j.efsa.2018.5206.
9. Panel, E.B.; Koutsoumanis, K.; Allende, A.; Alvarez-Ordóñez, A.; Bolton, D.; Bover-Cid, S.; Chemaly, M.; Davies, R.; De Cesare, A.; Hilbert, F.; et al. The list of QPS status recommended biological agents for safety risk assessments carried out by EFSA. **2019**, doi:10.5281/ZENODO.3336268.
10. Carattoli, A.; Zankari, E.; García-fernández, A.; Larsen, M.V.; Lund, O.; Villa, L.; Aarestrup, F.M.; Hasman, H. In Silico Detection and Typing of Plasmids using PlasmidFinder and Plasmid Multilocus Sequence Typing. *Antimicrob. Agents*

- Chemother.* **2014**, *58*, 3895–3903, doi:10.1128/AAC.02412-14.
11. Cosentino, S.; Larsen, M.V.; Aarestrup, F.M.; Lund, O. PathogenFinder - Distinguishing Friend from Foe Using Bacterial Whole Genome Sequence Data. *Plus One* **2013**, *8*, e77302, doi:10.1371/journal.pone.0077302.
 12. Liu, B.; Zheng, D.; Zhou, S.; Chen, L.; Yang, J. VFDB 2022: A general classification scheme for bacterial virulence factors. *Nucleic Acids Res.* **2022**, *50*, D912–D917, doi:10.1093/nar/gkab1107.
 13. Alcock, B.P.; Raphenya, A.R.; Lau, T.T.Y.; Tsang, K.K.; Bouchard, M.; Edalatmand, A.; Huynh, W.; Nguyen, A.L. V.; Cheng, A.A.; Liu, S.; et al. CARD 2020: Antibiotic resistome surveillance with the comprehensive antibiotic resistance database. *Nucleic Acids Res.* **2020**, *48*, D517–D525, doi:10.1093/nar/gkz935.
 14. Zankari, E.; Hasman, H.; Cosentino, S.; Vestergaard, M.; Rasmussen, S.; Lund, O.; Aarestrup, F.M.; Larsen, M.V. Identification of acquired antimicrobial resistance genes. **2012**, 2640–2644, doi:10.1093/jac/dks261.
 15. Arndt, D.; Grant, J.R.; Marcu, A.; Sajed, T.; Pon, A.; Liang, Y.; Wishart, D.S. PHASTER : a better, faster version of the PHAST phage search tool. *Nucl. Acids Res.* **2016**, *44*, 16–21, doi:10.1093/nar/gkw387.
 16. Couvin, D.; Bernheim, A.; Toffano-nioche, C.; Touchon, M.; Rocha, E.P.C.; Vergnaud, G.; Michalik, J.; Bertrand, N.; Gautheret, D.; Pourcel, C.; et al. CRISPRCasFinder, an update of CRISRFinder, includes a portable version, enhanced performance and integrates search for Cas proteins. **2018**, *46*, 246–251, doi:10.1093/nar/gky425.
 17. Colombo, M.; Nero, L.A.; Todorov, S.D. Safety profiles of beneficial lactic acid bacteria isolated from dairy systems. *Brazilian J. Microbiol.* **2020**, *51*, 787–795, doi:10.1007/s42770-020-00227-y.
 18. CLSI Performance standards for antimicrobial susceptibility testing. *CLSI Suppl. M100, Clin. Lab. Standars Inst.* **2022**, *32nd edn*, 362.
 19. Pouille, C.L.; Ouaza, S.; Roels, E.; Behra, J.; Turret, M.; Molinié, R.; Fontaine, J.-X.; Mathiron, D.; Gagneul, D.; Taminiau, B.; et al. Chicory: Understanding the Effects and Effectors of This Functional Food. *Nutrients* **2022**, *14*, 957.
 20. Fonseca, H.C.; de Sousa Melo, D.; Ramos, C.L.; Dias, D.R.; Schwan, R.F. Probiotic Properties of Lactobacilli and Their Ability to Inhibit the Adhesion of Enteropathogenic Bacteria to Caco-2 and HT-29 Cells. *Probiotics Antimicrob. Proteins* **2021**, *13*, 102–112, doi:10.1007/s12602-020-09659-2.

21. Roberts, A.; Barrangou, R. Applications of CRISPR-Cas systems in lactic acid bacteria. *FEMS Microbiol. Rev.* **2020**, *44*, 523–537, doi:10.1093/femsre/fuaa016.
22. Dong, J.; Feng, C.; Wang, P.; Li, R.; Zou, Q. Comparative genomics analysis of *Acinetobacter baumannii* multi-drug resistant and drug sensitive strains in China. *Microb. Pathog.* **2022**, *165*, 105492, doi:10.1016/j.micpath.2022.105492.
23. Muñoz-atienza, E.; Gómez-sala, B.; Araújo, C.; Campanero, C.; Campo, R.; Hernández, P.E.; Herranz, C.; Cintas, L.M. Antimicrobial activity, antibiotic susceptibility and virulence factors of Lactic Acid Bacteria of aquatic origin intended for use as probiotics in aquaculture. *BMC Microbiol.* **2013**, *13*.
24. Tenea, G.N.; Hurtado, P. Next-Generation Sequencing for Whole-Genome Characterization of *Weissella cibaria* UTNGt21O Strain Originated From Wild Solanum quitoense Lam . Fruits: An Atlas of Metabolites With Biotechnological Significance. *Front. Microbiol.* **2021**, *12*, 1–16, doi:10.3389/fmicb.2021.675002.
25. Teixeira, C.G.; Fusieger, A.; Milião, G.L.; Martins, E.; Drider, D.; Nero, L.A.; de Carvalho, A.F. *Weissella*: An Emerging Bacterium with Promising Health Benefits. *Probiotics Antimicrob. Proteins* **2021**, *13*, 915–925, doi:10.1007/s12602-021-09751-1.
26. Fontana, A.; Falasconi, I.; Molinari, P.; Treu, L.; Basile, A.; Vezzi, A.; Campanaro, S.; Morelli, L. Genomic Comparison of *Lactobacillus helveticus* Strains Highlights Probiotic Potential. *Front. Microbiol.* **2019**, *10*, 1–14, doi:10.3389/fmicb.2019.01380.
27. Mercanti, D.J.; Rousseau, G.M.; Capra, M.L.; Quiberoni, A.; Tremblay, D.M.; Labrie, S.J. Genomic Diversity of Phages Infecting Probiotic Strains of *Lactobacillus paracasei*. *Appl. Environ. Microbiol.* **2016**, *82*, 95–105, doi:10.1128/AEM.02723-15.Editor.
28. Falasconi, I.; Fontana, A.; Patrone, V.; Rebecchi, A.; Garrido, G.D.; Principato, L.; Callegari, M.L.; Spigno, G.; Morelli, L. Genome-assisted characterization of *Lactobacillus fermentum*, *Weissella cibaria*, and *Weissella confusa* strains isolated from sorghum as starters for sourdough fermentation. *Microorganisms* **2020**, *8*, 1–16, doi:10.3390/microorganisms8091388.
29. Fukano, K.; Kinura, K. Measurement of Enolase Activity in Cell Lysates. In *Methods in Enzymology*; Galluzzi, L., Kroemer, G., Eds.; Academic Press, 2014; pp. 115–124 ISBN 9780124166189.
30. Abriouel, H.; Lerma, L.L.; Casado Muñoz, M. del C.; Montoro, B.P.; Kabisch, J.; Pichner, R.; Cho, G.S.; Neve, H.; Fusco, V.; Franz, C.M.A.P.; et al. The controversial nature of the *Weissella* genus: Technological and functional aspects versus whole genome

- analysis-based pathogenic potential for their application in food and health. *Front. Microbiol.* **2015**, *6*, 1–14, doi:10.3389/fmicb.2015.01197.
31. Mansfield, M.J.; Adams, J.B.; Doxey, A.C. Botulinum neurotoxin homologs in non-Clostridium species. *FEBS Lett.* **2015**, *589*, 342–348, doi:10.1016/j.febslet.2014.12.018.
 32. Kim, M.I.N.J.; You, Y.O.U.K.; Kang, J.O.O.Y.I.; Kim, H.J.I.N.; Kang, M.I.S.U.N. *Weissella cibaria* CMU exerts an anti-inflammatory effect by inhibiting *Aggregatibacter actinomycetemcomitans*-induced NF- κ B activation in macrophages. *Mol. Med. Reports* **2020**, *22*, 4143–4150, doi:10.3892/mmr.2020.11512.
 33. Kang, M.; Lim, H.; Kim, S.; Lee, H.; Oh, J. Effect of *Weissella cibaria* on *Fusobacterium nucleatum*-induced Interleukin-6 and Interleukin-8 Production in KB Cells. *J. Bacteriol. Virol.* **2011**, *41*, 9–18, doi:10.4167/jbv.2011.41.1.9.
 34. Vieco-saiz, N.; Belguesmia, Y.; Raspoet, R.; Auclair, E.; Padgett, C.; Bailey, C.; Drider, D. Protective Effects of Novel Lactobacillaceae Strains Isolated from Chicken Caeca against Necrotic Enteritis Infection: In Vitro and In Vivo Evidences. *microorganisms* **2022**, *10*.
 35. Silva, M.P.; Tulini, F.L.; Martins, E.; Penning, M.; Fávoro-Trindade, C.S.; Poncelet, D. Comparison of extrusion and co-extrusion encapsulation techniques to protect *Lactobacillus acidophilus* LA3 in simulated gastrointestinal fluids. *LWT-Food Sci. Technol.* **2018**, *89*, 392–399.
 36. Vianna, C.C.R.; Renhe, I.R.T.; Bessa, M.E.; Martins, E.; Stephani, R.; Carvalho, A.F. de; Perrone, Í.T. Microencapsulamento de bactérias probióticas: uma breve revisão. *Res. Soc. Dev.* **2021**, *10*, e242101320814.
 37. Khan, M.M.T.; Pyle, B.H.; Camper, A.K. Specific and rapid enumeration of viable but nonculturable and viable-culturable gram-negative bacteria by using flow cytometry. *Appl. Environ. Microbiol.* **2010**, *76*, 5088–5096.
 38. Monteagudo-Mera, A.; Rastall, R.A.; Gibson, G.R.; Charalampopoulos, D.; Chatzifragkou, A. Adhesion mechanisms mediated by probiotics and prebiotics and their potential impact on human health. *Appl. Microbiol. Biotechnol.* **2019**, *103*, 6463–6472.
 39. Mu, Q.; Tavella, V.J.; Luo, X.M. Role of *Lactobacillus reuteri* in human health and diseases. *Front. Microbiol.* **2018**, *9*, 757.
 40. Yu, Q.; Wang, Z.; Yang, Q. Ability of *Lactobacillus* to inhibit enteric pathogenic bacteria

- adhesion on Caco-2 cells. *World J. Microbiol. Biotechnol.* **2011**, *27*, 881–886.
41. Gagnon, M.; Zihler, B.A.; Chervet, N.; Chassard, C.; Lacroix, C. Comparison of the Caco-2, HT-29 and the mucus-secreting HT29-MTX intestinal cell models to investigate *Salmonella* adhesion and invasion. *J. Microbiol. Methods* **2013**, *94*, 274–279.
 42. Ye, J.; Song, L.; Liu, Y.; Pan, Q.; Zhong, X.; Li, S.; Shang, Y.; Tian, Y.; He, Y.; Chen L, Chen W, Peng Z, W.R. Core 2 Mucin-type Oglycan is related to EPEC and EHEC O157:H7 adherence to human colon carcinoma HT-29 epithelial cells. *Dig. Dis. Sci.* **2015**, *60*, 1977–1990.

GENERAL CONCLUSION

Microorganisms of the genus *Weissella* have a wide range of applications in food products, since they are capable of producing a wide variety of compounds. However, few studies have been performed yet and the industrial application of *Weissella* is not yet a reality.

In addition, this work allowed us to understand the diversity of *Weissella* species found in the different regions producing artisanal cheeses in Brazil, comparing two molecular techniques used for clustering analyzes. It also allowed the understanding of how the isolated strains are distributed in the environments that produce artisanal cheese, besides helping to understand the role they play in the artisanal cheese itself. It also showed that some of them have a favorable technological potential for future uses in the food industry as starter cultures and for the study of cheese defect prevention.

Besides that our work culminated in a possible new bacteriocin from the *Weissella* genus leaving the perspective of further investigation on the producing of this bacteriocin. Also with this work we have a panorama of the possibility of using different strains of *Weissella* for human or animal consumption and as food ingredients. We also notice the relative importance of studies technologies to help the use of this strains as probiotics.

ANNEXES

Annex 1. License to reuse the review article (ref. chapter: 1, part A).

Research, Society and Development, v. 10, n. 5, e8310514557, 2021
(CC BY 4.0) | ISSN 2525-3409 | DOI: <http://dx.doi.org/10.33448/rsd-v10i5.14557>

The *Weissella* genus in the food industry: A review

O gênero *Weissella* na indústria de alimentos: Uma revisão

El género *Weissella* en la industria alimentaria: Una revisión

Received: 04/01/2021 | Reviewed: 04/10/2021 | Accept: 04/15/2021 | Published: 04/28/2021

Research, Society and Development - ISSN 2525-3409



Este obra está licenciado com uma Licença [Creative Commons Atribuição 4.0 Internacional](https://creativecommons.org/licenses/by/4.0/).

CDRR Editors, Avenida Sulim Abramovitch, 100 - Centro, Vargem Grande Paulista - SP, 06730-000
E-mail: rsd.articles@gmail.com | WhatsApp +55 11 98679-6000

Attribution 4.0 International (CC BY 4.0)

This is a human-readable summary of (and not a substitute for) the [license](https://creativecommons.org/licenses/by/4.0/). [Disclaimer](#).

You are free to:

Share — copy and redistribute the material in any medium or format

Adapt — remix, transform, and build upon the material for any purpose, even commercially.

The licensor cannot revoke these freedoms as long as you follow the license terms.



Annex 2. License to reuse the review article (ref. chapter: 1, part B).

Annex 3. License to reuse the research article (ref. chapter: 2).



Biodiversity and technological features of Weissella isolates obtained from Brazilian artisanal cheese-producing regions

Author:
Camila Gonçalves Teixeira, Andressa Fusiéger, Evandro Martins, Rosângela de Freitas, Martina Vakarelova, Luis Augusto Nero, Antonio Fernandes de Carvalho

Publication: LWT - Food Science and Technology

Publisher: Elsevier

Date: July 2021

© 2021 Elsevier Ltd.

Journal Author Rights

Please note that, as the author of this Elsevier article, you retain the right to include it in a thesis or dissertation, provided it is not published commercially. Permission is not required, but please ensure that you reference the journal as the original source. For more information on this and on your other retained rights, please visit: <https://www.elsevier.com/about/our-business/policies/copyright#Author-rights>






BACK **CLOSE WINDOW**

Annex 4. License to reuse the research article (ref. chapter: 3).



Communication

Genomic Analyses of *Weissella cibaria* W25, a Potential Bacteriocin-Producing Strain Isolated from Pasture in Campos das Vertentes, Minas Gerais, Brazil

Camila Gonçalves Teixeira ^{1,2}, Rafaela da Silva Rodrigues ^{1,3}, Ricardo Seiti Yamatogi ³, Anca Lucau-Danila ², Djamel Drider ², Luís Augusto Nero ^{3,*} and Antônio Fernandes de Carvalho ^{1,*}

MDPI Open Access Information and Policy

All articles published by MDPI are made immediately available worldwide under an open access license. This means:

- everyone has free and unlimited access to the full-text of *all* articles published in MDPI journals;
- everyone is free to re-use the published material if proper accreditation/citation of the original publication is given;
- open access publication is supported by the authors' institutes or research funding agencies by payment of a comparatively low **Article Processing Charge (APC)** for accepted articles.

Permissions

No special permission is required to reuse all or part of article published by MDPI, including figures and tables. For articles published under an open access Creative Common CC BY license, any part of the article may be reused without permission provided that the original article is clearly cited. Reuse of an article does not imply endorsement by the authors or MDPI.